



Forest Carbon Partnership Facility (FCPF) Carbon Fund	
ER Monitoring Report (ER-MR)	
ER Program Name and Country:	Zambézia Integrated Landscape Management Program (ZILMP) Republic of Mozambique
Reporting Period covered in this report:	01-01-2021 to 31-12-2022
Number of FCPF ERs:	0
Quantity of ERs allocated to the Uncertainty Buffer	0
Quantity of ERs to allocated to the Pooled Reversal Buffer	0
Quantity of ERs to allocated to the Reversal Pooled Reversal buffer	0
Number of FCPF ERs from enhanced removals through afforestation/ reforestation	0
Date of Submission:	27-11-2025
Version	3.2

WORLD BANK DISCLAIMER

The boundaries, colors, denominations, and other information shown on any map in ER-MR does not imply on the part of the World Bank any legal judgment on the legal status of the territory or the endorsement or acceptance of such boundaries.

The Facility Management Team and the REDD Country Participant shall make this document publicly available, in accordance with the World Bank Access to Information Policy and the FCPF Disclosure Guidance.

TABLE OF CONTENTS

TABLE OF CONTENTS	3
1 IMPLEMENTATION AND OPERATION OF THE ER PROGRAM DURING THE REPORTING PERIOD	7
1.1 Implementation status of the ER Program and changes compared to the ER-PD	7
1.2 Update on major drivers and lessons learned	13
2 SYSTEM FOR MEASUREMENT, MONITORING AND REPORTING EMISSIONS AND REMOVALS OCCURRING WITHIN THE MONITORING PERIOD	14
2.1 Forest Monitoring System	14
2.2 Updates to the monitoring approach.....	20
2.3 Measurement, monitoring and reporting approach.....	21
3 DATA AND PARAMETERS	26
3.1 Fixed Data and Parameters.....	26
3.2 Monitored Data and Parameters	38
4 QUANTIFICATION OF EMISSION REDUCTIONS.....	46
4.1 ER Program Reference level for the Monitoring / Reporting Period covered in this report	46
4.2 Estimation of emissions by sources and removals by sinks included in the ER Program’s scope	46
4.3 Calculation of emission reductions	48
5 UNCERTAINTY OF THE ESTIMATE OF EMISSION REDUCTIONS	48
5.1 Identification, assessment and addressing sources of uncertainty.....	48
5.2 Uncertainty of the estimate of Emission Reductions.....	55
5.3 Sensitivity analysis and identification of areas of improvement of MRV system	58
6 TRANSFER OF TITLE TO ERS	60
6.1 Ability to transfer title	60
6.2 Implementation and operation of Program and Projects Data Management System	62
6.3 Implementation and operation of ER transaction registry	63
6.4 ERs transferred to other entities or other schemes.....	64
7 REVERSALS.....	64
7.1 Occurrence of major events or changes in ER Program circumstances that might have led to the Reversals during the Reporting Period compared to the previous Reporting Period(s)	64
7.2 Quantification of Reversals during the Reporting Period	64

7.3	Quantification of pooled reversal buffer replenishments	67
7.4	Reversal risk assessment	68
8	EMISSION REDUCTIONS AVAILABLE FOR TRANSFER TO THE CARBON FUND	70
	ANNEX 1: INFORMATION ON THE IMPLEMENTATION OF THE SAFEGUARDS PLANS	72
	ANNEX 2: INFORMATION ON THE IMPLEMENTATION OF THE BENEFIT-SHARING PLAN	85
	ANNEX 3: INFORMATION ON THE GENERATION AND/OR ENHANCEMENT OF PRIORITY NON-CARBON BENEFITS	102
	ANNEX 4: CARBON ACCOUNTING - ADDENDUM TO THE ERPD	111
	ANNEX 5: DETAILED DESCRIPTION OF THE APPLICATION OF THE REVERSAL RISK ASSESSMENT TOOL	112
	ANNEX 6: ER MONITORING REPORT (ER-MR) ON THE AREA OUTSIDE THE SCOPE OF ZAMBÉZIA INTEGRATED LANDSCAPE MANAGEMENT PROGRAM (ZILMP)	114
5.1	CARBON POOLS, SOURCES AND SINKS	114
5.1.1	Description of Sources and Sinks selected	114
5.1.2	Description of carbon pools and greenhouse gases selected	114
5.2	REFERENCE LEVEL	114
5.2.1	Reference Period	114
5.2.2	Forest definition used in the construction of the Reference Level	115
5.2.3	Average annual historical emissions over the Reference Period	115
5.2.4	Estimated Reference Level	131
5.3	MONITORING AND REPORTING PERIOD	132
5.3.1	Measurement, monitoring and reporting approach	132
5.3.2	Data and parameters	136
5.4	QUANTIFICATION OF EMISSION REDUCTIONS	148
5.4.1	Reference level for the Monitoring / Reporting Period covered in this report	148
5.4.2	Estimation of emissions by sources and removals by sinks included	148
5.4.3	Calculation of emission reductions	149
5.5	UNCERTAINTY OF THE ESTIMATE OF EMISSION REDUCTIONS	150
	ANNEX 7: DEGRADATION IN THE SCOPE OF ZAMBÉZIA INTEGRATED LANDSCAPE MANAGEMENT PROGRAM (ZILMP)	151
	Calculation of emission reductions	157

LIST OF ACRONYMS

AD	Activity Data
AFS	Agro-Forestry Systems
ANAC	National Administration of Conservation Areas
AQUA	Nacional Agency for Environmental Quality Control
BSP	Benefit Sharing Plan
CATS	Carbon Assets Tracking System
CEAGRE	Centro de Estudos de Agricultura e Gestão de Recursos Naturais
CBNRM	Communities Based in Natural Resources Management
CBOs	Community-Based Organizations
CNG	National Management Committee
DINAF	National Directorate of Forest
DMC	Directorate of Climate Change
EF	Emission Factor
EFF	Planted Forest Grant Scheme
ER	Emission Reduction
ERPA	Emission Reductions Program Agreement
ERDP	Emission Reductions Program Document
ESMP	Environmental and Social Management Plan
FCPF	Forest Carbon Partnership Facility
FDS	Semi-deciduous Forest
FGRM	Feedback and Grievance Redress Mechanism
FSSV	Evergreen Forest
FNDS	National Fund for Sustainable Development
FMS	Forest Monitoring System
FREL	National Forest Reference Emissions Level
FPS	Forest Plantation Scheme
GALs	Gender Action Learn System
GHG	Greenhouse Gases
GNR	Gile National Reserve
LULCC	Land Use and Land Cover Change
GoM	Government of Mozambique
GPM	Good Practice Manual
M&E	Monitoring and Evaluation
MEF	Ministry of Finance
MGS	Matching Grant Scheme
MoU	Memorandum of Understanding
MR	Monitoring Report

MSLF	Multi-stakeholder Landscape Forum
MRV	Monitoring, Reporting and Verification
MTA	Ministry of Land and Environment
NFI	National Forest Inventory
NFMS	National Forest Monitoring System
NTFP	Non-timber Forest products
PDD	Project Document
PDFs	Probability Density Functions
PDIZ	Zambézia Integrated Development Platform
PCUT	Community Land Use Plans
PIU	Project Implementation Unit
PMRV	Participatory Monitoring Report and Verification
PMP	Pest Management Plan
QAQC	Quality Assurance and Quality Control
REDD+	Reducing Emissions from Deforestation and Forest Degradation
SDAE	District Service for Economic Activities
SESA	Social and Environmental Strategic Assessment
SECF	Small Emerging Commercial Farmers
SIS	Safeguards Information System
SOP	Standard Operating Procedure
UNFCCC	National Framework Convention on Climate Change
VCS	Verified Carbon Standard
VVB	Validation and Verification Body
WB	World Bank
WWF	World Wildlife Fund
ZILMP	Zambézia Integrated Landscape Management Program

1 IMPLEMENTATION AND OPERATION OF THE ER PROGRAM DURING THE REPORTING PERIOD

1.1 Implementation status of the ER Program and changes compared to the ER-PD

Unsustainable small-scale agriculture is the major driver of deforestation in Mozambique and within the Zambézia Emission Reduction (ER) Program. Illegal logging and informal charcoal production are the main drivers of forest degradation within the Program geographical area (CEAGRE and Winrock International, 2015). In order to address those drivers, the ER Program is based on a comprehensive approach that recognizes the link between agricultural development, natural resources management and governance. Since the ER Program only accounts for ERs resulting from reduced deforestation, activities focusing on the adoption of sustainable agricultural techniques will be key to its success. Nonetheless, the ER Program has four World Bank (WB) investment projects (the Agriculture and Natural Resources Landscape Management Project ([Sustenta](#)), Mozambique Conservation Areas for Biodiversity and Development Project ([MozBio I](#)), Dedicated Grant Mechanism for Local Communities project ([MozDGM](#)¹) and Forest Investment Project ([MozFIP](#)), and those have a broader approach on land management: their activities extend beyond the agricultural sector. This is actually coherent with the overall scheme of the ER Program, based on an integrated land management approach. Other measures focus on livelihood and income generation through the strengthening of key value chains of cash crops that are not responsible for deforestation, on regularizing land tenure and on community awareness to secure stakeholders' commitment on the long run. Regarding the Emission Reductions Program Document (ERPD) was prepared and this is financing more activities aiming to generate ERs within the ER Program.

Crediting Period and Evidence of Implementation of Activities

The Crediting Period for Zambézia Integrated Landscape Management Program is from January 1st 2018 to December 31st 2024.

According to the FCPF's (Forest Carbon Partnership Facility) *Glossary of Terms* ([link](#)) the Crediting Period Start Date has to comply with several conditions:

- i) it must not be earlier than the date of the first implementation of project activities;
- ii) is justified with objective evidence;
- iii) it cannot be earlier than January 1st 2016;
- iv) cannot fall within the Reference Period; and
- v) the activities must comply with safeguards requirements.

Table 1, which includes on-the-ground activities and enabling environment interventions for 4 of the projects: MozDGM, MozFIP², MozBio I³ and Sustenta⁴. Sustenta, Support to the development of agricultural and forestry value chains of the Project approved in 2021 and 2022 and new business plans for emerging small commercial farmers.

¹ (<https://www.fnds.gov.mz/index.php/en/our-projects/project-list/mozdgm>)

² (<https://www.fnds.gov.mz/index.php/en/our-projects/project-list/MozFIP>)

³ (<https://www.fnds.gov.mz/index.php/en/our-projects/project-list/mozbio>)

⁴ (<https://www.fnds.gov.mz/index.php/en/resources/highlights/131-programa-sustenta-2>)

Compliance with safeguard requirements is described in detail in Annex 1: Information on the implementation of the Safeguards Plans.

Table 1: Specific project activities conducted in the monitoring period in the ER Program Area and the respective annual report which describe the progress on the actions and interventions under the ER Program.

Project	Activity	Evidence
MozFIP	Delimitation of communities	2021 Project Activity Report: https://www.dropbox.com/scl/fi/qe5k9c3auh0t5e8phvog1/Relat-rio-Anual-2021_WB_02.docx?rlkey=ykmklg1e856cws13xdryujsx&dl=0 Website from Service Provider with Results Dashboard (https://sites.google.com/site/verdeazullandscape/dashboard/entrega-de-duats)
	Forest Plantation Scheme (FPS)	Contract with Consultant (https://www.dropbox.com/s/480cc7h8gai0g5y/Moz%20Agroforestry%20Contract%20signed.pdf?dl=0) and (https://www.dropbox.com/s/uowe3m4dourt04k/CONTRACTO%20HORFPEC.Lda.pdf?dl=0) Proof of payment of consultants (https://www.dropbox.com/s/wdioc76k6ilfls/Horfpec%20--%20267.721%2C90.pdf?dl=0) (https://www.dropbox.com/s/mjbdhdcnjq18zb/HORFTEC%20--299.776%2C14.pdf?dl=0)
		Results of 2020/21 from Consultant NIRAS (https://www.dropbox.com/s/rbzj63pp35uixpv/Apresenta%C3%A7%C3%A3o_Missao%20WB_23042021_v002.pdf?dl=0)
		Newsletter 2020/21 (https://www.dropbox.com/s/sa97rb88vfi4ov0/6%20edi%C3%A7%C3%A3o%20do%20Boletim%20Informativo%20do%20EFF%20e%20SAFs%20.pdf?dl=0)
	Benefit Sharing Plan - First payment event	https://www.dropbox.com/sh/wu2h7lcc2eu9198/AAAgTzHzetiUf7VF-n4tKv9a?dl=0
	Communication	https://www.dropbox.com/sh/e64veyfc78gzvxo/AADAFm3DfZgB8WOxJGDDIsXta?dl=0
	Capacity building for the local communities	Report from capacity building for communities in management of natural resources (https://www.dropbox.com/s/6v0ve29d2dp6rc/Relatorio%20Final%20do%20Curso%20de%2)

		<p>0Governanca%20e%20MCRN%20-%20Zambezia.docx.pdf?dl=0</p> <p>Report of capacity building for communities in Nipiode and Anawape https://www.dropbox.com/s/dk4fptc9iksmrw/Relat%C3%B3rio%20Final%20do%20curso%20de%20Fiscaliza%C3%A7%C3%A3o%20e%20Leisla%C3%A7%C3%A3o%20Florestal.pdf?dl=0</p> <p>Finance administration training reports https://drive.google.com/file/d/1UrX1vK5gG7qfPV0pcytBbW3zhGjf68yl/view</p>
Sustenta	Agricultural development (16 SECF- Small emerging commercial farmers)	<p>Proof of payment https://www.dropbox.com/s/bddinbe55om4y40/Paces.pdf?dl=0</p>
	Restoration of degraded areas in the Sustenta Landscape	<p>Proof of payment https://www.dropbox.com/s/gxktce3laksahvy/Factura.pdf?dl=0</p> <p>2019 Project Activity Report https://www.dropbox.com/s/dn9gw1afx7w82zr/Relat%C3%B3rio%20Anual%20de%20Actividades%202019vvvv_25.02.2020.doc?dl=0</p> <p>2020 Project Activity Report https://www.dropbox.com/s/6ve8du56uymy1gx/Relat%C3%B3rio%20Anual%20de%20Actividades%202020.%20PGIARN.versao%20final.1.doc?dl=0.</p>
		<p>2021 Project Activity Report https://www.dropbox.com/s/cr2pox82w10sff1/Relat%C3%B3rio%20Anual%202021_Integrado.%20Versao%2021%20Abril.docx?dl=0</p> <p>2022 Project Activity Report https://www.dropbox.com/scl/fi/lkfwo3h6zqxcomodlguuw/Relat-rio-Anual-2022_Integrado.-Versao-FINAL-2.docx?rlkey=8yiguxbbn15nb1gvhq55d5td4&dl=0</p>
		Support Micro, Small and Medium Enterprises to develop non-timber and non-wood forest products business plans
	Infrastructure	https://www.dropbox.com/sh/vuctt0dik4zynwk/AAAUeWD8NhPbmArHVqj8L7Cba?dl=0
	Maintenance	https://www.dropbox.com/sh/vuctt0dik4zynwk/AAAUeWD8NhPbmArHVqj8L7Cba?dl=0

MOZDGM	World Bank report and annual report	<p>WB report https://www.dropbox.com/s/vwo3ms1dodzu9pz/AM_MozDGM_Nov%202021_FINAL%20%28002%29.pdf?dl=0</p> <p>2021 Project Activity Report https://www.dropbox.com/scl/fi/mryd6omnij9bv1qwc18gi/MozDGM_semiAnnual-Report_Final_Report-2021.pdf?rlkey=jsz4l5phtm8r8rk6fqcbxlxchi&dl=0</p> <p>2022 Project Activity Report https://www.dropbox.com/scl/fi/os2k7hfmgnwgq2jweeoz/Report_MozDGM_July2022.pdf?rlkey=hpp4eeguay3yon409xepdzbdl&dl=0</p>
---------------	-------------------------------------	---

1.1.1 Update on the strategy to mitigate and/or minimize potential Displacement.

The ER Program has done all efforts to minimize displacement of emissions to an area outside the Program boundaries and if present, it will be minimal, as most of the measures proposed to tackle the drivers of deforestation and forest degradation are primarily based on incentives and on the valorization of non-carbon benefits rather than coercive measures that will cause a displacement of drivers of deforestation. Therefore, the expectation is to lower the appeal of deforestation and forest degradation. As part of the strategy, the Monitoring, Report and Verification (MRV) Unit developed a tool to detect annual deforestation for the whole country and currently, the data is accessible through the geospatial platform where deforestation for 2017, 2018, 2019, 2020, 2021 and 2022 in the Districts inside and outside of the ER Program and in other Provinces is displayed (See the link: <https://bit.ly/GeoportalMRVOnline>). Degradation is another component of forest monitoring that the MRV Unit is developing (see the first results of degradation in annex 7). One major driver of deforestation identified during the design of the program was unsustainable small-scale agriculture and two causes of forest degradation identified are illegal logging and charcoal production (these activities were identified during the verification process for 2020 monitoring report). The drivers of deforestation and forest degradation within the ER Program remain the same (see section 1.2). All strategies outlined in the ER-PD are being strictly implemented to avoid displacement and the risk of displacement still assessed and categorized as low for slash and burn agriculture, low for charcoal production and Medium for Illegal logging (Table 2).

Table 2: Updates on strategies being applied to the different drivers of deforestation or degradation to minimize potential displacement.

Small scale agriculture based on “slash and burn” techniques	
Risk of displacement	Low
Progress of the strategy in Place	<p>There is a plan of involving 1500 farmers in technical assistance to adopt sustainable practices of agriculture such as Agroforest systems in about 750 ha. By the end of 2018, 550 farmers from Mulevala District (from 3 communities) were involved in a total area of 250ha. By 2020 the total area of agroforestry systems in the program area increased to a total of 931ha (click here for more information).</p> <p>Community delimitation is the first step towards a sustainable management of natural resources and land. The outputs of community delimitations are Certificates signed by the Provincial Geography and Cadastral Service (SPGC), the community zoning Land Use map, the Community Land Use Plan and the Community development agenda. According to the ERPD plan, the aim is to achieve 322,500 ha of community land supported by land use plans by the end of the crediting period. Sustenta and MozFIP projects delimited a total</p>

	<p>of 187 communities land in Mulevala (48), Gilé (5), Mocubela (27), Gurue (4), Maganja da Costa (81) e Pebane (22) Districts. This number is expected to increase in the following years. This will reduce nomadism thus avoiding displacement. To foster sustainable community management, individual farmers also benefited from Regularization of the Right to Use and Benefit from Land (<i>Regularização do Direito de Uso e Aproveitamento da Terra - R-DUAT</i>) in Mocuba, Mulevala e Gilé 37,671 farmers, in an area of 60,559 ha. More details regarding R-DUATs can be found in Annex 3.</p> <p>The District authorities are incentivizing the adoption of conservation agriculture practices to restore and maintain the soil fertility through public extension services. There are also efforts to promote plantation of cashew trees as part of the agricultural extension package.</p>
Charcoal production	
Risk of displacement	Low
Progress of the strategy in Place	<p>The focus in this component is the training of charcoal makers to incentivize them to use fuel-efficient technology, promote the sustainable management of forests for charcoal production and use of forest logging and sawmill residues. 168 people from communities were trained to adopt improved kilns to produce charcoal in Pebane, Mocubela, Maganja da Costa and Ile. In each community, 500 hectares were identified for sustainable logging to produce charcoal. Four companies from the private sector were also involved in processing sawmill residues to produce charcoal. The use of sustainable charcoal in these communities is also happening and the private sector is in the process of adopting new practices of charcoal production. To ensure the value for money for charcoal production, informal partnership between the private sector and trained communities was established.</p>
Unsustainable forestry practices, including illegal logging	
Risk of displacement	Medium
Progress of the strategy in Place	<p>The project is contributing significantly in strengthening the law enforcement in the forest sector. The Government moved this component from the National Directorate of Forest to the National Agency for Environmental Quality Control (AQUA). The support of the project was concentrated on the preparation of the strategy for law enforcement in forest, and investing on the creation of AQUA Delegation in Zambézia. MozFIP hired an international consultant to support AQUA in the production of the Law enforcement strategy.</p> <p>At the National level, by the recommendation of the last National Forestry Inventory (NFI), the Government of Mozambique (GoM) has recently taken strict actions over the most harvested tree species in Mozambique. For instance, harvesting of <i>Pterocarpus tinctorius</i> (Nkula), <i>Combretum imberbe</i> (Mondzo) and <i>Swartzia madagascariensis</i> (Pau-ferro) was banned as well exportation of <i>Pterocarpus angolensis</i> (Umbila), <i>Millettia stuhlmannii</i> (Jambirre, Panga-Panga), <i>Afzelia quanzensis</i> (Chanfuta) in form of logs was ceased. <i>Swartzia madagascariensis</i> (Pau-ferro) occurs mostly within the Gilé National Park reason why the GoM decided to take such measures as the last NFI indicates that the species' stock has steeply declined over the past 10 years (https://fnds.gov.mz/mrv/index.php/documentos/relatorios/26-inventario-florestal-nacional/file).</p> <p>The GoM conducted a nation-wide audit of licensed areas (forest concessions and simple licenses) to assess the extent to which sustainable forest management practices are improving within the ER Program area and results have shown improvements. This assessment happens every two years since 2016.</p>

	<p>The GoM put in place a new law on timber exports, including log export ban on all native species to incentivize domestic timber processing for adding value to the product whilst also creating more jobs for rural communities</p> <p>A tool of Minimum standards for sustainable management was developed in 2018, to translate into a legal instrument for evaluation of operators' performance to inform any suspension of licenses, with potential for a national certification standard to be developed.</p> <p>In 2018 additional forest inventory plots were sampled in Zambézia, which improved the biomass estimates (link) and was a critical input to the measures taken by the ministry regarding species exploitation and exportation ban.</p>
--	---

1.1.2 Effectiveness of the organizational arrangements and involvement of partner agencies

The success of implementation of an ER program is dependent on the stakeholder engagement. The ER Program has been inclusive on all the decisions regarding interventions on the ground aiming to generate ERs. The major milestones achieved are:

- The creation of a multi-stakeholder landscape forum for sustainable management of natural resources, which is a crucial instrument for stakeholder's consultation and participation in the implementation of the activities within the ER Program. This forum involves different civil society organizations; the Government; Private sector; community organizations and academic institutions. The connection to the platform has been very positive and active.
- a committee for assessing the implementation of forest plantation scheme (Composed by National Directorate of Agriculture and Forestry (DINAS), National Directorate of Forest (DINAF) and Manica Forestry Industries (IFLOMA). This committee has the role of Assess and approve the conformance and eligibility of the proposals to the signature of contracts; Approve payments to beneficiaries of the projects; Monitor progress of the implementation of the scheme. With a committee, performance evaluation of forest plantations was carried out for all beneficiaries of the Forest Plantation Scheme (FPS), showing maintenance rates of the planted area that varies between 70 and 98%. As a result, subsidy payments were made, with the first installment (new beneficiaries) and the second installment (former beneficiaries)⁵.
- Exchange of experience with New Forests Company (NFC) and World Wildlife Fund (WWF) Uganda with the support of NGP (New Generation Plantation) in how to engage the SME in forest plantations. A study visits and technological exchange was held in Uganda, which had as its main theme: "Sustainable Plantations for the Prosperity of Africa and was focused on the challenges and opportunities faced by forest companies in establishing sustainable plantations. This event was co-organized by NFC and WWF-Uganda with the support of the New Generation Plantations (NGP) Platform -. The Travel Report was shared with the World Bank.⁶ Partnership between private sector and communities in small business enterprises (Sustainable charcoal production; non-timber forest products; community concessions, among others).
- Several Memoranda of Understanding (MoU) were signed between Community-Based Organizations (CBOs) and private sector, such as: MoU between Mocuba Honey Company and the associations of Nipiode and Uapé was signed, with a view to making the honey business viable. The National Fund for Sustainable Development (FNDS) promoted a new negotiation process for the partnership to make the

⁵ Source: MozFIP annual progress report 2021

(https://www.dropbox.com/s/8vjilq4fatcw4gx/Relat%C3%B3rio%20Anual%202021_WB_02.docx?dl=0https://www.dropbox.com/s/n49lg01rpouw836/Relat%C3%B3rio%20Anual%202020_FINAL.docx?dl=0)

⁶ Source: MozFIP annual progress report 2018 (<https://www.fnds.gov.mz/index.php/en/our-projects/project-list/MozFIP>)

mushroom business viable with the Divateches-Agri and Miruku consortium, having already signed the MoU. Some negotiations have not been successful, but efforts are still being made to promote more partnerships.

- FNDS signed MoUs with Portucel and Niras in 2020 (more details please click [here](#)). According to these agreements, Portucel and Niras will be responsible for: i) providing seedlings and other inputs at a subsidized price; ii) technical assistance; and iii) Training for extension technicians and beneficiaries (please see 2021 MozFIP reports for more details).
- The signature of the MoU between FNDS and Zambeze University (Unizambeze), to provide technical support for research and development; Supply interns (students) to help communities on the ground to comply with sustainable practices aiming to halt deforestation. The MoU has not yet been signed, but several activities are already underway, such as Unizambeze's involvement in the Forest Plantation Scheme, helping the NIRAS Service Provider and Portucel, during forestry operations.

The major milestones still to be achieved are:

- Insert information from unofficial certificates, Community Land Use Plans (*Planos Comunitários de Uso de Terra* PCUT) and DUATs elaborated throughout the project, in the geospatial platform;
- Promote training and technical demonstrations of planting associated with greater frequency of technical assistance to beneficiaries as well as exchange of experience between various stakeholders;
- Continue aerial monitoring in the areas of Forest Plantation Scheme beneficiaries via drones with the aim of evaluating the progress of tree cover;
- Forest monitoring by drone in the beneficiaries' area to evaluate the progress of the tree plantation;
- Disseminate the participatory MRV in others communities, in order to involve all actors, such as communities, civil society, private and public sector, in the assessment of deforestation and forest degradation. Therefore, capacity must be built at the level of provinces, districts and communities and training of community technicians.

1.1.3 Updates on the Assumptions in the Financial Plan and Any Changes in Circumstances Affecting the ER Program Implementation

The financial plan of the Zambézia Integrated Landscape Management Program (ZILMP) under the REDD+ framework was developed based on stable assumptions, including expected funding sources and cost estimates. During the reporting period, no changes occurred in the main financial assumptions, with funding continuing to come exclusively from the the World Bank

1.2 Update on major drivers and lessons learned

Unsustainable small-scale agriculture still by far, the first driver of deforestation in the ER Program area. The monitoring data produced by the MRV unit in [2018](#) and [2019-2020](#) show that agriculture is the main drive of deforestation. Other drivers such as forest activities for timber and charcoal could not detected directly as drivers. The solution is to improve the tool to detect the forest degradation which combines with updated high-resolution imagery or/and field survey.

The charcoal production process is a result of agriculture expansion and small agriculturalists maximizing value from the land clearing process. As evidence, during the site visit of monitoring, the ZILMP team noted that expanding subsistence agriculture is the primary driver of deforestation. The ZILMP team had the opportunity to interview numerous small formers during the site visit and is reasonably assured that expanding subsistence agriculture remains the primary driver of deforestation. On other hands in June of 2021 started a study that aims to analyze the driver, economic and cultural factors in deforestation and forest degradation. This study conclude that the agriculture is the main driver of deforestation (see full report [here](#)). At the moment (December 2023), the MRV and BSP team are carrying out a survey of PMRV data in several districts of Zambézia to better understand the reason for the increase in deforestation in the years prior to this monitoring period

Unsustainable timber exploitation poses a medium risk for potential displacement of the activity to the districts outside of the ER Program because law enforcement was intensified. However, such intensification had taken place throughout the country also, thus minimizing this potential risk. No harmful activities were prohibited inside of the ER Program as part of the strategies to minimize potential displacement; conversely, improvements on practices are based on incentives for agricultural intensification and settlement within the ER Program area through systematic land use delimitation and titling for individuals and communities. The integration of sustainable practices in forestry, agriculture and land use in the program area with involvement of different stakeholders using the participatory approach generated ERs for this monitoring/reporting period.

The risk of displacement is low as other Government initiatives are taking place on the other districts outside of the ER Program. For more information on the drivers of deforestation and forest degradation within the ER Program, kindly refer to the [Mozambique's ERPD](#). To sustain the generation of ERs in the program area and minimize the risk of displacement MozFIP will continue to monitor the dynamic of emissions from deforestation and forest degradation and invest in sustainable practices in agriculture, forestry and land.

1.3 Methodological deviations

Intentionally left blank

No correction to be done on material or non-material error or misstatement impacting the estimation

2 SYSTEM FOR MEASUREMENT, MONITORING AND REPORTING EMISSIONS AND REMOVALS OCCURRING WITHIN THE MONITORING PERIOD

2.1 Forest Monitoring System

Organizational Structure, Responsibilities, and Competencies

Mozambique has not formalized its national forest monitoring system (NFMS). There is a work in progress between the DINAF and FNDS and other relevant stakeholders to formalize the NFMS. This process started in 2019, with the establishment of the NFMS Task Force, responsible for designing, developing and operationalizing the NFMS. Technical officers of DINAF and FNDS (MRV unit) were appointed as its members.

The NFMS Working Group, as a group of stakeholders related to the NFMS with its role to provide related information, inputs and advice to the development and operationalization of the NFMS. The initial members included DINAF, National Directorate of Environment (DINAB), National Directorate of Land (DINAT), FNDS, Eduardo Mondlane University (UEM), Institute of Agricultural Research of Mozambique (IIAM), FAO (Food and Agriculture Organization of the United Nations), World Bank, International Union for Conservation of Nature (IUCN) and Federation of Timber Operators (AMOMA). However, the Working Group is, by its nature, an open forum which the members can change flexibly depending on the needs and interests.

The first version of the NFMS document was finalized in 2021 ([link](#)) and defines the NFMS as a system which enables accountable reporting of REDD+ results; monitoring the implementation and effectiveness of Policies and Measures (PaMs) for sustainable forest management, which include national and international purposes and beyond REDD+; and builds on robust IT system to support data management and transparency. The following principles are stated in the document as the basis of the NFMS:

- The NFMS shall be designed and operationalized under the full ownership of Ministry of Land and Environment (MTA), and in collaboration with relevant stakeholders;
- The NFMS should be target-driven, oriented towards specified sub-national, national and international objectives;
- The NFMS shall build on existing system as far as practical;
- The NFMS shall be developed through step-wise improvement, take into consideration the national circumstances, reflect the phased approach for the implementation of REDD+ activities, and sustainable in the long-run. The development shall be realistically feasible within the available time, financial and human resources; and
- The NFMS shall meet the international requirement under REDD+, and as appropriate, apply international and national good practices.

Selection and Management of GHG-related Data and Information

The current monitoring system has three sub-systems:

- Satellite and land monitoring system
- National forest inventory
- National Greenhouse Gases (GHG) inventory

Satellite and land monitoring system

The satellite and monitoring system is a sub-system within the NFMS that produces the activity data. The MRV Unit within FNDS is responsible for this system. It specifically generates the information on the number of hectares of deforestation within a given geographic area. This system produced information of deforestation that was used to produce the ER Program's RL and the National Forest Reference Emissions Level (FREL). This information was also used to generate historical deforestation statistics by Provinces, districts (link <https://fnds.gov.mz/mrv/index.php/documentos/estudos/15-anuario-ambiental-para-instituto-nacional-de-estatistica-ine/file>), conservation areas and Zambézia Integrated Landscape Management Program (ZILMP) using a systematic stratified sampling. With new tool to detect deforestation developed, it was possible to produce annual deforestation maps for 2017, 2018, 2019, 2020, 2021 and 2022 as shown through the link <https://bit.ly/GeoportalMRVOnline> for the whole country and the area estimates for Zambézia Province which are based on sampling approach. The MRV unit from FNDS is responsible to produce the activity data for the ZILMP as well as for the country, as it has gained experience and expertise from training provided by FCPF finance.

The process of generating activity data comprises five steps (Figure 5); they are *response design, map production, sampling design, data collection and analysis*. These steps mainly define the criteria for classification, produces a change map and area estimates.

To ensure a good quality of data the team developed and implemented Quality Assurance/Quality Control (QA/QC) processes in all production processes including the development of Standard Operating Procedures (SOPs). This ensures a high standard of quality of the data produced. To guarantee the replication of processes, the MRV unit developed a Portuguese version guideline to produce activity data, accessed through the link <https://www.fnds.gov.mz/mrv/index.php/documentos/guioes/62-protocolo-de-monitoria-e-estimativa-de-emissoes-por-desmatamento-vmar2023/file>. Data collection is conducted by a core team of professional interpreters who work permanently for FNDS and who have received adequate training in the implementation of the SOPs.

To disseminate the use of activity data to communities and other stakeholders to monitor deforestation, the MRV unit started in December of 2020 to set up participatory MRV (PMRV) systems as described below ([Forest Monitoring System under the ZILMP](#)).

National forest Inventory

The national forest inventory is the second sub-system within the NFMS, which produces the emission factors. They give the tonnage of carbon stored per unit hectare of forest. The tonnage of carbon per hectare varies from one type of forest to another. Mozambique has conducted four national forest inventories and the updating of NFI is carried out every 10 years. The last inventory in 2016-17 produced the emission factors used for the FREL submitted to the United Nations Framework Convention on Climate Change (UNFCCC) in 2018 (report may be accessed in the link: https://redd.unfccc.int/files/moz_frel_report_final.v03_03102018.pdf). In order to have more accurate estimates for the ZILMP, the plots located in Zambézia Province were used to generate specific Emission Factors for ZILMP. The methods to generate the emissions factors for ZILMP are described in the Zambézia forest inventory report: <https://fnds.gov.mz/mrv/index.php/documentos/relatorios/41-relatorio-de-inventario-florestal-da-zambezia-actualizado/file>.

The process used to produce the emission factors followed these steps: Response design, Sampling design, Data collection and Data analysis (Details in Figure 5). The entity responsible for the National Forest Inventory is DINAF. The NFI report (<https://fnds.gov.mz/mrv/index.php/documentos/relatorios/26-inventario-florestal-nacional/file>) was produced by FNDS and DINAF. The data collection involved the IIAM, the Faculty of Agronomy and Forest Engineering (FAEF), the Department of Biological Sciences and Provincial Forest Services. The estimation of emissions also relies on the allometric equations that have been developed by Masters and PhD

students and research projects from FAEF and the Department of Biological Sciences (DCB) of the UEM, as well as peer-reviewed publications.

To ensure the quality of the data collected, the team followed QAQC procedures defined by DINAF. To maintain the processes of the national forest inventory, the MRV unit developed a practical field manual for training teams in data collection that can be accessed on the link <https://fnds.gov.mz/mrv/index.php/documentos/guioes/21-manual-do-inventario-florestal/file>.

The Permanent Sample plots are another component of the National Forest Monitoring System that will improve the estimation of emissions factors and IIAM leads the process. Currently, under the MozFIP project, a joint group of institutions that involves IIAM, FNDS, UEM and DINAF are establishing the network of Permanent Sampling plots across the country, including Zambézia province.

National GHG inventory

The National GHG inventory for the purpose of REDD+ combines the Activity data and the emission factors (Figure 5) to estimate the annual emissions and the FREL.

At the national level, the recent experience of GHGs inventory was with the submission of the FRELs to the UNFCCC (https://redd.unfccc.int/files/moz_frel_report_final.v03_03102018.pdf). The National Directorate of Climate Change is responsible for the communication of GHG emissions of Mozambique, as the focal point for climate change with the UNFCCC. The National Directorate of Climate Change coordinates with DINAF and FNDS on the production of such information.

At the subnational level, the MRV unit from FNDS is currently responsible for the generation of all information related to emissions from deforestation for the ZILMP program. The MRV Unit is also generating estimates of emissions from deforestation at national, Provincial and District level. To maintain the quality standards in the production of emissions estimates from deforestation, the MRV unit has developed SOPs on how to produce the estimates.

Major institutional changes in institutional arrangements since the Approval of ERPD were: (1) Changes in the Ministries; (2) Change in the institutions. Before the approval of the ERPD, FNDS, DINAF, and the National Directorate of Environment were under the Ministry of Land, Environment and Rural Development (MITADER). IIAM was under the Ministry of Agriculture and Food Security (MASA); after the elections in 2019, the new Government was formed, and the result was the extinction of MITADER with the creation of Ministry of Land and Environment (MTA), the extinction of MASA with the creation of the Ministry of Agriculture and Rural Development (MADER). As a result, FNDS and IIAM were moved to MADER, while the National Directorate of Environment and DINAF moved to MTA. The climate change component of National Directorate of Environment was moved to a new Directorate, the National Directorate of Climate Change. This new setting is important as FNDS and DINAF now interact with the national Directorate of Climate Change on issues related to Reporting. Despite these changes on the institutional arrangements and lack of a formal institutional arrangement, the components of the Forest Monitoring System can deliver the function of producing the emissions from deforestation at all levels.

Forest Monitoring System under the ZILMP

The forest monitoring system (FMS) under the ZILMP is simpler in terms of processes and entities as it relies on the first and second system above and it is fully operated by the MRV unit within FNDS with collaboration of DINAF. Therefore, the system uses the standard technical procedures of the NFMS as required by Criterion 15 of the MF.

In December of 2020 (see section 1.2), the MRV Unit tested the introduction of participatory MRV (PMRV) for annual monitoring of deforestation under the ZILMP, which is part of the recommendation of civil society, decision makers and the scientific community in the measurement, reporting and verification (MRV) of stocks of carbon with the participation of the local community. In phase one, the PMRV was tested in twelve communities across the districts of Alto Molocué, Mocuba, Mulevala and Gilé, where three communities were selected per district. The results of the PMRV test can be found on MRV website (PRMV page - <https://bit.ly/pmrvfndsredd>).

The aim of the PMRV activities was to involve communities in deforestation reporting activities (confirming deforestation cases and reporting new cases in near-real time using GIS tools such as Survey123 for ArcGIS, ArcGIS Field Maps and ArcGIS Collector). In 2021 the field activity of PMRV was carried out in communities such as Munhiba in Mocuba district, Dindini and Sacane – Pebane district, Muapila and Cannaua – Mocubela district, Soares, Vacha, Mutchana, Mutchiua and Nehita in Alto Molocue district, and Namigonha, Vassele, Malema-Serra, and Namurua in Gilé district. This activity involved staff from government institutions, academia and civil society (BIOFUND, AQUA, ANAC, Unizambeze, Unilurio, Network for Community Management of Natural Resources (Regecom), DINAF and FNDS technicians to present and demonstrate the tool and its potential. For more details, please find the PMRV report [here](#).

Information on the ZILMP can be found both on the FCPF website (<https://www.forestcarbonpartnership.org/country/mozambique>) and the MRV Unit website (<https://www.fnds.gov.mz/mrv/>). The ERPD is available online on the FCPF website (https://www.forestcarbonpartnership.org/system/files/documents/Mozambique_Revised%20ERPD_16April2018_CLEAN.pdf). The latest version of 2018 (first ER report) and 2019/2020 (second ER report) Monitoring Reports are also available online, on the FCPF website (<https://www.forestcarbonpartnership.org/country/mozambique>) and on the MRV Unit website (<https://www.fnds.gov.mz/mrv/index.php/documentos/relatorios>).

The organogram of the MRV Unit responsible for the ZILMP monitoring is described in

Figure 1. The MRV Unit was created in 2016, with a coordinator (Aristides Muhate) and 4 technicians (Alismo Herculano, Credêncio Maúnze, Délfio Mapsanganhe and Hercilo Odorico). Towards the end of 2016 a fifth element was added to the team (Muri Soares). In 2019 the unit added 3 new elements (Alex Boma, Orlando Macave and Sérgio João). In 2022, the MRV Unit added 3 new technicians to respond the demand of PMRV (Edna Munjovo, Felício Guelume and Sadamo Ussene). Therefore, various efforts have been made in terms of personnel and resources in order to maintain the capacity of the MRV system to monitor and report emissions and emission reductions. The production of the various SOPs has contributed to the knowledge management of the MRV Unit. In addition, there is no task performed by only one person, which increases redundancy. The MRV Unit recognizes that there is a need for continuous improvement of its knowledge management process, to ensure that all activities are standardized and documented. The organizational structure for the Activity data (reference and annual) and NFI is described in Figure 2, Figure 3 and Figure 4.




Figure 1: Organogram of MRV Unit responsible for ZILMP monitoring.




Figure 2: Organizational structure for Activity Data of Reference Level.




Figure 3: Organizational structure for National Forest Inventory.




Figure 4: Organizational structure for Annual Activity Data.

Systems and Processes Ensuring Data Accuracy

To maintain data accuracy, the monitoring system incorporates:

- **Quality Assurance/Quality Control (QA/QC) protocols** for all data collection and processing steps⁷.
- **Standard Operating Procedures (SOPs)** to standardize methodologies and ensure consistency⁸.
- **Training and Capacity Building** to enhance the skills of personnel responsible for data collection and analysis.
- **Independent Review and Validation** by international organizations, such as Aster Global and UNFCCC

Systems and processes that support the Forest Monitoring System, including Standard Operating Procedures and QA/QC procedures;

The developed SOPs are:

- Map production – SOP0
- Sampling Design – SOP1
- Response Design – SOP2
- Data Collection – SOP3
- Sample-based Area Estimation Analysis – SOP4

FNDS also has detailed QAQC procedures for the collection of reference data for the sample-based area estimation, which is described in the Standard Operating Procedures for Area Estimation document ([link](#)), which contains the above SOPs.

Role of Communities in the Forest Monitoring System

The Participatory MRV (PMRV) initiative, launched in December 2020 under the ZILMP, involves local communities in monitoring deforestation. Community members are trained to:

- Use GIS tools such as Survey123 and ArcGIS for deforestation detection.
- Confirm and report deforestation cases in near-real time.
- Provide ground-truth data to validate satellite-based observations.

2.2 Updates to the monitoring approach

"Intentionally left blank"

⁷ <https://www.fnfs.gov.mz/mrv/index.php/documentos/guioes/62-protocolo-de-monitoria-e-estimativa-de-emissoes-por-desmatamento-vmar2023/file>

⁸

https://www.dropbox.com/scl/fo/tn3qh2zy4yycl0avfrx45/AlqGjJauGtov_Ggg764pUCI?rlkey=eZ94z07c9ihlf6e2tdd7e0mrk&dl=0

2.3 Measurement, monitoring and reporting approach

2.3.1 Line Diagram

The Figure 5 illustrates the emissions reductions calculation workflow during the Monitoring Period. It is important to note that as part of the ZILMP, all this workflow including the phase of reported is implemented by the MRV unit within FNDS.




Figure 5: Emissions reductions calculation workflow.

Step-by-step description of monitoring and measurement approach presented in the Line Diagram above.
More details find [here](#).

Category	Monitoring parameters	Methodological Approach
Activity Data	Deforestation Monitoring	<p>The area of interest (AOI) is delineated using project boundaries. Satellite imagery (Sentinel-2 or Landsat) is selected for reference and current periods, preprocessed to remove clouds and shadows, and vegetation indices (NDVI, EVI, SAVI, NDWI, NBR) are computed. Land cover is classified using Random Forest, followed by post-processing to remove noise and apply spatial filters. Stratified sampling is designed based on cleaned maps, and sample plots are interpreted to produce area estimates per class using the equation below. Accuracy is assessed with user's, producer's, and overall metrics.</p> $n = \frac{(\sum W_i S_i)^2}{[S(\hat{O})]^2 + (\frac{1}{N}) \sum W_i S_i} \approx \left(\frac{\sum W_i S_i}{S(\hat{O})} \right)^2$
Emission Factors	Forest Carbon Stock Assessment	<p>Above- and below-ground biomass is measured using national forest inventory (NFI) plots. Biomass is converted to carbon using the carbon fraction CF=0.47, following IPCC guidelines: C=B×CF. Emission factors are stratified by forest type to capture spatial variability.</p>
Emission Estimation	Calculate Emissions	<p>Carbon loss for each land cover transition is computed as</p> $\Delta C_B = \sum_{j,i} (B_{Before,j} - B_{After,i}) \times CF$ <p>This is converted to CO₂-equivalent emissions: Emissions = $\frac{44}{12} \times A(j, i)$. Emissions are aggregated over all strata and regions. Uncertainty is quantified using sample variance or Monte Carlo simulation to derive 95% confidence intervals.</p>
Emissions Reductions	Avoided Deforestation	<p>The reduction in emissions is estimated by comparing gross annual emissions to the Forest Reference Emission Level (FREL): ER = FREL–Actual Emissions. This provides tCO₂e/year of avoided emissions, forming the basis for reporting and verification.</p>

2.3.2 Calculation

Emission reduction calculation

$$ER_{ERP,t} = RL_t - GHG_t \quad \text{Equation 1}$$

Where:

ER_{ERP}	=	Emission Reductions under the ER Program in year t ; $tCO_2e \cdot year^{-1}$.
RL_{RP}	=	Gross emissions of the RL from deforestation over the Reference Period; $tCO_2e \cdot year^{-1}$.
GHG_t	=	Monitored gross emissions from deforestation at year t ; $tCO_2e \cdot year^{-1}$;
T	=	Number of years during the monitoring period; <i>dimensionless</i> .

Reference Level (RL_t)

Gross emissions of the RL from deforestation over the Reference Period (RL_{RP}) are estimated as the sum of annual change in total biomass carbon stocks (ΔC_{B_t}) during the reference period.

$$RL_{RP} = \frac{\sum_t^{RP} \Delta C_{B_t}}{RP} \quad \text{Equation 2}$$

Where:

ΔC_{B_t} = Annual change in total biomass carbon stocks at year t ; $tC \cdot year^{-1}$;
 RP = Reference period; years.

Following the 2006 IPCC Guidelines, the annual change in total biomass carbon stocks forest land converted to other land-use category (ΔC_{B_t}) would be estimated through the following equation:

$$\Delta C_{B_t} = \Delta C_G + \Delta C_{CONVERSION} - \Delta C_L \quad \text{Equation 3}$$

Where:

ΔC_{B_t} Annual change of total biomass carbon stocks during the period, in tC per year;
 ΔC_G Annual increase in carbon stocks in biomass due to growth on land converted to another land-use category, in tC per hectare and year;
 $\Delta C_{CONVERSION}$ Initial change in carbon stocks in biomass on land converted to other land-use category, in tC per hectare and year; and
 ΔC_L Annual decrease in biomass carbon stocks due to losses from harvesting, fuel wood gathering and disturbances on land converted to other land-use category, in tC per hectare and year.

Following the recommendations set in chapter 2.2.1 of the GFOI Methods Guidance Document for applying IPCC Guidelines and guidance in the context of REDD⁹, the above equation will be simplified and it will be assumed that:

- The annual change in total biomass carbon stocks (ΔC_B) is equal to the initial change in carbon stocks ($\Delta C_{CONVERSION}$);

Considering equation 2.16 of the 2006 IPCC GL for estimating ($\Delta C_{CONVERSION}$) the change of biomass carbon stocks could be expressed with the following equation:

$$\Delta C_{B_t} = \sum_{j,i} (B_{Before,j} - B_{After,i}) \times CF \times \frac{44}{12} \times A(j, i)_{RP} \quad \text{Equation 4}$$

Where:

$A(j, i)_{RP}$ Area converted/transited from forest type j to non-forest type i during the Reference Period, in hectares per year. In this case, three forest land conversions are possible:

- (Semi-)deciduous forest to Non-forest type i ;
- (Semi-)evergreen forest to Non-forest type i ; and
- Mangrove forest to Non-forest type i .

Five types of non-forest land are considered:

- Cropland (C);
- Grassland (P);
- Wetland (A);
- Settlement (U); and
- Other lands (O).

$B_{Before,j}$ Total biomass of forest type j before conversion/transition, in tons of dry matter per ha. This is equal to the sum of aboveground ($AGB_{Before,j}$) and belowground biomass ($BGB_{Before,j}$) and it is defined for each forest type.

$B_{After,i}$ Total biomass of non-forest type i after conversion, in tons dry matter per ha. This is equal to the sum of aboveground ($AGB_{After,i}$) and belowground biomass ($BGB_{After,i}$) and it is defined for each of the five non-forest IPCC Land Use categories.

⁹ <https://www.reddcompass.org/mgd>

This parameter was technically corrected so as to replace the estimates sourced from research by estimates given by the 2006 IPCC Guidelines.

- CF* Carbon fraction of dry matter in tC per ton dry matter. The value used is:
- **0.47** is the default for (sub)tropical forest as per IPCC AFOLU guidelines 2006, Table 4.3.
- 44/12 Conversion of C to CO₂

Monitored emissions (GHG_t)

Annual gross GHG emissions over the monitoring period in the Accounting Area (GHG_t) are estimated as the sum of annual change in total biomass carbon stocks (ΔC_{B_t}).

$$GHG_t = \frac{\sum_t^T \Delta C_{B_t}}{T} \quad \text{Equation 5}$$

Where:

- ΔC_{B_t} = Annual change in total biomass carbon stocks at year t ; $tC \cdot year^{-1}$
- T = Number of years during the monitoring period; *dimensionless*.

Changes in total biomass carbon stocks

Following the 2006 IPCC Guidelines, the annual change in total biomass carbon stocks forest land converted to other land-use category (ΔC_B) would be estimated through **Equation 3** above. Making the same assumptions as described above for the RL the change of biomass carbon stocks could be expressed with the following equation:

$$\Delta C_B = \sum_{j,i} (B_{Before,j} - B_{After,i}) \times CF \times \frac{44}{12} \times A(j,i)_{MP} \quad \text{Equation 6}$$

Where:

$A(j,i)_{MP}$ Area converted/transited from forest type j to non-forest type i during the Monitoring Period, in hectare per year. In this case, three forest land conversions are possible:

- (Semi-)deciduous forest to Non-forest type i ;
- (Semi-)evergreen forest to Non-forest type i ; and
- Mangrove forest to Non-forest type i .

Five types of non-forest land are considered:

- Cropland (C);
- Grassland (P);
- Wetland (A);
- Settlement (U); and
- Other lands (O).

These parameters may be found in [Section 3.2](#).

$B_{Before,j}$ Total biomass of forest type j before conversion/transition, in tons of dry matter per ha. This is equal to the sum of aboveground ($AGB_{Before,j}$) and belowground biomass ($BGB_{Before,j}$) and it is defined for each forest type.

This was defined ex-ante and is described in [Section 3.1](#).

$B_{After,i}$ Total biomass of non-forest type i after conversion, in *tons dry matter per ha*. This is equal to the sum of aboveground ($AGB_{After,i}$) and belowground biomass ($BGB_{After,i}$) and it is defined for each of the five non-forest IPCC Land Use categories.

This was defined ex-ante and is described in [Section 3.1](#).

CF Carbon fraction of dry matter in tC per ton dry matter. The value used is:

- **0.47** is the default for (sub)tropical forest as per IPCC AFOLU guidelines 2006, Table 4.3.

44/12 Conversion of C to CO₂

3 DATA AND PARAMETERS

3.1 Fixed Data and Parameters

Parameter:	AGB _{before,j}
Description:	Aboveground biomass of forest type <i>j</i> before conversion,
Data unit:	tons of dry matter per ha
Source of data or description of the method for developing the data including the spatial level of the data (local, regional, national, international):	<p>The data used for the present document are Tier 2 (country specific data or country level estimates or locally derived estimates) and they were sourced from the NFI (for deciduous and evergreen forests) or from a regional study conducted in the Zambezi River Delta for Mangrove forests.</p> <p>For semi-deciduous and evergreen forest, the data are from the Zambézia Forest Inventory. It includes data that was collected in Zambézia province during the NFI, in 2017 and 2018. Although the inventory covers the whole Zambézia province (ER ZILMP program and outside). This is still representative of the forests located in the ZILMP as forests across the province are homogenous (floristic and structural composition). Moreover, the higher sample size of the forest inventory covering the whole province will enable more precise estimates for emission factors.</p> <p><i>i. Sampling design</i></p> <p>Carbon stocks before conversion for deciduous and evergreen forests were estimated using data from the National Forest Inventory sample units that were located in Zambézia province. The sample units for surveying carbon stocks were allocated using restricted stratified random sampling, using 4 * 4 km systematic grid superimposed on the agro-ecological zoning map, and stratified among the 12 forest types. Was considered as the strata, the semi-deciduous forest “open and closed”, Miombo forest “open and closed”, semi-evergreen forest “open and closed”, semi-evergreen Mountain forest “open and closed”, Mopane forest “open and closed”, and Mecrusse forest “open and closed”, of which only the first eight types occur in Zambézia province.</p> <p>The total number of sample units was determined using the optimal allocation (assuming a maximum error of 10% for the total volume, and 5% of confidence level). Proportional allocation was used to determine the number of sample units per stratum (Husch, Beers, and Kershaw 2003). For Zambézia province, 128 clusters (512 plots) were distributed between the eight (8) forest types. The cluster was used as a sampling unit, and each cluster has 4 plots of 0.1 ha (20 * 50 m), where each plot was divided into 4 sub-plots of 0.025 ha (10 * 25 m) (Figure 6).</p>




Figure 6: Design of each cluster used in the National Forest Inventory.

For estimating emission factors, the eight strata were aggregated into 2 (Semi-deciduous forest and semi-evergreen forest), and the similarity of the strata was used for the aggregation. The aggregation was done with the purpose of harmonizing the forest strata of the activity data with the emission factor data. Below the aggregation table.

Allocation stratum	EF Strata for MR
semi-deciduous open forest	semi-deciduous forest
semi-deciduous closed forest	
miombo open forest	
miombo closed forest	
semi-evergreen Mountain open forest	semi-evergreen forest
semi-evergreen Mountain closed forest	
semi-evergreen open forest	
semi-evergreen closed forest	

ii. Data collection

The plots were used for data collection of adult trees (diameter at breast height (dbh) $\geq 10\text{cm}$), and the subplots "A" were used for data collection of established regeneration trees ($10\text{cm} > \text{dbh} \geq 5\text{cm}$), which were included in the calculation of the carbon stocks. Data collected in the plots and subplots included tree information (dbh, scientific name, total and commercial height, stem quality), soil, forest type (this information was used to validate the information from agro-ecological zoning map), and other important information. Tree data were used to estimate above ground biomass (AGB) and below ground biomass (BGB).

The NFI did not cover Mangrove forests, so, data from the [literature](#) was used. For other strata, data from literature were also used.

Details of data collection can be found at <https://www.fnds.gov.mz/mrv/index.php/documentos/guioes/35-directrizes-do-inventario-florestal-nacional/file>.

iii. Prediction at plot level

Above ground biomass (AGB) and below ground biomass (BGB) were estimated using a series of allometric equations adjusted for ecosystems or tree species similar to those in the Zambézia province (Table 3), and this equation was applied at tree level.

The use of the equations meant, applying allometric equations of the specific species (*Millettia stuhlmannii* taub., *Pterocarpus angolensis* DC., *Afzelia quanzensis* Welw.) in all trees of these species to estimate AGB, regardless of forest types. The allometric equation of the semi-deciduous forest was applied for all trees of this forest type (except the species mentioned above), as well as in all trees of the species *Brachystegia spiciformis* Benth., and *Julbernardia globiflora* (Benth.) Troupin to estimate AGB and BGB, because they were the main species used to adjust this equation in this forest type. The equations of the semi-evergreen forest were applied in all remaining trees of this forest type to estimate AGB; and apply the semi-deciduous forest equation in all trees to estimate the BGB in this forest type (including species mentioned above in other forest type), and apply factor 0.275 (shoot ratio) to estimate the BGB of the semi-evergreen forest.

Table 3: List of allometric equations used to estimate above and below biomass

Forest Type	Forest type or species	Above-ground biomass (AGB) [kg]	Below-ground biomass (BGB) [kg]
Semi-deciduous forest	Semi-deciduous forest (open and closed)	$\hat{Y} = 0.0763 * DAP^{2.2046} * H^{0.4918}$	$\hat{Y} = 0.1766 * DAP^{1.7844} * H^{0.3434}$
		Author: Mugasha <i>et al.</i> (2013)	Author: Mugasha <i>et al.</i> (2013)
	<i>Millettia stuhlmannii</i> taub.	$\hat{Y} = 5.7332 * DAP^{1.4567}$	$\hat{Y} = 0.1766 * DAP^{1.7844} * H^{0.3434}$
		Author: Mate <i>et al.</i> (2014)	Author: Mugasha <i>et al.</i> (2013)
	<i>Pterocarpus angolensis</i> DC.	$\hat{Y} = 0.2201 * DAP^{2.1574}$	$\hat{Y} = 0.1766 * DAP^{1.7844} * H^{0.3434}$
		Author: Mate <i>et al.</i> (2014)	Author: Mugasha <i>et al.</i> (2013)
<i>Afzelia quanzensis</i> Welw.	$\hat{Y} = 3.1256 * DAP^{1.5833}$	$\hat{Y} = 0.1766 * DAP^{1.7844} * H^{0.3434}$	
	Author: Mate <i>et al.</i> (2014)	Author: Mugasha <i>et al.</i> (2013)	
Evergreen forest	Evergreen forest (open and closed)	$\hat{Y} = \exp(-2.289 + 2.649 \ln(DAP) - 0.021(\ln(DAP))^2)$	$\hat{Y} = AGB * R/S; \quad R/S = 0.275$
		Author: IPCC (2003)	Author: Mokany <i>et al.</i> (2006)
		$\hat{Y} = 0.0613 * DAP^{2.7133}$	$\hat{Y} = AGB * R/S; \quad R/S = 0.275$

	Evergreen Mountain forest (open and closed)	Author: Lisboa <i>et al.</i> (2018)	Author: Mokany <i>et al.</i> (2006)							
	<i>Millettia stuhlmannii</i> taub.	$\hat{Y} = 5.7332 * DAP^{1.4567}$	$\hat{Y} = 0.1766 * DAP^{1.7844} * H^{0.3434}$							
		Author: Mate <i>et al.</i> (2014)	Author: Mugasha <i>et al.</i> (2013)							
	<i>Pterocarpus angolensis</i> DC.	$\hat{Y} = 0.2201 * DAP^{2.1574}$	$\hat{Y} = 0.1766 * DAP^{1.7844} * H^{0.3434}$							
		Author: Mate <i>et al.</i> (2014)	Author: Mugasha <i>et al.</i> (2013)							
	<i>Azelia quanzensis</i> Welw.	$\hat{Y} = 3.1256 * DAP^{1.5833}$	$\hat{Y} = 0.1766 * DAP^{1.7844} * H^{0.3434}$							
		Author: Mate <i>et al.</i> (2014)	Author: Mugasha <i>et al.</i> (2013)							
	<p>iv. Estimation</p> <p>The estimation of mean and their respective uncertainties (standard error, sampling error, and confidence interval) for the variables biomass, carbon and carbon dioxide equivalent (above and below ground) for the two strata (semi-deciduous forest and semi-evergreen forest), were done using the forest inventory data analysis approach proposed by Bechtold & Patterson (2005) chapter 4 of the book “The Enhanced Forest Inventory and Analysis Program-National Sampling Design and Estimation Procedures”. Details of this methodology are described in Zambézia inventory report, available at https://fnfs.gov.mz/mrv/index.php/documentos/relatorios/41-relatorio-de-inventario-florestal-da-zambezia-actualizado/file.</p> <p>For mangrove forests, data are secondary, extracted from existing literature. Stringer <i>et al.</i> (2015)¹⁰ made an inventory on this ecosystem in the Zambezi delta in Mozambique; we can easily assume that carbon stocks are comparable to those of mangroves in Zambézia province. They divided mangroves into 5 strata and estimated carbon stocks in above and belowground biomass. Since we do not have information on these specific strata for ZILMP, the mean and standard error of biomass (AGB and BGB) of mangrove forest, comes indirectly from table 1 of the article by Stringer <i>et al.</i> (2015). For its determination, first the mean of carbon was found for the two pools (sum of overstory and understory carbon) for each stratum (Height Class 1, ..., Height Class 5), followed by the calculation of the mean of the ecosystem (mean weighted according to the stratum areas). Finally, the carbon was converted to biomass using the conversion factor of 0.47 proposed in the IPCC good practice guide.</p> <p>Spatial level: Regional</p>									
	Value applied:	<table border="1"> <tr> <td>Semi-deciduous forest (FSD)</td> <td>144.69</td> </tr> <tr> <td>Evergreen forest (FSSV)</td> <td>123.13</td> </tr> <tr> <td>Mangrove forest (FF)</td> <td>269.01</td> </tr> </table>			Semi-deciduous forest (FSD)	144.69	Evergreen forest (FSSV)	123.13	Mangrove forest (FF)	269.01
	Semi-deciduous forest (FSD)	144.69								
	Evergreen forest (FSSV)	123.13								
	Mangrove forest (FF)	269.01								

¹⁰ Stringer, C. E.; Trettin, C. C.; Zarnoch, S. J. and Tang, W. 2015. Carbon stocks of mangroves within the Zambezi River Delta, Mozambique. *Forest Ecology Management* 354:139–148.

	The values above are estimated and extracted in the " Emission factor v.2 " workbook, and then they are recorded in the cells "B4", "B10" and "B16" respectively, of the "BIOMASS" worksheet tab in the " ZILMP Emissions Calculations MR (2021) and ZILMP Emissions Calculations MR (2022) " workbooks. These values are then applied in the range "C9:C20" of the "EMISSION MONITORING PERIOD(EMP)" worksheet tab in the " ZILMP Emissions Calculations MR (2021) and ZILMP Emissions Calculations MR (2022) " workbooks for estimating emissions.																													
QA/QC procedures applied	<p>The QA/QC procedures consisted on the following:</p> <ul style="list-style-type: none"> • SOPs were developed as described in <i>Section 2.1 - National Forest Inventory</i>. • A training on the SOPs was conducted prior to the field work. This training lasted for 3 weeks, and consisted of training on the usage of all equipment and evaluating the specific skills of each participant, in order to determine the team and brigade leaders. On the start of the 2nd phase of the IFN (2017) an additional 1-week training was conducted, to refresh the participants and train any new members. • The supervisor of each inventory team conducted a remeasurement of 4 trees per plot which means 16 trees per cluster. This served to ensure that the SOPs were adequately implemented. • An independent measurement of 10% of the plots. This activity was conducted by technicians of the National Directorate of Forests, who had participated in the Provincial Inventories of Gaza and Cabo Delgado. Diameter error must be below 10%. • The adequacy of the allometric models, including root-to-shoot ratios used was confirmed by experts of the Faculty of Agronomy and Forest Engineering (FAEF) and the Department of Biology Sciences (DCB) of the Eduardo Mondlane University (UEM). • The World Bank conducted two regular supervision missions of the National Forest Inventories to confirm the adequate implementation of the SOPs and suggest areas for improvement. The report can be found here. • An independent expert (Jim Alegria, ex-US Forestry Service) was hired in order to evaluate the methodology for the inventory and support in the estimation process, to address any gaps that were identified. The report can be found here. Many of the issues identified in the report have since been corrected, with the help of the independent expert. 																													
Uncertainty associated with this parameter:	<table border="1"> <thead> <tr> <th rowspan="2">Forest type</th> <th colspan="5">Uncertainty estimate</th> </tr> <tr> <th>Mean</th> <th>Lower (5th percentile)</th> <th>Upper bound (95th percentile)</th> <th>Half-width confidence interval at 90%</th> <th>Relative Margin</th> </tr> </thead> <tbody> <tr> <td>FSD</td> <td>144.7</td> <td>116.7</td> <td>172.1</td> <td>27.7</td> <td>0.19</td> </tr> <tr> <td>FSSV</td> <td>123.1</td> <td>101.1</td> <td>145.1</td> <td>22</td> <td>0.18</td> </tr> <tr> <td>FF</td> <td>269</td> <td>225.1</td> <td>313.8</td> <td>44.35</td> <td>0.16</td> </tr> </tbody> </table>	Forest type	Uncertainty estimate					Mean	Lower (5 th percentile)	Upper bound (95 th percentile)	Half-width confidence interval at 90%	Relative Margin	FSD	144.7	116.7	172.1	27.7	0.19	FSSV	123.1	101.1	145.1	22	0.18	FF	269	225.1	313.8	44.35	0.16
Forest type	Uncertainty estimate																													
	Mean	Lower (5 th percentile)	Upper bound (95 th percentile)	Half-width confidence interval at 90%	Relative Margin																									
FSD	144.7	116.7	172.1	27.7	0.19																									
FSSV	123.1	101.1	145.1	22	0.18																									
FF	269	225.1	313.8	44.35	0.16																									
Any comment:	-																													

Parameter:	BGB _{before,j}
Description:	Belowground biomass of forest type <i>j</i> before conversion,
Data unit:	tons of dry matter per ha
Source of data or description of the method for developing the data	<p>For semi-deciduous and evergreen forest, data are from the Zambézia Forest Inventory. It includes data that was collected in Zambézia province during the NFI, in 2017 and 2018. Please refer to parameter AGB_{before,j} for more information on how the below ground biomass was estimated.</p> <p>For mangrove forests, please refer to parameter AGB_{before,j} for more information.</p>

including the spatial level of the data (local, regional, national, international):	Spatial level: Regional																													
Value applied:	<table border="1"> <tr> <td>Semi-deciduous forest (FSD)</td> <td>49.95</td> </tr> <tr> <td>Evergreen forest (FSSV)</td> <td>42.06</td> </tr> <tr> <td>Mangrove forest (FF)</td> <td>85.43</td> </tr> </table> <p>The values above are estimated and extracted in the workbook "Emission factor v.2", and then they are recorded in the cells "B34", "B40" and "B46" respectively, of the "BIOMASS" worksheet tab in the "ZILMP Emissions Calculations MR (2021)" and "ZILMP Emissions Calculations MR (2022)" workbooks. These values are then applied in the range "E9:E20" of the "EMISSION MONITORING PERIOD(EMP)" worksheet tab in the "ZILMP Emissions Calculations MR (2021)" and "ZILMP Emissions Calculations MR (2022)" workbooks for estimating emissions.</p>	Semi-deciduous forest (FSD)	49.95	Evergreen forest (FSSV)	42.06	Mangrove forest (FF)	85.43																							
Semi-deciduous forest (FSD)	49.95																													
Evergreen forest (FSSV)	42.06																													
Mangrove forest (FF)	85.43																													
QA/QC procedures applied	Please see section QA/QC procedures under parameter $AGB_{before,j}$.																													
Uncertainty associated with this parameter:	<table border="1"> <thead> <tr> <th rowspan="2">Forest type</th> <th colspan="5">Uncertainty estimate</th> </tr> <tr> <th>Mean</th> <th>Lower (5th percentile)</th> <th>Upper bound (95th percentile)</th> <th>Half-width confidence interval at 90%</th> <th>Relative Margin</th> </tr> </thead> <tbody> <tr> <td>FSD</td> <td>49.9</td> <td>41.5</td> <td>58.4</td> <td>8.4</td> <td>0.17</td> </tr> <tr> <td>FSSV</td> <td>42.1</td> <td>35.3</td> <td>48.9</td> <td>6.8</td> <td>0.16</td> </tr> <tr> <td>FF</td> <td>85.4</td> <td>69</td> <td>101.6</td> <td>16.3</td> <td>0.19</td> </tr> </tbody> </table>	Forest type	Uncertainty estimate					Mean	Lower (5 th percentile)	Upper bound (95 th percentile)	Half-width confidence interval at 90%	Relative Margin	FSD	49.9	41.5	58.4	8.4	0.17	FSSV	42.1	35.3	48.9	6.8	0.16	FF	85.4	69	101.6	16.3	0.19
Forest type	Uncertainty estimate																													
	Mean	Lower (5 th percentile)	Upper bound (95 th percentile)	Half-width confidence interval at 90%	Relative Margin																									
FSD	49.9	41.5	58.4	8.4	0.17																									
FSSV	42.1	35.3	48.9	6.8	0.16																									
FF	85.4	69	101.6	16.3	0.19																									
Any comment:																														

Parameter:	$AGB_{after,i}$
Description:	Aboveground biomass of non-forest type <i>i</i> after conversion
Data unit:	tons of dry matter per ha
Source of data or description of the method for developing the data including the spatial level of the data (local, regional,	<p>For cropland: The values and assumptions of 2006 IPCC GL, Volume 4, Chapter 5 are used because there is no country-specific data. The agricultural land in Mozambique is mostly under the annual-crop farming practices that drive conversion of forest land to agricultural lands. So, according to 2006 IPCC GL (Volume 4, Chapter 5, Section 5.28), for lands planted in annual crops, the default value of growth in crops planted after conversion is 5 tonnes of C per hectare, based on the original IPCC Guidelines recommendation of 10 tonnes of dry biomass per hectare (dry biomass has been converted to tonnes carbon in Table 5.9) (2006 IPCC, Volume 4, Chapter 5, Section 5.28).</p> <p>For grassland: The values and assumptions of 2006 IPCC GL, Volume 4, Chapter 6 are used because there is no country-specific data. As the climate in most of Mozambique is tropical dry to subtropical dry, the value for peak-above ground biomass for tropical dry climate was used. The default value 2.30 tonnes of C per hectare from TABLE 6.4 (2006 IPCC, Volume 4, Chapter 6, Section 6.3.1.2).</p>

national, international):	For other lands: No default values exist for these conversions. Spatial level: International																													
Value applied:	<table border="1"> <tr> <td>Cropland (C)</td> <td>10.00</td> </tr> <tr> <td>Grassland (P)</td> <td>2.30</td> </tr> <tr> <td>Other lands (A O U)</td> <td>0.00</td> </tr> </table> <p>The values above are recorded in the ranges "B5:B9", "B11:B15" and "B17:B21" of the "BIOMASS" worksheet tab in the "ZILMP Emissions Calculations MR (2021) and ZILMP Emissions Calculations MR (2022)" workbooks. These values are then applied in the range "D9:D20" of the "EMISSION MONITORING PERIOD(EMP)" worksheet tab in the "ZILMP Emissions Calculations MR (2021) and ZILMP Emissions Calculations MR (2022)" workbooks for estimating emissions.</p>	Cropland (C)	10.00	Grassland (P)	2.30	Other lands (A O U)	0.00																							
Cropland (C)	10.00																													
Grassland (P)	2.30																													
Other lands (A O U)	0.00																													
QA/QC procedures applied	The adequacy in the use of these default values was confirmed with the experts in GHG Inventory in DINAB.																													
Uncertainty associated with this parameter:	<table border="1"> <thead> <tr> <th rowspan="2">Non-forest type</th> <th colspan="5">Uncertainty estimate</th> </tr> <tr> <th>Mean</th> <th>Lower (5th percentile)</th> <th>Upper (95th percentile)</th> <th>Half-width confidence interval at 90%</th> <th>Relative Margin</th> </tr> </thead> <tbody> <tr> <td>Cropland (C)</td> <td>10</td> <td>3.9</td> <td>16.1</td> <td>6.1</td> <td>0.61</td> </tr> <tr> <td>Grassland (P)</td> <td>2.3</td> <td>0.9</td> <td>3.7</td> <td>1.4</td> <td>0.61</td> </tr> <tr> <td>Other lands (A O U)</td> <td>0</td> <td>0</td> <td>0</td> <td>0</td> <td>NA</td> </tr> </tbody> </table>	Non-forest type	Uncertainty estimate					Mean	Lower (5 th percentile)	Upper (95 th percentile)	Half-width confidence interval at 90%	Relative Margin	Cropland (C)	10	3.9	16.1	6.1	0.61	Grassland (P)	2.3	0.9	3.7	1.4	0.61	Other lands (A O U)	0	0	0	0	NA
Non-forest type	Uncertainty estimate																													
	Mean	Lower (5 th percentile)	Upper (95 th percentile)	Half-width confidence interval at 90%	Relative Margin																									
Cropland (C)	10	3.9	16.1	6.1	0.61																									
Grassland (P)	2.3	0.9	3.7	1.4	0.61																									
Other lands (A O U)	0	0	0	0	NA																									
Any comment:																														

Parameter:	BGB _{after,i}		
Description:	Belowground biomass of non-forest type <i>i</i> after conversion		
Data unit:	tons of dry matter per ha		
Source of data or description of the method for developing the data including the spatial level of the data (local, regional, national, international):	<p>For cropland: The values and assumptions of 2006 IPCC GL, Volume 4, Chapter 5 are used because there is no country-specific data. Tier 2 may modify the assumption that carbon stocks immediately following conversion is zero. In this case, it is assumed that conversion leads to annual croplands and in the case the carbon stock in biomass after one year for annual crops provided in TABLE 5.9 is used.</p> <p>For grassland: The values and assumptions of 2006 IPCC GL, Volume 4, Chapter 6 are used because there is no country-specific data. As the climate in most of Mozambique is tropical dry to subtropical dry, the value for peak-above ground biomass for tropical dry climate was used. The default value 2.30 tonnes of C per hectare from TABLE 6.4 (2006 IPCC, Volume 4, Chapter 6, Section 6.3.1.2).</p> <p>For other lands: No default values exist for these conversions. Spatial level: International</p>		
Value applied:	<table border="1"> <tr> <td>Cropland (C)</td> <td>0.00</td> </tr> </table>	Cropland (C)	0.00
Cropland (C)	0.00		

	<table border="1"> <tr> <td>Grassland (P)</td> <td>6.40</td> </tr> <tr> <td>Other lands (A O U)</td> <td>0.00</td> </tr> </table> <p>The values above are recorded in the ranges "B35:E39", "B41:B45" and "B47:B51" of the "BIOMASS" worksheet tab in the "ZILMP Emissions Calculations MR (2021) and ZILMP Emissions Calculations MR (2022)" workbooks. These values are then applied in the range "F9:F20" of the "EMISSION MONITORING PERIOD(EMP)" worksheet tab in the "ZILMP Emissions Calculations MR (2021) and ZILMP Emissions Calculations MR (2022)" workbooks for estimating emissions.</p>	Grassland (P)	6.40	Other lands (A O U)	0.00																									
Grassland (P)	6.40																													
Other lands (A O U)	0.00																													
QA/QC procedures applied	The adequacy in the use of these default values was confirmed with the experts in GHG Inventory in DINAB.																													
Uncertainty associated with this parameter:	<table border="1"> <thead> <tr> <th rowspan="2">Non-forest type</th> <th colspan="5">Uncertainty estimate</th> </tr> <tr> <th>Mean</th> <th>Lower (5th percentile)</th> <th>Upper bound (95th percentile)</th> <th>Half-width confidence interval at 90%</th> <th>Relative Margin</th> </tr> </thead> <tbody> <tr> <td>Cropland (C)</td> <td>0</td> <td>0</td> <td>0</td> <td>0</td> <td>NA</td> </tr> <tr> <td>Grassland (P)</td> <td>6.4</td> <td>-0.2</td> <td>12.8</td> <td>6.5</td> <td>1.02</td> </tr> <tr> <td>Other lands (A O U)</td> <td>0</td> <td>0</td> <td>0</td> <td>0</td> <td>NA</td> </tr> </tbody> </table>	Non-forest type	Uncertainty estimate					Mean	Lower (5 th percentile)	Upper bound (95 th percentile)	Half-width confidence interval at 90%	Relative Margin	Cropland (C)	0	0	0	0	NA	Grassland (P)	6.4	-0.2	12.8	6.5	1.02	Other lands (A O U)	0	0	0	0	NA
Non-forest type	Uncertainty estimate																													
	Mean	Lower (5 th percentile)	Upper bound (95 th percentile)	Half-width confidence interval at 90%	Relative Margin																									
Cropland (C)	0	0	0	0	NA																									
Grassland (P)	6.4	-0.2	12.8	6.5	1.02																									
Other lands (A O U)	0	0	0	0	NA																									
Any comment:																														

Parameter:	$A(j,i)_{RP}$
Description:	Area converted from forest type j to non-forest type i during the reference period.
Data unit:	hectare per year.
Source of data and description of measurement /calculation methods and procedures applied:	<p><i>i. Approach and source</i></p> <p>Activity data (AD) for deforestation were obtained from an annual historical time series analysis of land use, land-use change and forestry (LULUCF) carried out by five trained operators in approximately 98 effective working days (4.4 months), for the period of 2001 – 2016 across the country, using the Collect Earth Open tool.</p> <p>Activity data have been generated following IPCC Approach 3 for representing the activity data as described in the 2006 IPCC Guidelines for National Greenhouse Gas Inventories (Volume 4, Chapter 3, Section 3.13), i.e., using spatially-explicit observations of land-use categories and land-use conversions over time across the country, derived from sampling of geographically located points. The result was forest cover data for 2016 and forest cover change data for every year from 2001 to 2016.</p> <p>The period of AD analysis from 2005 to 2015 (11 years) considered for the ER Program area, could be adapted within the general period 2001 – 2016 with little effort, due to the operators collecting the date of the LULC change.</p> <p><i>ii. Sampling design</i></p>

A systematic 4 x 4 km grid consisting of a total of 48,894 sampling units was established at a national level to generate the historical activity data for the entire area of the country using high and medium resolution imagery. This is the same grid used to allocate the NFI clusters from the Stratified Random Sampling design. At jurisdictional level, this corresponds to 3,308 sampling units being interpreted. Each sampling unit was visually assessed and its information was collected and entered in a complete database on LULC changes at the national level.

iii. **Response design**

Spatial sampling unit

The spatial sampling unit was defined as a square with 1 ha (100m x 100m), where an internal grid of 5 x 5 points (20m x 20m grid) is overlapped. Each point from the internal grid has a weight coverage of 4%.



Spatial sampling unit for the reference period

Source of reference data

The sampling approach for historical AD calculation based on the regular National 4 x 4 km grid has been designed and conducted using the high and medium resolution images repository available through Google Earth and Earth Engine as a visual assessment exercise. These imageries with digital forms designed to collect the LULCC information on the points of the grid are automatically accessible through the Collect Earth tool (www.openforis.org) along with scripts accessible through Earth Engine code that facilitate vegetation type's interpretation (e.g. MODIS or Landsat NDVI time series). Each point of the grid is photo-interpreted thanks to Collect Earth tool and the year and type of changes are also collected.

The use of various scripts programmed on Earth Engine Code facilitates the interpretation of the vegetation type and the determination of LULC changes. Specifically, the MOD13Q1 (NDVI 16-day Global Modis 250 m) graphic from 2001-2016, most recent Sentinel-2 image, most recent Landsat-8 pan sharpened image, Landsat-7 pan sharpened image (2000, 2004, 2008, 2012), etc.

The completeness of the series is guaranteed using remote sensing products from medium resolution imagery repositories from 2001 (e.g. Annual TOA Reflectance Composite, Annual NDVI Composite, Annual EVI Composite, Annual Greenest-Pixel TOA Reflectance Composite, etc. from Landsat 5 TM) and the most recent Sentinel-2 image from 2016. In this way, a temporal analysis of LULC changes has been completed for each sampling point of the national 4 x 4 km grid (48,894 records).

The image displays four digital forms from the Collect Earth Tool, arranged around a central satellite image of a forest area. The forms are:


- Top Left:** A form for selecting land cover elements. It lists categories like 'Árvores' (40-49%), 'Arbustos' (50-59%), 'Matagais', 'Gramíneas', 'Solo Exposto', 'Cultivos', 'Rio', 'Lago', and 'Infraestrutura', each with a dropdown menu for coverage percentage.
- Top Right:** A 'Lulc Change' form with a grid of buttons for transitions (e.g., C\rightarrowC, F\rightarrowF, P\rightarrowP, A\rightarrowA, U\rightarrowU, O\rightarrowO, T\rightarrowC, T\rightarrowF, T\rightarrowP, T\rightarrowA, T\rightarrowD). It also includes a 'Degradação' section with 'Sim' and 'Não' buttons and a 'Mudança estimada na CD' dropdown set to '20-29%'.
- Bottom Left:** An 'Uso da terra IPCC' form with buttons for 'Cultivos', 'Florestas', 'Pastagens', 'Áreas alagadas', 'Áreas urbanas', and 'Outras Terras'. It includes 'Confiança IPCC' (Sim/Não), 'Classe nacional' (Floresta (Semi-) decídua fechada), 'Confiança classe nacional' (Sim/Não), 'Subclasse nacional' (Miombo denso), and 'Confiança subclasse nacional' (Sim/Não).
- Bottom Right:** A form for satellite image data entry, including 'Imagem de satélite (fornecedor)' (Digital Globe), 'Data da imagem atual' (12/14/2016), 'Imagem de satélite' (Outro), 'Produto' (Outro), and 'Data da imagem antiga' (12/05/2001).

Arrows point from the forms to a central satellite image of a forest area with a yellow bounding box and a green label '63866'.

LULCCF detection using Collect Earth Tool (www.openforis.org). Digital forms designed with Collect Tool.

Reference labelling protocol

The activity data was generated considering the national land use and land cover classification system, which reflects the six broad IPCC Land Use categories. A set of hierarchical rules were established and used to determine the LULCCF category based on a certain percentage and taking into account the national forest definition as well. A single land use class is easier to classify, but it becomes challenging when there is a combination of two or more land use classes within the area of interest. Thus, this is where the hierarchical rules are important to determine the land use. Any sampling unit that has 30% of tree canopy cover is considered a forest, according to the national forest definition, even if it has more than 20% of settlements, crops or other land use, the forest has priority. In the case the sampling unit was classified as forest land and different forest types were present in the sampling unit, a majority rule was used in this case, i.e. the largest forest class is the winner.



Decision tree for the attribution of the LULCCF category based on the percentage cover of the elements present in the sampling unit of 1 ha.

iv. Analysis

The estimation of the areas corresponding to a certain category changes from a forest type to a non-forest type in the framework of this systematic sampling approach was based on assessments of area proportions. According to 2006 IPCC Guidelines for National Greenhouse Gas Inventories (Volume 4, Chapter 3, Section 3.33), the proportion of each land-use or land-use change category is calculated by dividing the number of points located in the specific category by the total number of points, and area estimates for each land-use or land-use change category are obtained by multiplying the proportion of each category by the total area of interest, in this case, the ER Program accounting area.

$$A_i = p_i \times A \quad \text{Equation 7}$$

Where:

- A_i Area estimate on forest type j converted to non-forest type i ; hectare
- p_i Proportion of points on forest type j converted to non-forest type i ; dimensionless
- A Total area of interest; hectare

$$p_i = \frac{n_i}{N} \quad \text{Equation 8}$$

Where:

- n_i Number of points on forest type j converted to non-forest type i ; number
- N Total number of points; number

Uncertainties in activity data were derived using non-parametric bootstrapping, where reference data points were re-sampled with replacement 100,000 times. For each permutation of reference data points, the bias-corrected area estimates were produced following the methods described in Olofsson *et al.* (2014). Uncertainty was estimated from the resulting distribution of area estimates. Although more complex to implement, bootstrapping has the advantages of not requiring any assumption about the shape of the probability distribution function of each land cover transition class, and avoids the generation of negative areas in rare classes where a probability distribution function crosses zero. The method was implemented in R, and the scripts used are available in the "[Mozambique ERPA 2020](#)" shared folder.

The impact of using non-parametric bootstrapping to estimate uncertainties vs other methods was tested with a comparison of deforested areas derived from bootstrapping against sampling from a normal distribution with standard error calculated with the methods described in Olofsson *et al.* (2014) (Figure 7). For the latter case two uncertainties were derived: one retaining any negative area estimates for rare transition classes, and another setting these to zero. The result indicates that there is very little difference between any of the methods in either reference or monitoring periods, with the result that any chosen approach would produce equivalent emissions estimates.




Figure 7: Total activity data area estimates for reference period using normal distributions for each transition class (red), normal distributions with a minimum area of 0 ha (green), and non-parametric bootstrapping (blue). All three methods result in equivalent uncertainty estimates.

Value applied

Semi-deciduous forest to cropland	11,785.07
Semi-deciduous forest to grassland	1,745.94
Semi-deciduous forest to other lands	145.49
Evergreen forest to cropland	3,200.88
Evergreen forest to grassland	145.49
Evergreen forest to other lands	0.0
Mangrove forest to cropland	0.0
Mangrove forest to grassland	0.0
Mangrove forest to other lands	0.0

QA/QC procedures applied:	<p>Quality Control consisted in having a team of 5 technicians with experience in forests and remote sensing, all trained together by an MRV specialist. The team worked in the same office, and discussed any classification issues with each other.</p> <p>Quality Assurance was conducted using the SAIKU extension of Collect Earth. This tool allows the detection of whether:</p> <ul style="list-style-type: none"> i) Data point was not filled ii) The class assigned followed the classification hierarchy, based on the % of individual element cover iii) Year of the Old image/Change image was less than the current image iv) Change classes are consistent with previous and current classes v) Open and closed forest was correctly classified, based on the 30% (open) and 65% (closed) cover threshold <p>In the case of any error being detected, the ID of the data point was registered and the user performed the necessary corrections.</p>																																													
Uncertainty associated with this parameter:	<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr style="background-color: #d9ead3;"> <th rowspan="2">Category change</th> <th colspan="5">Uncertainty estimate (from non-parametric bootstrapping)</th> </tr> <tr style="background-color: #d9ead3;"> <th>Median</th> <th>Lower bound (5th percentile)</th> <th>Upper bound (95th percentile)</th> <th>Half-width confidence interval at 90%</th> <th>Relative Margin</th> </tr> </thead> <tbody> <tr> <td>FSD>(A O U)</td> <td>291</td> <td>0</td> <td>873</td> <td>436.5</td> <td>1.5</td> </tr> <tr> <td>FSD>C</td> <td>23570.1</td> <td>19496.3</td> <td>27935</td> <td>4219.4</td> <td>0.18</td> </tr> <tr> <td>FSD>P</td> <td>3491.9</td> <td>2036.9</td> <td>5237.8</td> <td>1600.4</td> <td>0.46</td> </tr> <tr> <td>FSSV>C</td> <td>6401.8</td> <td>4364.8</td> <td>8729.7</td> <td>2182.5</td> <td>0.34</td> </tr> <tr> <td>FSSV>P</td> <td>291</td> <td>0</td> <td>873</td> <td>436.5</td> <td>1.5</td> </tr> </tbody> </table>					Category change	Uncertainty estimate (from non-parametric bootstrapping)					Median	Lower bound (5 th percentile)	Upper bound (95 th percentile)	Half-width confidence interval at 90%	Relative Margin	FSD>(A O U)	291	0	873	436.5	1.5	FSD>C	23570.1	19496.3	27935	4219.4	0.18	FSD>P	3491.9	2036.9	5237.8	1600.4	0.46	FSSV>C	6401.8	4364.8	8729.7	2182.5	0.34	FSSV>P	291	0	873	436.5	1.5
Category change	Uncertainty estimate (from non-parametric bootstrapping)																																													
	Median	Lower bound (5 th percentile)	Upper bound (95 th percentile)	Half-width confidence interval at 90%	Relative Margin																																									
FSD>(A O U)	291	0	873	436.5	1.5																																									
FSD>C	23570.1	19496.3	27935	4219.4	0.18																																									
FSD>P	3491.9	2036.9	5237.8	1600.4	0.46																																									
FSSV>C	6401.8	4364.8	8729.7	2182.5	0.34																																									
FSSV>P	291	0	873	436.5	1.5																																									
Any comment:																																														

3.2 Monitored Data and Parameters

Parameter:	$A(j,i)_{MP}$																																			
Description:	Area converted from forest type j to non-forest type i during the Monitoring Period.																																			
Data unit:	hectare per year.																																			
Value monitored during this Monitoring / Reporting Period:	<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr style="background-color: #d9ead3;"> <th>Type of change</th> <th>2021</th> <th>2022</th> </tr> </thead> <tbody> <tr> <td>Semi-deciduous forest to cropland</td> <td>33577.09</td> <td>17939.15</td> </tr> <tr> <td>Semi-deciduous forest to grassland</td> <td>250.14</td> <td>719.44</td> </tr> <tr> <td>Semi-deciduous forest to other lands</td> <td>0</td> <td>679.08</td> </tr> <tr> <td>Evergreen forest to cropland</td> <td>1878.42</td> <td>1792.63</td> </tr> <tr> <td>Evergreen forest to grassland</td> <td></td> <td></td> </tr> <tr> <td>Evergreen forest to other lands</td> <td></td> <td></td> </tr> <tr> <td>Mangrove forest to cropland</td> <td>98.91</td> <td></td> </tr> <tr> <td>Mangrove forest to grassland</td> <td></td> <td></td> </tr> <tr> <td>Mangrove forest to other lands</td> <td></td> <td></td> </tr> <tr> <td>Total</td> <td>35,804.56</td> <td>21130.3</td> </tr> </tbody> </table>			Type of change	2021	2022	Semi-deciduous forest to cropland	33577.09	17939.15	Semi-deciduous forest to grassland	250.14	719.44	Semi-deciduous forest to other lands	0	679.08	Evergreen forest to cropland	1878.42	1792.63	Evergreen forest to grassland			Evergreen forest to other lands			Mangrove forest to cropland	98.91		Mangrove forest to grassland			Mangrove forest to other lands			Total	35,804.56	21130.3
Type of change	2021	2022																																		
Semi-deciduous forest to cropland	33577.09	17939.15																																		
Semi-deciduous forest to grassland	250.14	719.44																																		
Semi-deciduous forest to other lands	0	679.08																																		
Evergreen forest to cropland	1878.42	1792.63																																		
Evergreen forest to grassland																																				
Evergreen forest to other lands																																				
Mangrove forest to cropland	98.91																																			
Mangrove forest to grassland																																				
Mangrove forest to other lands																																				
Total	35,804.56	21130.3																																		

Source of data and description of measurement /calculation methods and procedures applied:

i. Source
Activity data used for the monitoring period are obtained from a combination of an annual wall-to-wall deforestation map with sampling to generate deforested area estimates through a stratified estimator.

ii. Variable of interest
The variable of interest are all the transitions specified above. It is important to note that the variables of interest are not aligned to the strata as this is not required. Strata is linked to the likelihood of presence of deforestation events, whereas the variable of interest is linked to the possible transitions of deforestation per forest type and post-deforestation type.

ii. Annual deforestation map
The workflow used to produce annual deforestation map for the ZILMP program area follows the steps below:

1. Produce two Sentinel-2 satellite imagery composites for the monitoring area, containing all images of wet season (i.e. January - May). For 2021 deforestation map, the first composite (reference period) comprises the period between January 2021 to May 2021 and January 2022 to May 2022 for 2022 deforestation map and the second composite (actual period) comprises the period from January 2022 to May 2022 and January 2023 to May 2023 respectively. The reason behind the selection of January - May as a reference and actual period of monitoring resides on the fact that it is the wet season, where the NDVI stability is very high in relation to the dry season, which starts in June to October, when most trees lose their foliage and makes it difficult the analysis of deforestation.
2. Generate image features from reference period and actual period from the composites generated in previous step, to identify changes in forest cover. The image features have different vegetation indexes, namely, NDVI, EVI, SAVI, NBR, NDWI with respective sub-products such as NDVI 90th percentile, Normalized NDVI, and variation on NDVI.
3. Generate training data on classes of deforestation, stable forest and stable non-forest by visual interpretation of composites from the reference and actual periods, and NDVI change detection image. The NDVI change detection image is a result of the difference of NDVI from the composites of reference and actual periods. The calculated NDVI change detection image helps the interpreter to locate where the changes of forest cover are occurring.
4. Produce a categorical deforestation map from training data and image features through a process of classification using Random Forest classifier. The Categorical deforestation map includes non-forest stable and stable forest classes. Because errors of omission of deforestation have a very large impact on the final estimates, it is important to reduce these errors as much as possible.
5. To improve the efficacy of the sampling the deforestation class on the map is reclassified as:
 - a) High probability deforestation (cluster of more than 10 pixels of deforestation, corresponding to at least 40% of one hectare);
 - b) Low probability of deforestation (cluster of less than 10 pixels and greater than 2 pixels, corresponding at least 12% to 40% of one hectare) and;

- c) Non-forest (cluster of less than 3 pixels, corresponding to less than 12% of one hectare).
6. To reduce the risk of omission errors, a Buffer of 60 meters is added around the high probability of deforestation class. The result is a deforestation map with five classes: *High probability of deforestation; buffer; low probability of deforestation; stable forest and stable non-forest.*

v. Sampling design

Sampling method

Monitoring of activity data for annual reporting is conducted using a stratified estimator, where deforestation map (which includes classes of forest and non-forest) is used for stratification and reference-sampling units are used for estimate activity data and associated confidence intervals.

Sample size determination

The sample size n was determined from the equation:

$$n = \frac{(\sum W_i S_i)^2}{[S(\hat{O})]^2 + \left(\frac{1}{N}\right) \sum W_i S_i} \approx \left(\frac{\sum W_i S_i}{S(\hat{O})}\right)^2 \quad \text{Equation 9}$$

Where:

- N Number of units in the ROI
- $S(\hat{O})$ Standard error of the estimated overall accuracy that we would like to achieve
- W_i Mapped proportion of area of class i ; and
- S_i Standard deviation of stratum i .

The standard deviation of stratum i is given by the formula:

$$S_i = \sqrt{U_i(1 - U_i)} \quad \text{Equation 10}$$

Where:

- U_i Proportion of area of deforestation in stratum i .

In order to obtain approximate values of proportion of deforestation in each stratum (U_i), a pilot sampling is conducted. This pilot consists of 100 sample units per stratum.

Sample units per stratum

After the pilot sampling, sample units may need to be added to each stratum, in order to achieve 20% relative margin error at 95% confidence level. It was decided to use the Optimum (Neyman) allocation for each change stratum, where the stratum standard deviation $S_i = \sqrt{U_i \cdot (1 - U_i)}$ increases the number of plots (ensuring larger numbers of plots in rare classes or strata) and sampling unit costs are constant:

$$n_i = n \frac{w_i \cdot S_i}{\sum_{i=1}^I w_i \cdot S_i} \quad \text{Equation 11}$$

For each stable stratum, the proportional allocation is applied if deforestation omission errors are completely absent from these strata. In stratified sampling the sample size for proportional allocation is given by:

$$n_i = n \cdot w_i \quad \text{Equation 12}$$

The number of reference points is presented in Table 4.

Table 4: Number of reference sampling units per map stratum for each classes of the 2021 and 2022 maps.

Stratum	Number of sample units (2021)	Number of sample units (2022)
High probability of deforestation	100	105
60 m Buffer	309	445
Low probability of deforestation	100	100
Forest	300	300
Non-forest	300	300
Total	1,109	1,250

v. Response design

Sampling unit and spatial support

The sampling unit is a 20 m pixel of the stratification map that was produced. The spatial support used is a 100m x 100m plot (1ha). Each Spatial sampling unit contains an internal grid of 5 x 5 points (20m x 20m grid) to aid in the labelling attribution (Figure 8).



Figure 8: Spatial sampling unit.

Source of reference data

Each sampling unit was evaluated using Collect Earth (<http://www.openforis.org/>). This tool enables access to high-resolution images in Google Earth, Bing Maps and Planet Labs, as well as a medium resolution image repository available through Google Earth Engine Explorer and Code Editor (Landsat and Sentinel-2). The tool enables to display digital forms designed to collect the Land-Use Land Cover Change and Forestry (LULCCF) information on the sampling

points (Figure 9). The Earth Engine Code Editor facilitates the interpretation of the vegetation type and the determination of LULC changes, by displaying the historical MOD13Q1 (NDVI 16-day Global Modis 250 m) graphic as well as monthly mosaics of Sentinel-2 images. The main source of data to identify changes in land cover, is Sentinel-2 15 days' reflectance composite. However, Planet data is also used in cases of doubt or excessive cloud cover with Sentinel-2.




Figure 9: LULCCF detection using Collect Earth Tool (www.openforis.org). Digital forms designed with Collect Tool.

Reference labelling protocol

The activity data was generated considering the national land use and land cover classification system, which reflects the six broad IPCC Land Use categories.

A set of hierarchical rules were established and used to determine the LULCCF category based on a certain percentage and taking into account the national forest definition as well (Figure 10). A single land use class is easier to classify, but it becomes challenging when there is a combination of two or more land use classes within the area of interest. Thus, this is where the hierarchical rules are important to determine the land use. Any sampling unit that has 30% of tree canopy cover is considered a forest, according to the national forest definition, even if it has more than 20% of settlements, crops or other land use, the forest is priority.

In the case the sampling unit was classified as forest land and different forest types were present in the sampling unit, a majority rule was used in this case, i.e. the largest forest class is chosen (please click [here](#) for more details).




Figure 10: Decision tree for the attribution of the LULCCF category based on the percentage cover of the elements present in the sampling unit of 1 ha.

vi. Analysis

Applying the methodology described in Olofsson *et al.* (2014)¹¹ and the GFOI MGD (<https://www.reddcompass.org/mgd>), the estimations of the areas corresponding to land-use and land-cover change categories, more specifically the activity data for deforestation, in the framework of this stratified random sampling approach (based on the visual assessment of the 1 ha plots) was based on assessments of area proportions. A sample error matrix is constructed where the map classes ($h=1, 2, \dots, q$) are represented by rows and the reference data ($k=1, 2, \dots, q$) by columns as shown in Table 5. The size of strata and original proportion matrix can be found in [2021](#) and [2022](#) spreadsheets.

Table 5: Error matrix of area proportions.

Map data	Reference data					Total	User's accuracy (\hat{d}_i)
	Deforestation			Stable forest	Stable non-forest		
	High probability of deforestation	40 m Buffer	Low probability of deforestation				
High probability of deforestation	\hat{p}_{11}	\hat{p}_{12}	\hat{p}_{13}	\hat{p}_{14}	\hat{p}_{15}	\hat{p}_1	\hat{p}_{11}/\hat{p}_1
40 m Buffer	\hat{p}_{21}	\hat{p}_{22}	\hat{p}_{23}	\hat{p}_{24}	\hat{p}_{25}	\hat{p}_2	\hat{p}_{22}/\hat{p}_2
Low probability of deforestation	\hat{p}_{31}	\hat{p}_{32}	\hat{p}_{33}	\hat{p}_{34}	\hat{p}_{35}	\hat{p}_3	\hat{p}_{33}/\hat{p}_3
Stable forest	\hat{p}_{41}	\hat{p}_{42}	\hat{p}_{43}	\hat{p}_{44}	\hat{p}_{45}	\hat{p}_4	\hat{p}_{44}/\hat{p}_4
Stable non-forest	\hat{p}_{51}	\hat{p}_{52}	\hat{p}_{53}	\hat{p}_{54}	\hat{p}_{55}	\hat{p}_5	\hat{p}_{55}/\hat{p}_5
Total	\hat{p}_1	\hat{p}_2	\hat{p}_3	\hat{p}_4	\hat{p}_5	1	
Producer's accuracy (P_i)	\hat{p}_{11}/\hat{p}_1	\hat{p}_{22}/\hat{p}_2	\hat{p}_{33}/\hat{p}_3	\hat{p}_{44}/\hat{p}_4	\hat{p}_{55}/\hat{p}_5		Overall accuracy (\hat{d}) $= \hat{p}_{11} + \hat{p}_{22} + \hat{p}_{33} + \hat{p}_{44} + \hat{p}_{55}$

The mean estimator for the area of each class can be directly obtained from the error matrix. Unbiased stratified estimators are provided using reference class area proportions ($\hat{p}_{.k}$):

$$\hat{p}_{.k} = \sum_{h=1}^H w_h \cdot \frac{n_{hk}}{n_h} = \sum_{h=1}^H \hat{p}_{hk} \quad \text{Equation 13}$$

Where:

$\hat{p}_{.k}$ Area proportions of reference data class k . These proportions of reference data for deforestation classes as a whole are collapsed in three possible types of conversions/transitions from forest type j to non-forest type i , namely:

- Broadleaved (Semi-) deciduous to Non-forest type i ;
- Broadleaved (Semi-) evergreen to Non-forest type i ; and
- Mangrove to Non-forest type i .

Five types of non-forest land are considered:

- Cropland (C);
- Grassland (P);
- Wetland (A);
- Settlement (U); and
- Other lands (O).

w_h Proportion of area mapped as class h ;

n_{hk} Sample count at cell (h,k) ;

n_h Sum of sample counts across row h ; and

\hat{p}_{hk} Proportion of area in cell (h,k) .

Once the estimated reference class area proportions ($\hat{p}_{.k}$) are obtained, the mean total area per class is calculated by multiplying them with the total reporting area a :

$$\hat{A}_j = \hat{p}_{.k} \cdot a \quad \text{Equation 14}$$

Uncertainty in activity data were derived using non-parametric bootstrapping, where reference data points were re-sampled with replacement 100,000 times. For each permutation of reference data points, the bias-corrected area estimates were produced following the methods described in Olofsson *et al.* (2014). Uncertainty was estimated from the resulting distribution of area estimates. Although more complex to implement, bootstrapping has the advantages of not requiring any assumption about the shape of the probability distribution function of each land cover transition class, and avoids the generation of negative areas in rare classes where a probability distribution function crosses zero. The method was implemented in R, and the scripts used are available in the “*Mozambique Monitoring report*” shared folder.

The impact of using non-parametric bootstrapping to estimate uncertainties vs other methods was tested with a comparison of deforested areas derived from bootstrapping against sampling from a normal distribution with standard error calculated with the methods described in Olofsson *et al.* (2014). For the latter case, two uncertainties were derived: one retaining any negative area estimates for rare transition classes, and another setting these to zero. The result indicates that there is very little difference between any of the methods in either reference or monitoring periods, with the result that any chosen approach would produce equivalent emissions estimates.

¹¹ Olofsson, P., Foody, G.M., Herold, M., Stehman, S.V., Woodcock, C.E., & Wulder, M.A. 2014. Good practices for estimating area and assessing accuracy of land change. *Remote Sensing of Environment*. 148:42-57.




Figure 11: Total activity data area estimates for the monitoring period using normal distributions for each transition class (red), normal distributions with a minimum area of 0 ha (green), and non-parametric bootstrapping (blue). All three methods result in equivalent uncertainty estimates.

QA/QC procedures applied:

The QA/QC procedures consisted on the following:

- SOPs were developed as described in Section 2.1 - Satellite and land monitoring system and training; and
- Interpretation is done by highly qualified professionals which are specialized in land cover interpretation with satellite imagery. They were trained and a robust control system is in place to ensure that they are correctly calibrated throughout the data collection process.
- All reference data interpreted as deforestation or forest degradation, and an additional 20% of the remaining reference data were evaluated. The quality control is carried out by two independent supervisors, who after the independent evaluation compare the two evaluations and consensually compile a single comment for each sample. The parameters to be taken into account in the evaluation for identifying errors are: a) the percentage of coverage for each element within the plot; b) the current land cover/land use class (levels 1 and 2); c) the land cover/land use change class; d) the former land cover/land use class (levels 1 and

	<p>2); and e) the date of occurrence of land cover/land use change, or evidence date of remaining land cover/land use. If there are gross errors related to the parameters b), c) and d) in at least 20% of samples from the 20% mentioned initially, the respective interpreter should review all samples from the batch, otherwise the interpreter reviews only the samples evaluated by the supervisors, that present gross errors. On the other hand, in relation to all samples interpreted as deforestation, the interpreter reviews only the samples that present gross errors according to the evaluation from the supervisors. The process is cyclical until the interpreter achieves values less than 20% of gross errors in the batch.</p> <ul style="list-style-type: none"> The uncertainty analysis approach was reviewed by Philip Mundhenk, a professor of the University of Hamburg specialized in Monte Carlo simulations. 																																									
Uncertainty for this parameter:	<table border="1"> <thead> <tr> <th rowspan="2">Category change</th> <th colspan="5">Uncertainty estimate (from non-parametric bootstrapping)</th> </tr> <tr> <th>Median</th> <th>Lower bound (5th percentile)</th> <th>Upper bound (95th percentile)</th> <th>Half-width confidence interval at 90%</th> <th>Relative Margin</th> </tr> </thead> <tbody> <tr> <td>FF>C</td> <td>95.10</td> <td>-</td> <td>290.90</td> <td>145.45</td> <td>1.53</td> </tr> <tr> <td>FSD>(A O U)</td> <td>660.20</td> <td>171.40</td> <td>1,286.40</td> <td>557.50</td> <td>0.84</td> </tr> <tr> <td>FSD>C</td> <td>51,476.80</td> <td>45,202.20</td> <td>58,146.20</td> <td>6,472.00</td> <td>0.13</td> </tr> <tr> <td>FSD>P</td> <td>943.50</td> <td>-</td> <td>2,004.20</td> <td>1,002.10</td> <td>1.06</td> </tr> <tr> <td>FSSV>C</td> <td>3,586.30</td> <td>1,978.30</td> <td>5,670.60</td> <td>1,846.15</td> <td>0.51</td> </tr> </tbody> </table>	Category change	Uncertainty estimate (from non-parametric bootstrapping)					Median	Lower bound (5 th percentile)	Upper bound (95 th percentile)	Half-width confidence interval at 90%	Relative Margin	FF>C	95.10	-	290.90	145.45	1.53	FSD>(A O U)	660.20	171.40	1,286.40	557.50	0.84	FSD>C	51,476.80	45,202.20	58,146.20	6,472.00	0.13	FSD>P	943.50	-	2,004.20	1,002.10	1.06	FSSV>C	3,586.30	1,978.30	5,670.60	1,846.15	0.51
Category change	Uncertainty estimate (from non-parametric bootstrapping)																																									
	Median	Lower bound (5 th percentile)	Upper bound (95 th percentile)	Half-width confidence interval at 90%	Relative Margin																																					
FF>C	95.10	-	290.90	145.45	1.53																																					
FSD>(A O U)	660.20	171.40	1,286.40	557.50	0.84																																					
FSD>C	51,476.80	45,202.20	58,146.20	6,472.00	0.13																																					
FSD>P	943.50	-	2,004.20	1,002.10	1.06																																					
FSSV>C	3,586.30	1,978.30	5,670.60	1,846.15	0.51																																					
Any comment:	-																																									

4 QUANTIFICATION OF EMISSION REDUCTIONS

4.1 ER Program Reference level for the Monitoring / Reporting Period covered in this report

See the workbook template for more details [here](#) Estimation of emissions by sources and removals by sinks included in the ER Program's scope

Year of Monitoring / Reporting Period t	Average annual historical Emissions from deforestation over the Reference Period (tCO _{2-e} /yr)	If applicable, average annual historical Emissions from forest degradation over the Reference Period (tCO _{2-e} /yr)	If applicable, average annual historical enhanced Removals from afforestation/reforestation (AR) (tCO _{2-e} /yr)	If applicable, average annual historical enhanced Removals from other activities besides A/R (tCO _{2-e} /yr)	Adjustment, if applicable (tCO _{2-e} /yr)	Reference Level (tCO _{2-e} /yr)
2021	5,253,267	0	0	0	0	5,253,267
2022	5,253,267	0	0	0	0	5,253,267
Total	10,506,534	0	0	0	0	10,506,534

4.2 Estimation of emissions by sources and removals by sinks included in the ER Program's scope

The following table shows the emissions results obtained per category changes from a forest type to a non-forest type during the Monitoring Period. The emissions are generated relating the data and parameters described in Section 3 and summarized in the Table 6, by applying **Equation 6**.

Table 6: Calculation of the emissions from the ER Program during the Monitoring Period (2021 and 2022).

Category changes	AGB _{before,j} (tdm/ha)	BGB _{before,j} (tdm/ha)	AGB _{after,i} (tdm/ha)	BGB _{after,i} (tdm/ha)	2021		2022	
					A _{(j,i)MP} (ha)	Emissions (tCO ₂ e)	A _{(j,i)MP} (ha)	Emissions (tCO ₂ e)
Semi-deciduous forest to cropland	144.69	49.95	10.00	0.00	33577.09	10,683,828	17,939.15	5,708,024
Semi-deciduous forest to grassland	144.69	49.95	2.30	6.40	250.14	80,133	719.44	230,479
Semi-deciduous forest to other lands	144.69	49.95	0.00	0.00	0.00	0.00	679.08	227,779
Evergreen forest to cropland	123.13	42.06	10.00	0.00	1878.42	502,364	1,792.63	479,419
Evergreen forest to grassland	123.13	42.06	2.30	6.40	0.00	0.00	0.00	0.00
Evergreen forest to other lands	123.13	42.06	0.00	0.00	0.00	0.00	0.00	0.00
Mangrove to cropland	269.01	85.43	10.00	0.00	98.91	58,709	0.00	0.00
Mangrove to grassland	269.01	85.43	2.30	6.40	0.00	0.00	0.00	0.00
Mangrove to other lands	269.01	85.43	0.00	0.00	0.00	0.00	0.00	0.00
Total					35,804.55	11,325,034	21,130.30	6,645,702

See the workbook template for more details [here](#)

Year of Monitoring / Report Period t	Emissions from deforestation (tCO ₂ -e/yr)	If applicable, Emissions from forest degradation (tCO ₂ -e/yr)*	If applicable, enhanced Removals from afforestation/reforestation (AR) (tCO ₂ -e/yr)	If applicable, enhanced Removals from other activities besides A/R (tCO ₂ -e/yr)	Net Emissions and Removals (tCO ₂ -e/yr)
2021	11,325,034				11,325,034
2022	6,645,702				6,645,702
Total	17,970,736				17,970,736

4.3 Calculation of emission reductions

See the workbook template for more details [here](#)

<i>T.1</i>	Deforestation	If applicable, forest degradation	If applicable, enhanced Removals from afforestation/ reforestation (A/R)	If applicable, enhanced Removals from other activities besides A/R*	Adjustment, if applicable	Total (tCO _{2-e})
Emissions or Removals in the Reference Level (tCO _{2-e})	10,506,534	0	0	0	0	10,506,534
Emissions or Removals under the ER Program during the Reporting Period (tCO _{2-e})	17,970,736				0	17,970,736
Emission Reductions during the Reporting Period (tCO _{2-e})	-7,464,202				0	-7,464,202

5 UNCERTAINTY OF THE ESTIMATE OF EMISSION REDUCTIONS

5.1 Identification, assessment and addressing sources of uncertainty

Table 7: Sources of uncertainty to be considered under the FCPF MF.

Sources of uncertainty	Analysis of contribution to overall uncertainty	Systematic	Random	Contribution to overall uncertainty (High / Low)	Addressed through QA/QC?	Residual uncertainty estimated?
Activity Data						

<p>Measurement</p>	<p>This error represents the operator error during the interpretation of LULCC on sampled points and inconsistencies between operators. This error is reduced by extensive QA/QC procedures.</p> <p>Quality control was guaranteed by having a team of technicians with experience in forests and remote sensing, all trained using the same methodology. The team worked in the same office, and discussed any classification issues with each other. Moreover, specific SOPs were defined in order to ensure the consistency in the interpretations.</p> <p>Quality control was conducted using the SAIKU extension of Collect Earth. This tool allows the detection of whether:</p> <ul style="list-style-type: none"> (i) Data point was not filled (ii) The class assigned followed the classification hierarchy, based on the % of individual element cover (iii) Year of the Old image/Change image was less than the current image (iv) Change classes are consistent with previous and current classes (v) Open and closed forest was correctly classified, based on the 30% (open) and 65% (closed) cover threshold <p>In the case of any error being detected, the ID of the data point was registered and the user performed the necessary corrections.</p> <p>All sampling units detected as deforestation and 20% of the remaining sampling units are subjected to quality assurance (QA). This QA is performed by 2 independent reviewers, who compare their evaluations of each sampling unit, to reach a decision on whether the chose sampling unit was correctly evaluated or not. The critical evaluated parameters, which determine whether a sample has to be reviewed by the user are: land cover class (level 1 and 2), land cover change class and previous land cover class (in case of change). If errors are detected in at least 20% of the reviewed sampling units from the 20% mentioned initially, then the operator has to reanalyze their lot. This process is cyclical, until less than 20% of the sampling units are found to have errors.</p>	<p>YES</p>	<p>YES</p>	<p>H(bias/random)</p>	<p>YES</p>	<p>NO</p>
--------------------	---	------------	------------	-----------------------	------------	-----------

Representativeness	This source of error is related to the representativeness of the estimate which is related to the sampling design. We produce annual deforestation maps as the basis for stratification, to ensure that our sample is representative of the area of interest. We applied a probabilistic-based sampling, where all areas have an inclusion probability larger than zero	YES	NO	L (Bias)	YES	NO
Sampling	<p>Sampling error is the statistical variance of the estimate of area for the applicable forest transitions that are reported by the ER Program. This source of error is random. Mozambique has followed Good Practices regarding estimating the contribution of this error.</p> <p>For the reference period we used systematic sampling, which does not have an unbiased estimator for the variance. The variance estimation formulae for simple random sampling were used as a conservative option.</p> <p>For the monitoring period we used stratified sampling and the method described by Olofsson (2014).</p>	NO	YES	H (bias/random)	YES	YES
Extrapolation	This source of uncertainty is not applicable to our approach. We generate estimates of deforestation per forest type, based on reference data.	YES	NO	N.A.	N.A.	NO
Approach 3	<p>This source of uncertainty exists when there is no tracking of lands or IPCC Approach 3, which is the case for Mozambique. We do not consider that the time-span of the Reference Period + Monitoring Period is sufficient for a land to have been deforested, grown back to forest and then deforested again.</p> <p>With the methodology used in the reference period, it was not possible to double count deforestation events, as we analyzed the entire period. On the other hand, this is a possibility in the monitoring period. Because we are only accounting for deforestation this is conservative with regards to our emissions reductions estimate.</p> <p>Mozambique does not have a clear definition of the time-span required for a land to be considered to have been</p>	YES	NO	H (bias)	YES	NO

	converted “back” to forest after a deforestation event.					
Emission factor						
DBH measurement	<p>Strong QA/QC processes were implemented:</p> <ul style="list-style-type: none"> SOPs were developed as described in <i>Section 2.1 - National Forest Inventory</i>. A training on the SOPs was conducted prior to the field work. This training lasted for 3 weeks, and consisted of training on the usage of all equipment and evaluating the specific skills of each participant, in order to determine the team and brigade leaders. On the start of the 2nd phase of the IFN (2017) an additional 1-week training was conducted, to refresh the participants and train any new members. The supervisor of each inventory team conducted a remeasurement of 4 trees per plot which means 16 trees per cluster. This served to ensure that the SOPs were adequately implemented. An independent measurement of 10% of the plots. Technicians of the National Directorate of Forests, who had participated in the Provincial Inventories of Gaza and Cabo Delgado, conducted this activity. Diameter below 10%. The World Bank conducted two regular supervision missions of the National Forest Inventories to confirm the adequate implementation of the SOPs and suggest areas for improvement. <p>As a result of these QA/QC procedures the possible bias in the measurement of DBH and H have been addressed and the measurement random error is considered to be low. Hence, this source of error will not be propagated.</p>	YES	YES	H (bias) & L (random)	YES	NO
H measurement		YES	YES	H (bias) & L (random)	YES	NO
Plot delineation		YES	YES	H (bias) & L (random)	YES	NO
Wood density measurement	The allometric equations used by Mozambique do not include wood density, so this source of error will not be propagated.	YES	YES	N.A.	N.A.	NO

<p>Other parameters (e.g. Carbon Fraction, root-to-shoot ratios)</p>	<p>Carbon fraction parameter was taken from the 2006 IPCC Guidelines. Error, as provided from the IPCC Guidelines, has been propagated. Sensitivity analysis showed a very small effect of this parameter.</p> <p>Root-to-shoot ratios were used for one of the strata (Evergreen Forest), with the value taken from the 2006 IPCC Guidelines. Within this stratum, we only applied the root-to-shoot ratio to species which were not covered by specific equations, as described in Section 3.1 of this report.</p> <p>Since the previous MR (2018) Mozambique has integrated emission factor estimation within the automated processing chain. As a result, we have propagated Root-to-shoot ratios as per the guidelines.</p>	<p>YES</p>	<p>YES</p>	<p><i>H (bias) & L (random)</i></p>	<p>YES</p>	<p>YES</p>
<p>Biomass allometric equation (Model error)</p>	<p>Allometric equations used ranged from national (specific species, and evergreen MoUntain forest), to regional (for mangrove), international (Semi-deciduous forest) and IPCC defaults (evergreen forests). However, effect on emission reductions is expected to be low, as emission factors remain constant from reference to monitoring period. Additionally, the overall effect of emission factor uncertainty on total uncertainty is low (10.4%).</p> <p>The equations used for semi-deciduous forest and evergreen forest were not validated with data from Mozambique, which is a source of bias. Unfortunately, this was not feasible due to financial reasons. As QA/QC procedure, the selection of the equations was discussed with experts from the Eduardo Mondlane University and IIAM who confirmed that these are the most representative and best available equations, which will provide accurate estimates, as far as practice.</p> <p>According to the experts, although there might be an associated bias from using the equation, it is safer to use the equation of Mugasha et al. 2013 (more representative "ecosystems and species") than using the adjusted equations in Mozambique (less representative "ecosystems and species"). It is because the adjusted equations in Mozambique mostly recommended for specific areas (example of one of the best-adjusted Miombo equation "Guedes et al. 2018" recommended only to estimate biomass in low Miombo of Beira corridor).</p>	<p>YES</p>	<p>YES</p>	<p><i>H (random/bias)</i></p>	<p>YES</p>	<p>YES</p>

	<p>In addition, if they are applicable to extensive ecosystems, they present a high level of uncertainty (example is the equation of Miombo adjusted by Chaúque 2004, which has $R^2 = 0.78$), which is associated with low representation of species and diameter range of the trees used during equation adjustment.</p> <p>On the other hand, Mugasha et al 2013 used data from 60 species (about half of which occur in Zambézia) from 1 to 110 cm of dbh, coming from Miombo woodland (which according to Chidumayo & Gumbo, 2010 "The Dry Forests and Woodlands of Africa", this forest type are similar in terms of floristic composition and structure to those of Mozambique). In addition, the last paragraph of conclusion of the authors' article where they show no reservations about the use of the equation in other regions of southeastern Africa.</p> <p>Currently the MRV unit has plans to establish MoU with research institutions to develop and/or adjust more accurate allometric equations for various ecosystems in the country, and thus update the emission factors.</p> <p>Since the previous MR (2018) Mozambique has included propagation of this source of error in MC simulations for all the strata and pools for which allometric equations are used. As a result the previous application of increased sampling uncertainty of AGB and BGB (of FSD and FSSV forest types) by 10% at 90% confidence level using the quadrature approach has been removed, with the exception of FSSV BGB, which does not have an allometric equation, but rather uses R:S ratio.</p>					
Sampling	<p>Sampling error is the statistical variance of the estimate of aboveground biomass, dead wood or litter. This source of error is random and is considered to be high and it has been propagated.</p> <p>The estimation of mean and their respective uncertainties (standard error, sampling error, and confidence interval) for the variables biomass, carbon and carbon dioxide equivalent (above and below ground) for the two strata (semi-deciduous forest and semi-evergreen forest), were done using the forest inventory data analysis approach proposed by Bechtold & Patterson (2005), as</p>	NO	YES	H (random/bias)	YES	YES

	suggested by the independent expert (Jim Alegria, ex-US Forestry Service) hired to evaluate the methodology for the inventory.					
Representativeness	This source of error is related to the representativeness of the estimate which is related to the sampling design. For semi-deciduous and evergreen forest, data are from the Zambézia Forest Inventory. It includes data that was collected in Zambézia province during the NFI, in 2017 and 2018. Although the inventory covers the whole province of Zambézia, this is still representative of the forests located in the ZILMP as forests across the province are homogenous (floristic and structural composition). Moreover, the higher sample size of the inventory covering the whole province will enable more precise estimates for emission factors. This source of uncertainty is considered to be low.	YES	NO	<i>L (bias)</i>	YES	NO
Integration						
Model error	The combination of AD & EF does not necessarily need to result in additional errors. Usually, sources of both random and systematic error are the calculations themselves (e.g. mistakes made in spreadsheets). The spreadsheets used for activity data and emissions estimation are derived from multiple past implementations and have been refined over several years. The MRV team has implemented an automated script to calculate emissions and uncertainty. This should greatly reduce the possibility of mistakes in the calculations. The outputs of the activity data and emissions spreadsheets were checked against R implementation and they matched. The worksheet for emission factor estimation was developed in consultation with, and checked by, an independent expert (Jim Alegria, ex-US Forestry Service).	YES	NO	<i>L (bias)</i>	YES	NO
Integration	This source of error is linked to the lack of comparability between the transition classes of the Activity Data and those of the Emission Factors. Considering the homogeneity of forests in Zambézia, the distinguishing feature of the two land strata (semi-deciduous and evergreen) are	YES	NO	<i>L (bias)</i>	YES	NO

	<p>the phenological behavior. The <i>Collect Earth</i> software provides a time-series of NDVI over the plot, which is used to determine whether a forest is deciduous or evergreen. More detail of this can be seen in our step-by-step description of activity data collection (https://www.fnds.gov.mz/mrv/index.php/documentos/guioes/46-protocolo-de-monitoria-e-estimativa-de-emissoes-por-desmatamento-vjun2021/file).</p>					
--	---	--	--	--	--	--

5.2 Uncertainty of the estimate of Emission Reductions

Parameters and assumptions used in the Monte Carlo method

Uncertainty in estimates of emission reductions were quantified using a Monte Carlo approach, based on 10,000 random permutations of model parameters (Table 8). The parameter values for AD in the monitoring period are an average of the activity data for 2021 and 2022, as they are calculated in the same way as the reference level AD (sum of area divided by number of years). All Monte Carlo simulation was done using scripts in R (see the full project [here](#))

Several types of Probability Density Functions (PDFs) are used as part of the Monte Carlo simulation. These are:

- t-distribution: Emission factors for FSD and FSSV vegetation classes
- Normal: Emission factors derived from IPCC defaults (cropland, grassland, other land use)
- Uniform: Default root:shoot ratio, for species where local data are not available.
- Non-parametric bootstrapping: Used for activity data.
- Triangular: Carbon fraction derived from IPCC defaults.

In each of these cases, the distributions were selected for their suitability for the data source.

Root to shoot ratio

A uniform distribution is used for estimation of BGB for species where specific local allometric models aren't available (derived from IPCC given the range 0.27 - 0.28). Without further information provided, a uniform distribution was selected for its conservative nature.

Carbon fraction

The triangular distribution used for the carbon fraction was selected to account for the asymmetric nature of the uncertainty range associated with the IPCC default used (0.47 (0.44 - 0.49)). In any case, emissions estimates show very little sensitivity to changes in this parameter (see sensitivity analysis), so it would not be expected that any reasonable alternative PDF would have any impact on overall uncertainties.

Emission factors

FSD/FSSV emission factors use a t-distribution to account for low sample sizes. IPCC tier 1 emission factors are presented with a nominal estimate of error equivalent to two times the standard deviation, for which a normal distribution is considered a reasonable PDF.

Activity data

Uncertainties for activity data were captured using non-parametric bootstrapping, where sample units were resampled (with replacement) from the Collect Earth points. This has the advantage of not needing to specify a PDF a priori, and removing the impact of generating impossible negative areas of deforestation where the uncertainty range crosses 0.

The impact of this decision over two other reasonable approaches (a normal distribution, and a truncated normal distribution removing any negative deforestation areas) was assessed by comparison. In all cases the uncertainty ranges are almost identical, so any reasonable PDF would not be expected to have any impact on overall uncertainty of emissions.

Table 8: Parameter specifications used in the Monte Carlo simulations for the monitoring period.

Parameter included in the model	Parameter values	Error sources quantified in the model (e.g. measurement error, model error, etc.)	Probability distribution function	Assumptions
Carbon fraction	0.47	Measurement	Triangular (lower bound = 0.44, upper bound = 0.49, mode = 0.47)	(IPCC 2006)
Ratio of molecular weights of CO ₂ and C	44/12			Default
Root to shoot ratio	0.275	Measurement	Uniform (lower bound = 0.27, upper bound = 0.28)	(IPCC 2006)
Length of reference period	11 years		-	ER program design
Project area	5310265.16 ha		-	ER program design
Area of FSD>(A O U) in reference period	1600.4 ha	Sampling	Non-parametric bootstrapping	

Area of FSD>C in reference period	129635.8 ha	Sampling	Non-parametric bootstrapping	
Area of FSD>P in reference period	19205.3 ha	Sampling	Non-parametric bootstrapping	
Area of FSSV>C in reference period	35209.7 ha	Sampling	Non-parametric bootstrapping	
Area of FSSV>P in reference period	1600.4 ha	Sampling	Non-parametric bootstrapping	
Area of FSD>C in monitoring period	51516.2 ha	Sampling	Non-parametric bootstrapping	
Area of FSD>P in monitoring period	969.6ha	Sampling	Non-parametric bootstrapping	
Area of FSSV>C in monitoring period	3671.0ha	Sampling	Non-parametric bootstrapping	
Area of FSD>(A O U) in monitoring period	679.1	Sampling	Non-parametric bootstrapping	
FF>C in monitoring period	98.9	Sampling	Non-parametric bootstrapping	
Aboveground biomass of FSD	144.7 t/ha	Sampling	t-distribution (mean = 144.7, sd = 16.33, df = 28.7)	
Aboveground biomass of FSSV	123.1 t/ha	Sampling	t-distribution (mean = 123.1, sd = 10.73, df = 5.2)	
Aboveground biomass of FF	269 t/ha	Sampling	Normal distribution (mean = 269, sd = 27.03)	
Aboveground biomass of C	10 t/ha	Sampling	Normal distribution (mean = 10, sd = 3.75)	
Aboveground biomass of P	2.3 t/ha	Sampling	Normal distribution (mean = 2.3, sd = 0.86)	
Aboveground biomass of (A O U)	0 t/ha	Sampling	Normal distribution (mean = 0, sd = 0)	
Belowground biomass of FSD	49.9 t/ha	Sampling	t-distribution (mean = 49.9, sd = 4.98, df = 25.99)	
Belowground biomass of FSSV	42.1 t/ha	Sampling	t-distribution (mean = 42.1, sd = 3.29, df = 4.01)	

Belowground biomass of FF	85.4 t/ha	Sampling	Normal distribution (mean = 85.4, sd = 10)	
Belowground biomass of C	0 t/ha	Sampling	Normal distribution (mean = 0, sd = 0)	
Belowground biomass of P	6.4 t/ha	Sampling	Normal distribution (mean = 6.4, sd = 3.9)	
Belowground biomass of (A O U)	0 t/ha	Sampling	Normal distribution (mean = 0, sd = 0)	

Quantification of the uncertainty of the estimate of Emission Reductions

		Reporting Period	Crediting Period
		Total Emission Reductions	Total Emissions Reductions
A	Median	-7,359,743	-10,899,741
B	Upper bound 90% CI (Percentile 0.95)	-4,584,818	-1,907,332
C	Lower bound 90% CI (Percentile 0.05)	-10,486,811	-22,102,494
D	Half Width Confidence Interval at 90% (B – C / 2)	2,950,996.5	10,097,581
E	Relative margin (D / A)	40%	93%
F	Uncertainty discount	8%	12%

5.3 Sensitivity analysis and identification of areas of improvement of MRV system

Sensitivity analysis was conducted by setting one parameter at a time to its nominal value, while retaining uncertainty of all other parameters generated from Monte Carlo (Table 9). The major contributor to uncertainty of ERs was Activity data for the monitoring period, followed by activity data for the reference period. Uncertainty from emission factors and carbon fraction was negligible.

This is the first MR where uncertainty from AD of monitoring period has a greater contribution than that of the reference period. This is due to the much greater magnitude of the AD for the monitoring period (i.e. there were significant reversals), compared to the reference period.

Table 9: Sensitivity analysis for the monitoring period.

Sensitivity test	Uncertainty estimate					Reduction in confidence interval (%)
	Median	Lower bound (5th percentile)	Upper bound (95th percentile)	Half-width confidence interval at 90%	Relative Margin	
Nominal	-7,359,743.43	10,486,811.16	4,584,817.61	2,950,996.77	0.4	0
AD (reference)	-7,340,034.77	10,005,642.70	5,039,117.17	2,483,262.76	0.34	15.9
AD (monitoring)	-7,380,552.91	9,534,097.34	5,398,720.89	2,067,688.22	0.28	29.9
EF AGB	-7,381,904.75	10,110,487.27	4,797,301.62	2,656,592.82	0.36	10
EF BGB	-7,365,103.76	10,461,370.68	4,602,028.75	2,929,670.96	0.4	0.7
CF	-7,412,635.11	10,535,789.01	4,642,740.43	2,946,524.29	0.4	0.2

6 TRANSFER OF TITLE TO ERS

6.1 Ability to transfer title

In Mozambique, the main legal and regulatory frameworks concerning to the land and forests that support the Program Entity ability to transfer title to ERs are: **The Constitution of the Republic of Mozambique (CRM, 2004), the Law on Forests and Wildlife (1999), the Land Law (1997) and the REDD+ Decree (2018)**. The REDD+ Decree provides all the principles and procedures to be respected for the design and implementation of the ER Program. It deals with, inter alia: (i) the institutional framework, which is greatly clarified; (ii) the process for the approval and issuing of licenses for projects involving carbon credits and the procedures for the approval of REDD+ projects, putting great emphasis on community consultations; (iii) **establishes the uncontested ownership of ER titles to the State of Mozambique**; and (iv) **details administrative procedures for the management of the ER Transactions Registry and the REDD+ Project and Data Management Registry**.

In Mozambique, **Carbon is a State property** - Carbon is a constituent element of forests. If carbon is seen a constituent part of all natural resources, which exists per se, current constitutional and sectorial legislation is adequate for establishing that ownership over carbon resides with the State. The starting point is Article 98 of the CRM, of which the clause 1 clearly states: "Natural resource in the soil and the subsoil, in inland waters, in the territorial sea, on the continental shelf and in the exclusive economic zone shall be the property of the State". In addition, Article 102 of the CRM goes on to say that "The State shall promote the knowledge, surveying and valuing of natural resources, and shall determine the conditions under which they may be used and developed subject to national interests".

The concept of "use and development" of natural resources - The intention of the Constitution in this overall context is clear: **the State as owner shall determine how natural resources are "used and developed" and, further, this determination can include selling the natural resource once it has gone through this process of "use and development"**. In other words, the carbon can be sold if it is subject to some sort of **conversion or transformation into a marketable commodity**. In the specific context of natural forests, which are State property, and which are in the public domain, the key legislation is the 1999 Forest and Wildlife Law (Law 10/99), which gives mandated agencies in the Government the right to assess requests to "use and develop" natural resources.

ERs are products of "use and development" of carbon natural resources - Precisely, ERs can be seen as a product of this "use and development" process. ERs are not a natural resource, as opposed to carbon: they are the outcome of a decision by the State and/or others with rights over natural resources, and can only be produced by a transformational process or action implying to reduce deforestation and forest degradation. As such, they could be considered as "environmental commodities", identifiable and marketable in their own right. As a consequence, the CRM and existing natural resources laws are sufficient for determining ownership of ERs through the application of the "use and development" concept: the "user and developer" of the natural resources (in this case, forest carbon stocks) implements activities that result in ERs being produced.

Until recently, State ownership of ERs was only clearly established by law for those generated within conservation areas. Although this right seems clearly established for conservation areas such as Gilé National Park where, in principle there will be few, if any, other pre-existing rights or claims over the resources in question, this may not have been true for other types of areas. In this situation, potential claims of rights on the ERs could have led the GoM to negotiate partnership or intermediation agreements with potential DUAT holders. Given the unfamiliar nature of the carbon and ER issues, it was therefore forecasted that specific legislation could greatly clarify the question of title and ER sales.

The REDD+ Decree clearly establishes State property on all ER generated in the country (Articles 4 and 6): although non-state DUAT holders and communities will have to benefits from the sale of ERs generated in the country, through specific benefit sharing plans, no formal agreements will need to be reached between each individual DUAT holders or local communities and the State. However, they will have to be properly consulted, as per national law. In order that the process has been implemented, taking into account national legislation, several meetings have taken place, between 2018 and 2019, from where 564 individuals participated in 6 consultation events at national, provincial and district level. The main objectives of these consultations were to discuss the program approach, the percentages of benefit allocation to each group of beneficiaries, allocation

models/processes, priorities areas and benefits sharing challenges of the Benefit Sharing Plan (BSP). For further details of public consultations, please see on the following site below¹²:

As such, the REDD+ Decree clarifies the **“legitimacy and ownership of the State in the creation, generation, emission, validation, verification and withdrawal of emission reductions and corresponding titles of emission reductions” (Article 4)**. As such, in the current ER Program in Mozambique, the State retains control over the remaining natural forests and ownership over the ERs that are generated and the GoM, promoting behavioral change on the part of forest users, and is therefore free to sell the titles over these ERs, following the arguments presented above. Furthermore, the ability of the State of Mozambique to dispose of ER titles as financial products that can be traded is established in the REDD+ Decree, which states that ER titles **“may be disposed of, transferred to national and international exchanges of environmental and financial assets, under the applicable laws and standards and within the limits of the current national legislation”** and that such ER titles **“may also be transferred and offset in future under the international agreements concluded by the State of Mozambique within the framework of its international competences and its commitments and cooperation programs with public and private entities” (Article 15)**. In the same way, Article 7 of the REDD+ Decree confirms that, for the implementation of REDD+ programs and projects, **“The government can sign compensation agreements with international partners”**.

Admittedly, the overall ability of the State to transfer the titles over ERs requires these ERs to be monitored, reported, verified and certified accordingly with UNFCC procedures and FCPF CF methodological guideline. The discussion of certification and negotiations underlines how the Ministry of Economy and Finance (MEF) is really the entity able to enter into international negotiations over ER titles transfers, whenever the ERs are generated. As stated in the REDD+ Decree, **“The Ministry responsible for the financial sector is the legitimate issuer and manager of the Titles of Emission Reductions, being able to create and manage property rights, including the validation, verification, emission, transfer, transaction and withdrawing of the titles of emission reductions at national and international level” (Article 6)**. In the context of the ER Program, the MEF was therefore the ER Program entity authorizing the ER Program and signing the ERPA with the FCPF CF. According to the administrative and legal procedures, the title of ERs is registered and ERs certificates issued by the MEF, after validation and verification of the monitoring report, provided by FNDS. Until now, MEF has not ER Transaction Registry established. However, FNDS is committed to working with the MEF, this year, in order to speed up the process of registering transactions. **As such, the MEF will be responsible the sale of ERs to the Carbon Fund.**

This REDD+ Decree clarifies the institutional arrangements for the implementation of REDD+ projects in Mozambique and clearly specifies the responsibilities of the FNDS and other key institutions. The institutional arrangement for the ER Program will fully respect the layout describes in the REDD+ Decree. According to the REDD+ Decree, The Ministry of Economy and Finance (MEF) is responsible for signing the Emission Reduction Payment Agreement (ERPA) with the FCPF CF ERPA and management of ER Titles transfer. FNDS will work closely with the MEF after the verification process, in order to provide technical support on this process.

Prior to the establishment of ZILMP, there was a VCS REDD project, called the Gilé National Reserve REDD Project, which was developed in the buffer zone of the Gilé National Park¹³, Zambézia Province (Figure 12). This Park is managed by the National Administration of Conservation Areas (ANAC). It is a national public agency that is responsible for the management of protected areas, and the project proponent of the above REDD Project.

This project was originally designed to have a crediting period of 20 years, from January 1, 2012, until December 31, 2031. However, once the ZILMP project was being developed, it became clear that, since the Gilé National Reserve REDD Project was fully included in the ZILMP program area, it would no longer be able to generate Carbon Credits from the date of the start of the ZILMP project. This is made clear in the project’s PDD¹⁴. The project was successfully validated for the monitoring period of 01-January-2012 to 31-December-2016. It is

12

<https://docs.google.com/spreadsheets/d/1LOo1dvQyUOXMHOU20Djg61E3ECM7OgDbJEhf68jPZ5c/edit?usp=sharing>

¹³ Previously it was known as the Gilé National Reserve

¹⁴ Sections 1.8.5 and 1.12.4 of joint PD & monitoring report (doc. Ref. 1)
<https://registry.verra.org/app/projectDetail/VCS/1674>

currently inactive for the duration of the ZILMP ERPA. This project is registered in the (<http://bit.ly/registryredd>). Several meetings between ANAC and FNDS have occurred in order to prevent double counting and a conflict of interest between the two REDD Programs.

Other than the Gilé National Park REDD Project, the program has not become aware of an inability or any contesting party during this reporting period, also there has not been any challenge, no one disputing the REDD+ decree and no title contested".




Figure 12: Location of Gilé National Reserve REDD Project Area (buffer zone of Gilé National Park) and Zambézia Emissions Reduction Program Area.

6.2 Implementation and operation of Program and Projects Data Management System

The National Fund for Sustainable Development (FNDS) will be in charge of supervising and coordinating the ER Program at central level. In the REDD+ Decree, the FNDS is confirmed as the entity in charge of approving all REDD+ programs and projects in Mozambique and in charge of managing REDD+ resources. However, since the creation of the Directorate of Climate Change in (DMC) the MTA in 2019, most of the duties of FNDS are now performed by the DMC. This includes approval of expressions of interest for implementation of REDD+ projects, official communication with REDD+ project proponents and submission of license requests to the Minister of Land and Environment. FNDS provides technical assistance to the DMC, when requested and manages the REDD+ Project Registry. As clarified in the REDD+ Decree (Article 10), the FNDS supports all institutions engaged in REDD+ policies. Its main responsibilities are:

- a. Establish, operationalize and ensure the maintenance of the components of the National MRV System;
- b. Propose and approve standards and technical methodologies for establishing the levels of reference, the monitoring, the evaluation of emission reductions, the reporting, the verification and the validation of REDD+ programs and projects;

- c. Receive, assess and evaluate the REDD+ projects proposals and annual monitoring reports;
- d. Monitor the reduction of greenhouse gas emissions and the achievements of ERs objectives of REDD+ projects;
- e. Management of the Safeguards Information System (SIS), including the REDD+ Feedback and Grievance Mechanism (FGRM);
- f. Enable the dissemination of data and relevant information on REDD+ projects, which should be made public respecting the policies of intellectual property privacy established with the different actors; (vii) To disseminate all information on the Programs and Projects and their social and environmental safeguards, Dialogue Mechanism and Complaints on existing platforms and their benefit sharing plan. With regard to the ER Program, the FNDS will therefore play a crucial role in the monitoring of the ERs generated by the ZILMP and of the safeguard policies - see section 14. In addition, and importantly for the ER Program, as stated in the REDD+ Decree (article 10) the FNDS is responsible for
- g. Managing the national REDD+ Programs and Projects Data Management System and for
- h. Communicating to the entity in charge of the ER Transactions Registry all information related to ERs generated by REDD+ projects – this is the MEF.

According to the REDD+ Decree (article 10), the **FNDS will be responsible for managing the national REDD+ Programs and Projects Data Management System** and for communicating to the entity in charge of managing the ER Transactions Registry (who will be the MEF, according to the same decree – Articles 14 and 26) all information related to ERs generated by REDD+ projects, including by the Zambézia Emission Reduction Program. After the creation of DMC, communication with MEF is done by DMC, while FNDS still manages the national REDD+ Programs and Projects Data Management System.

Mozambique is developed and implement its own comprehensive national REDD+ Program and Projects Data Management System. The system is hosted and managed by FNDS as per de REDD+ decree “the FNDS is responsible for (vi) managing the national REDD+ Programs and Projects Data Management System and for (vii) communicating to the entity in charge of the ER Transactions Registry all information related to ERs generated by REDD+ projects”. Currently the system is implemented through a WebGIS platform (<http://bit.ly/registryredd>) alongside with the NFMS and the projects M&E Web portal. The system is still under development and improvement will be included over time, as currently Mozambique only has one ER program.

The actual Content of the REDD+ Program and Project Data Management System is below:

- The proponent of the ER Program or project;
- Geographical boundaries of the ER Program or project;
- Scope of REDD+ activities and Carbon Pools;
- The Reference Level used;
- MRV data to specific REDD+ projects/programs; and
- Safeguards plans in specific REDD+ projects/programs

The National REDD+ Program & Project Data Management System Manual has been finalized. The document went through a consultation process with relevant stakeholders. The finalized version is the one currently available on the MRV platform at the link previously shared <https://www.fnds.gov.mz/mrv/index.php/documentos/guioes>

Please note that, while the Manual has been finalized, may be updated or improved over time to incorporate lessons learned and new developments in the operation of the REDD+ Program & Projects Data Management System.

6.3 Implementation and operation of ER transaction registry

As mentioned at 6.1, in this report, only after the approval of the REDD + decree in 2018, this is the first program to be implemented in the country. For this reason, it is still preparing to implement and operationalize the registration of ER transactions for future programs. Thus the GoM has decided to use a centralized ER Transaction Registry managed by a third party on its behalf: **the GoM will use the CATS Transaction Registry.**

6.4 ERs transferred to other entities or other schemes

The Zambézia Emission Reduction Program is the first REDD+ program that occurs in Mozambique, after the approval of the Monitoring report and according with Contract ER, the volume will be transferred to the FCPF CF on a 100% basis. No ERs will be transferred to other entities during the crediting period.

As mentioned in section 6.1, there is a prior REDD project within ZILMP, but it is currently inactive and only generated Emission Reductions from 1 January 2012 to 31 December 2016.

7 REVERSALS

7.1 Occurrence of major events or changes in ER Program circumstances that might have led to the Reversals during the Reporting Period compared to the previous Reporting Period(s)

A study carried out in the program area shows that shifting cultivation is the main cause of deforestation, contributing to more than 80% of the total deforestation in the program area. Of particular importance is an increase in the production of pigeon pea inside and outside the program area to later be exported to Asia through local intermediaries. The results obtained during the study by the MRV unit are available here:

<https://storymaps.arcgis.com/stories/8ffacb098c97469e897a94cd596da01a>

It is also speculated that the COVID-19 pandemic triggered a series of socioeconomic impacts that may have contributed to the increase in deforestation in vulnerable communities. Economic restrictions, lack of supervision, displacement of workers and the search for alternative sources of income may have led to deforestation practices for subsistence, which possibly contributed to an increase in deforestation in reporting period: 2020, 2021 and 2022, but with a small downward trend in 2022 (year in which COVID-19 pandemic restrictions reduced).

This possible impact of COVID-19 is supported by anecdotal evidence collected in the field, specifically during the country visit in the previous verification report. We interviewed farmers who had recently opened new agricultural areas around the city of Mocuba and at least one reported that they had lost their income in the city as a result of COVID-19 impacts.

7.2 Quantification of Reversals during the Reporting Period

See the workbook template for more details [here](#)

A.	ER Program Reference level for this Reporting Period (tCO₂-e)	<i>from section 4.1</i>	10,506,534	
B.	ER Program Reference level for all previous Reporting Periods in the ERPA (tCO₂-e).	<i>from section 4.1 of previous ER Monitoring Reports</i>	15,759,804	+
C.	Cumulative Reference Level Emissions for all Reporting Periods [A + B]		26,266,338	

D.	Estimation of Emissions by Sources and Removals by Sinks for this Reporting Period (tCO ₂ -e)	<i>from section 4.2</i>	17,970,736	
E.	Estimation of Emissions by Sources and Removals by Sinks for all previous Reporting Periods in the ERPA (tCO ₂ -e)	<i>From section 4.2 of previous ER Monitoring Reports</i>	11,613,545	
F.	Cumulative Emissions by Sources and Removals by Sinks including the current Reporting Period (as an aggregate accumulated since the Crediting Period Start Date) [D + E]		29,584,281	–
G.	Cumulative quantity of Total ERs estimated including the current Reporting Period (as an aggregate of ERs accumulated since the Crediting Period Start Date) [C – F]		-3,317,943	
H.	Cumulative quantity of Total ERs estimated for prior Reporting Periods (as an aggregate of ERs accumulated since the Crediting Period Start Date)	<i>from previous ER Monitoring Reports</i>	4,146,258	–
I.	[G – H], negative number indicates Reversals		-7,464,201	
If I. above is negative and Reversals have occurred complete the following:				
J.	Cumulative quantity FCPF ERs (as an aggregate of FCPF ERs accumulated since the Crediting Period Start Date)	<i>from previous ER monitoring reports, section 8</i>	2,524,661	

K.	Cumulative ER Program’s Pooled Reversal Buffer contributions (as an aggregate of Pooled Reversal Buffer ERs accumulated since the Crediting Period Start Date)	<i>from previous ER monitoring reports, section 8</i>	1,359,433
L.	Cumulative ER Program’s Uncertainty Buffer contributions (as an aggregate of Uncertainty Buffer ERs accumulated since the Crediting Period Start Date)	<i>from previous ER monitoring reports, section 8</i>	262,163
M.	Cumulative ER Program’s Pooled Reversal Buffer replenishments (as an aggregate of Reversal Buffer ERs replenished since the Crediting Period Start Date)	<i>from previous ER monitoring reports, section 7.3</i>	0
N.	Cumulative amount of FCPF ERs, Uncertainty and Pooled Reversal Buffer contributions and replenishments (as an aggregate since the Crediting Period Start Date) [J + K + L + M]		4,146,257
O.	Quantity of Buffer ERs to be canceled from the Pooled Reversal Buffer account [If I < N, report the value of I; if I > N, report the value of N]		4,146,257

7.3 Quantification of pooled reversal buffer replenishments

Intentionally left blank

7.4 Reversal risk assessment

Risk Factor	Risk indicators	Default Reversal Risk Set-Aside Percentage	Discount	Resulting reversal risk set-aside percentage
Default risk	N/A	10%	N/A	10%
Lack of broad and sustained stakeholder support	<ul style="list-style-type: none"> ● Existence of a transparent Benefit Sharing Mechanism ● Existence of legal mechanism for the systematization of community consultation ● Signature of MoU with implementing partners ● Existence of a Feedback and Grievance Redress Mechanism during the ER Program implementation, likely to generate the implementation of long-term efficient practices beyond the project life time ● Existence of consultative forums and platforms involving various stakeholders with concrete and immediate perception of benefits, likely to make consultation become a long-term concern (including out of the scope of the ER Program) ● Implementation of an efficient and large enough land titling and delimitation process to ensure stability of land rights in the long run 	10%	Reversal risk is considered High: 10% discount	10%
Lack of institutional capacities and/or ineffective vertical/cross sectorial coordination	<ul style="list-style-type: none"> ● Existence of designated and empowered relevant structure for ER Program implementation ● Experience in multi-sectorial project implementation ● Experience of collaboration between different levels of government ● Existence of dedicated mechanism or body for inter-sectorial cooperation ● Support from additional projects and programs for institutional capacities strengthening ● Deployment of relevant staff on the ground ● Training for long-term capacities on forest management and monitoring 	10%	Reversal risk is considered Medium: 5% discount	5%
Lack of long term effectiveness in addressing	<ul style="list-style-type: none"> ● Experience in decoupling deforestation and degradation from economic activities 	5%	Reversal risk is considered	5%

underlying drivers	<ul style="list-style-type: none"> ● Support from completing projects and programs oriented on deforestation and forest degradation reduction ● Existence of a relevant legal and regulatory environment conducive to REDD+ objectives in the long run ● Creation of relevant incentives for adoption of sustainable agricultural practices in the long run, including beyond the project lifetime ● Clear perception of non-carbon benefits for stakeholders at long term and especially beyond the terms of the ERPA ● Deployments of efficient and committed extension-agents at long-term ● Adaptation of promoted sustainable practices to local constraints and dynamic in order to make it possible for them to be maintained in the long run ● Potential administrative changes are expected to be progressive and participatory. However, potential risk may exist due to the fact that the ER program area doesn't cover the whole Province and additional coordination might be required. ● Well defined structures to ensure ensures the continuation of the ER Program beyond government term ● Pre-identification of financing sources 		High: 0% discount	
Exposure and vulnerability to natural disturbances	<ul style="list-style-type: none"> ● Vulnerability to fires, storms and droughts ● Capacities and experiences in effectively preventing natural disturbances or mitigating their impacts ● Promotion of climate smart agricultural practices ● Existence of a Pest Management Plan 	5%	Reversal risk is considered High: 0% discount	5%
Total reversal risk set-aside percentage			35%	
Total reversal risk set-aside percentage from ER-PD or previous monitoring report (whichever is more recent)			35%	

8 EMISSION REDUCTIONS AVAILABLE FOR TRANSFER TO THE CARBON FUND

See the workbook template for more details [here](#)

Year			2021	2022	TOTAL
A.	Emission Reductions during the Reporting period (tCO ₂ -e)	<i>from section 4.3</i>	(6,071,767)	(1,392,435)	(7,464,202)
B.	If applicable, number of Emission Reductions from reducing forest degradation that have been estimated using proxy-based estimation approaches (use zero if not applicable)		0	0	0
C.	Number of Emission Reductions estimated using measurement approaches (A-B)		(6,071,767)	(1,392,435)	(7,464,202)
D.	Percentage of ERs (A) for which the ability to transfer Title to ERs is clear or uncontested	<i>from section 6.1</i>	100%	100%	100%
E.	ERs sold, assigned or otherwise used by any other entity for sale, public relations, compliance or any other purpose including ERs accounted separately under other GHG accounting schemes or ERs that have been set-aside to meet Reversal management requirements under other GHG accounting schemes	<i>from section 6.4</i>	0	0	0
	If applicable, any buffer replenishments	<i>section 7.3 P</i>	0	0	0

F.	Total ERs [(B+C)*D-E] minus, if applicable, any replenishments as per section 7.3, P		(6,071,767)	(1,392,435)	(7,464,202)
G.	Conservativeness Factor to reflect the level of uncertainty from non-proxy based approaches associated with the estimation of ERs during the Crediting Period	<i>from section 5.2</i>	8%	8%	8%
H.	Quantity of ERs to be allocated to the Uncertainty Buffer (0.15*B/A*F)+(G*C/A*F)		(485,741)	(111,394)	0
I.	Total Reversal Risk set-aside percentage applied to the ER program	<i>from section 7.4</i>	35%	35%	35%
J.	Quantity of ERs to be allocated to the Pooled Reversal Buffer (F-H)*I		(1,955,109)	(448,364)	0
K.	Number of FCPF ERs (F-H – J)		(3,630,917)	(832,677)	0
L.	Percentage of Emission reductions from enhanced removals from afforestation/reforestation as a percentage of the total FCPF ERs [Optional if the country wishes to generate enhanced removals]	<i>From section 4.3</i>	0.00000000%	0.00000000%	0
M.	Number of FCPF ERs from enhanced removals from afforestation/reforestation (L * K) [Optional if the country wishes to generate enhanced removals]		0	0	0

ANNEX 1: INFORMATION ON THE IMPLEMENTATION OF THE SAFEGUARDS PLANS

Environmental and Social Aspects of ER Programs

The implementation of safeguards within the scope of ERPD complies with World Bank (WB) guidelines that are aligned with UNFCCC guidance related to REDD+. Some principles are defined as guidelines for compliance with the implementation of the project's environmental and social safeguards, respectively:

- (i) Compliance with legislation and good governance;
- (ii) Promotion of transparency and public/social responsibility;
- (iii) Respect the local culture and customary norms;
- (iv) Ensure meaningful participation by affected people and stakeholders (especially the most vulnerable);
- (v) Ensuring the existence of “auscultation” forms as conflict resolution mechanisms;
- (vi) Protect and conserve forests, contribute to the improvement of multiple forest functions;

In Mozambique the REDD+ readiness process went hand in hand with the preparation of the Social and Environmental Strategic Assessment (SESA).

Since 2015, the GoM is piloting a large-scale landscape program in part of the Zambézia province – the Zambézia Integrated Landscape Management Program (ZILMP) aiming to address the drivers of deforestation and degradation through an integrated landscape management approach. The landscape management approach recognizes the link between agricultural development, natural resource management and governance, both in terms of institutional management and practical implementation.

The present document is to provide evidence to demonstrate that the investment activities generating the ERs (2021-2022) meet the WB environmental and social safeguards. These investment activities are all financed by the following three World Bank investment projects:

Forest Investment Project (MozFIP P160033, effective in August 2017) and closed on June 2022.

Agriculture and Natural Resources Landscape Management Project (Sustenta P149620, effective Nov 2016)

Dedicated Grant Mechanism for local communities (MozDGM P161241, effective Feb 2018 and will be closed on Feb 2023).

Mozambique Sustainable Rural Economy Project (MozRural P174002, Effective June 2021 and will be closed on June 2026).

The above 3 investment projects represent the only REDD+ activities within the ER program area and they have the following safeguard instruments:

Annex-Table 110: List of safeguard instruments approved and in place (Documents available in the link: <https://www.fnds.gov.mz/index.php/pt/documentos/salvaguadas-artigos>)

Projects	Instruments	Duration
MozFIP	ESMF and PF Addendum ¹⁵	2017 – 2022
MozDGM		2017 – 2023
ANRLMP (Sustenta)	ESMF, RPF and PMP	2015 – 2023

¹⁵ These two instruments were prepared to support the implementation of the MozFIP, MozDGM and other REDD+ initiatives and were based on the findings gathered during the elaboration of the SESA and the National REDD+ Strategy, financed by the Mozambique FCPF REDD+ Readiness Preparation Support (P129413).

MozRural	2021 - 2026
----------	-------------

All of the above-mentioned instruments have undergone consultations (summary of consultations available as annexes to the reports) and have been properly disseminated at central, provincial and local/district levels and are available online on both the FNDS website and the World Bank info shop.

In addition to these instruments, a Sustenta GBV Action Plan, a land protocol and a protocol for preventing critical habitat conversion have been adopted to help address any potential land conflicts in land tenure regularization and unintended conversion of critical habitats respectively. Protocols available on the website: <https://sismdr.fnds.gov.mz/biblioteca>.

Effectively addressing the issues and complaints of individuals or groups affected by project activities is an essential component of operational risk management and FCPF requirements. Grievance redressing mechanisms are a way to prevent and resolve community concerns, reduce risk, and support processes that create positive social change.

In order to ensure a transparent and participatory process that contributes to the mitigation of risks related to possible conflicts in the implementation of REDD+ initiatives, FNDS developed a common Grievance Redress Mechanism called the “Dialogue and Grievance Mechanism” (MDR) for the projects that make up the Bank’s Integrated Landscape Management Portfolio in Mozambique (MozFIP, MozBio, and Sustenta). It is to mention that MozDGM has a similar system that it has been managed by WWF safeguard specialist. It is a system created to answer questions and complaints from individuals or groups affected by the project and program activities. Among other uses, it serves as an instrument to search for a harmonious relationship between affected and interested parties in the areas of implementation of projects and programs. The safeguards assistants based in the PIU is the focal point for the FGRM in Zambézia province, and therefore responsible for receiving, processing, investigating and responding to the grievances reported in this province. The safeguards team prepared a manual of procedures, a communication strategy and a monitoring system for tracking grievances and assessing the extent to which progress has been made to resolve them, have already been prepared. The information about compliance stage can be found at the following link: <http://sismdr.fnds.gov.mz/fnds/>

Both mechanism were discussed with key stakeholders, including local communities, and are being implemented successfully.

Simultaneously, there is a continuous process of consultation and engagement involving different actors through the Zambézia Multi-Stakeholders Landscape Forum (MSLF) and in District, local level. Information on different phases and levels of consultations¹⁶, where is highlighted the meetings with the Development Platform in the context of the preparation of the Benefit Sharing Plan and safeguard instruments is further described in the following link: <https://docs.google.com/spreadsheets/d/1LOo1dvQyUOXMHOU20Djg61E3ECM7OgDbJEhf68jPZ5c/edit#gid=862126948>

Monitoring and Reporting Requirements

FNDS has developed monitoring and reporting procedures and templates that were deemed acceptable by the WB and used in the projects that make up the Bank’s Integrated Landscape Management Portfolio referred above. Monitoring consists of field visits and generation of quarterly reports submitted to the World Bank. The landscape safeguard team, supported by FNDS central safeguard team are responsible for this activity. In addition, the WB carries out supervision visits and Mid-Term Reviews every six months.

To effectively guarantee transparency, information is available, accessible and disseminated among the stakeholders. Specifically, for the Zambézia ER Program the most used way to share information including opportunities to participate in project activities, consultations and project performance results has been the Zambézia Integrated Development Platform, (PDIZ)

¹⁶ The BSP team undertakes to submit this information within the monitoring report for the year 2021

Investment activities compliant with the environmental and social safeguards

The three WB projects in the program area include the following activities: (i) Integration of rural households into sustainable agriculture and forest-based value chains, (ii) restoring degraded areas, (iii) land delimitation and tenure regularization, (iv) establishment of new planted forests/ the Planted Forest Grant Scheme (EFF)¹⁷, and (v) promotion of agro-forestry systems.

The activities are implemented by service providers and monitored by the PIU (Project Implementation Unit) that has a team of experts and extensionists, including a safeguards specialist. The Provincial Government (SPA and DPDTA – The Environment Provincial Service and the Provincial Directorates for Environment and Territorial Development) are involved in the environmental assessment process of each subproject as per the national Environment Impact Assessment (EIA) Regulations. The district government is involved also in the supervision of civil works, verifying compliance with labour and health and safety contract requirements. While, for implementation of MozDGM activities, the National Management Committee (CNG) was created and the World Wild Life Fund for Nature (WWF) was hired as a National Executing Agency (NEA), with the responsibility of increasing the participation of local communities and CBOs in Integrated Landscape Management; strengthening capacities for Community-Based Natural Resource Management; and Managing, monitor & evaluate the project.

Thus, proponents of community projects were supported in the improvement and detailing of the pre-selected community projects and carried out environmental and social screening of the 18 selected community projects.

All WB investment activities undergo a review and screening process to determine the level of environmental and social assessment required. The screening and project categorization phase determines the necessary type of environmental and social management instrument to be developed for each activity, namely whether a Simplified Environmental Impact Assessment (ESIA), an Environmental and Social Management Plan (ESMP) or an Environmental and Social Management Good Practices Guide (ESMGPG) is required. Annex-Table 211 shows the list of activities implemented in the period of January 1st to December 31st, 2021 in the program area, highlighting the status, including the categorization and the instrument prepared.

It is important to note that the majority of activities, mainly Agro-forestry systems (AFS), Forest Plantations, Agriculture (small emergent commercial farms) started in 2021, and the number of beneficiaries are increasing over the years.

All screening and licensing processes go through the validation of the WB safeguards team as a way to ensure compliance with their guidelines.

Annex-Table 211: Environmental and social screening and safeguard plans prepared (January 1st January 2019 to December 31st, 2021-2022)

Subproject	Category	Instrument	Implementation phase
MozFIP			
Agro-forestry systems (AFS) in 42 communities in Zambézia Landscape	C	ESMGPG	Updated Obtained 42 environmental licenses from SPA
Forest Plantation (EFF) (104 Beneficiaries: SMEs – Small and Medium Entrepreneurs)	C	ESMPGPG	Obtained 104 Environmental Licenses and elaborated 40 ESMP ¹⁸

¹⁷ EFF acronym in Portuguese.

¹⁸ <https://fnds.infordata.co.mz/biblioteca>

Sustenta			
Agricultural development (75 SECF- Small emerging commercial farmers)	C	ESMGPG	Ongoing and elaborated a Agriculture and SAFs ESBPM for the farmers
MozDGM			
Participation of local communities and OCB in Management Integrated Landscape (18 Community projects selected and screened)	C	ESMGPG	17 project have been screened, resulting in development of Environmental and Social Management Plans (PGAS) for 9 projects (Anawapé, ACODEMUSO, ADETUR, CHICOPERA; CAMBORA, and 6 Cashew nuts value chain projects for Zambézia and MAKHELELO, CGRN-UNIDADE and 3 horticulture value chain projects in Cabo Delgado) and Best Practice's Manuals (MBP) for 8 projects (OPURELA, APROCAMO and EGUMI DIOROMA in Zambézia and poultry consortium of Montepuez, Salaua, Meza and ORERA Association in Cabo Delgado). Ongoing

Parallel to the screenings, the process for environmental licensing of sub-projects is instructed in compliance with national legislation. The forest plantation sub-projects were classified as category C, with this oriented towards the elaboration of a Good Practices Manual which was harmonized with the Forest Management/Management Plan. As a result of the process, 10425 environmental licenses were issued for a total of 29 beneficiaries. The remaining 4 being in process.

Screening of new areas for the 2020-2021 campaign began in the last quarter of 2020. Particularly for the FPS there is an attempt to involve young people, who can collaborate with large companies (Chazeira Mugoma,

Chazeira Mozambique). To date, we have identified 10 young people to join the EFF, 10 potential trainees from Unizambeze and 2 potential nurseries.

PFGS engaged 3 (three) educational institutions that planted from 2021 to 2022 a total of 38 ha from 2019 to 2021, with the participation of youth. 10 (Ten) Uni-Zambeze interns were involved in PFGS assisted activities, 11 youth planted 61 ha, women and youth in nurseries, planting and restoration activities. Women were involved in almost all forestry activities namely field preparation, planting, plantations maintenance, nurseries through associations and as employees. This positive involvement was achieved thanks to the great contribution of the women empowering program (GALS).

The safeguards team produced a very simple and illustrative Environmental and Social Management Good Practices Guide for SECFs and AFS including basic environmental and social mitigation measures (Available in the link [https://fnfs.infordata.co.mz/biblioteca:"Guia de boas práticas para agricultura e sistemas agroflorestais "](https://fnfs.infordata.co.mz/biblioteca:). 200 guides on good practices (ESMGPG) in agriculture and agroforestry systems were distributed to extension workers, community facilitators, SDAE's (District Service for Economic Activities) technicians, and the same guides are being distributed to beneficiaries gradually.

Safeguard Compliance Levels

Overall, the supervision and technical missions to the WB projects have systematically found that the measures identified by the safeguards instruments (ESMF, RPF, PF and PMP) have been properly implemented to prevent, minimize and mitigate environmental and social risks and impacts in a manner that is satisfactory to the World Bank Safeguard Policy. This is evidenced in particular by implementation of key and innovative tools:

- (i) A generic Good Practice Manual (GPM) for emerging commercial agricultures and agroforestry systems producers;
- (ii) A protocol to avoid conversion of critical habitats;
- (iii) A land protocol;
- (iv) A protocol for community readiness;
- (v) COVID-19 Prevention Protocol, and
- (vi) An online system to track the progress of safeguards documents processes was established, but the FNDS projects have faced constrains due to COVID-19 restrictions and deployment of staff which resulted in delays in submitting safeguards sub-instruments. Therefore, the team has been quick to react and improve compliance (i.e. submission of timely quarter reports and regular coordination meetings with the WB).

During this period (2019-20) an independent third party carried out a mid-term environmental audit, that was planned in the project preparation. The safeguards execution audit process to assess the level of performance of this component was completed and the audit report produced. The document presents the level of safeguards execution, which considers the beginning of the project and based on the instruments approved and aligned with the financing agreement between the Mozambican Government and the World Bank, and presents recommendations for improving the implementation of safeguards within the scope of the project.

The report presents a positive assessment of 82% Compliance out of a total of 34 requirements assessed, which includes performance in sub-project licensing, monitoring processes, Grievance Redress Mechanism, Pesticide management etc. In terms of training, the assessment was satisfactory with 63%. See the complete report on <https://www.fnfs.gov.mz/index.php/pt/component/edocman/relatorio-de-auditoria-de-salvaguardas-ambiental-e-social-mtr/download>.

The document was shared with the BM, Service Providers, PIU's and other implementing partners such as AQUA. This document is available on the FNDS website.

It was confirmed during project supervision missions that no works have started without ESMPs or Good Practice Manual (GPM) and that the environmental and social information and clauses are being included in the bidding documents and contracts, respectively. Despite the efforts made to comply with the WB safeguards policies, there are always challenges, especially in view of the quantity, variety and location – remote and disperse – of most subprojects. Thus, it is still necessary to continuously strengthen the capacity of FNDS to monitor the

implementation of ESMP, Pest Management Plan (PMP) and GPM by the contractors, service providers and beneficiaries.

The issue of safeguards and REDD+ are still new in the landscape. The main challenges are related to the low level of literacy of most beneficiaries, which requires constant awareness to engage in the fulfilment of safeguards. Another challenge is to identify and use the clearest and simplest language possible.

The safeguards issue has been introduced and trained from the beginning of the presentation of the project on the Zambézia Integrated Development Platform, (PDIZ), beneficiaries and district government.

Setting up safeguards for service providers and training extensionists has been the most used way to overcome the challenge. To this end, the FNDS team annually develops and implements a safeguards training program for government (SDAE- District Service for Economic Activities, SDPI- District Service for Planning and Infrastructure, DPDTA - Provincial Directorate of Environment and Territorial Development, SPAE- Provincial Services and SPA – Provincial Services for environment, for Economic Activities partners (RADEZA, CESC, R-GCRN) including contractors and service providers. The training program is very practical, focusing on improving the capacities to properly implement the safeguard instruments.

Under MozFIP activities, during the period 2021 individuals were trained, with 2000 participants¹⁹. The trainings were about several themes and approaches focusing on inductions on environmental and social safeguards, Grievance Redress Mechanism, land conflict management and prevention of conversion of critical habitats; COVID-19 Protection, Gender Action Learn System – GALS, which aims to support beneficiaries and their households to remove gender barriers that prevent joint planning and use of resources and in <https://mozdgm.org.mz/>. Under SUSTENTA, 3 training sessions were held in various matters of socio-economic progression. Involved in GALS training were the safeguards team at the central FNDS level and PIUs, community facilitators, service providers currently following up with beneficiaries; Identification of gender barriers in the management of natural resources (Survey models); Restoration; Use of organic pesticides (developed by the service provider/NIRAS and applied to the beneficiaries of SAF's); and Techniques for planting and fighting uncontrolled fires.

Under SUSTENTA, 3 training sessions were held in various matters of socio-environmental safeguards, both for the technical team at the central level and PIUs, and 3 inductions were carried out for the beneficiaries of the subprojects (startups, MSMEs, PACEs). Monitoring visits were carried out to 75 PACEs and 4 MSMEs.

Tree visits were carried out, in April, June and October 2021, in order to assess the level of implementation of the activities planned for this year, assess the projections of the District Governments, discuss the Benefit-sharing Plan, participatory MRV and in particular the local initiatives to reduce deforestation in the Zambézia landscape, as well as discuss the Benefit Sharing Plan, which culminated with the launch of the BSP call, on the 23rd of November, for the financing of CBOs and Companies.

The Annex-Table 313A public presentation of the current version of the Benefit Sharing Plan was held at the III Plenary Session of the Zambézia Platform, on December 13, 2019. 85 participants were present, 71 members and 14 guests, out of which 23 were women.

Although, every effort undertaken with all stakeholders during the different trainings there are non-compliance activities to be reported, as indicated in the table below.

19

https://docs.google.com/document/d/1wR7objAss4Axy7L6DeBGGm038DCPAvp0/edit?usp=share_link&oid=115860222606486742127&rtpof=true&sd=true

Annex 1- Table 12 Non-compliance activities under implementation of different projects in Zambézia Landscape

Aspect	Non/compliance activities	ZILMP/BSP	All other Projects
Economic	<ul style="list-style-type: none"> • Introduction of plants/crops reduce yield among them, due to competition of sunlight, water and nutrients; • Difficulties in seedling production and access to organic seeds. 	<ul style="list-style-type: none"> • ✓ • ✓ 	<ul style="list-style-type: none"> • ✓ • ✓
Social	<ul style="list-style-type: none"> • Influx of communities to conservation areas or its buffer zones (case of PNG); • Losses of land tenure, due to familiar problems (i.e. death of one of the spouses); • Lack of completion and preparation of a management plan for the parcels/machambas 	<ul style="list-style-type: none"> • ✓ • ✓ 	<ul style="list-style-type: none"> • ✓ ✓
Environmental	<ul style="list-style-type: none"> • Reduction of biodiversity due to implementation of “commercial” agroforestry (i.e. plantation of cashew trees); • Transformation of pristine forest (non-secondary) to agriculture which lead to, among others, biodiversity losses and transmission of disease to society; • Lack of physical barriers to control external vectors and wind actions in the parcels (Contamination); • Weak adequacy of environmental aspects in plots/machambas, such as adoption of MBPAS 	<ul style="list-style-type: none"> • ✓ • ✓ • ✓ • ✓ 	<ul style="list-style-type: none"> • ✓ • ✓ • ✓ • ✓

below illustrates the summary information of the safeguards action in the ongoing activities.

Annex-Table 313: On the ground safeguards activities implemented in the Zambézia Landscape.

Activities	Safeguard Action on the Ground	Compliance
<p>Agroforestry System (Gurué= 17 Communities, Mulevala= 16 Communities e Mocubela= 9 Communities. In all of these communities have Environment License- 1,610 farmers were involved in a total area of 1.035ha.). Of this</p>	<ul style="list-style-type: none"> • Mapping and ground-truthing sensitive habitats for identifying protection and restauration activities (critical habitats, riverine forest, water springs, steep slopes, etc.); 	<p>In progress</p>

<p>number 34% are women, with 583 in the three districts.</p> <p>12 contracts were established with people living in, (districts of Mocuba – 3; Gurue- 4, Mulevala-4 and 1 outside Zambezia province) and distributed about 152,155 seedlings (12,079 fruit trees, 106,054 nitrogen fixers, 15,534 native tree species and 18,488 others species).</p>	<ul style="list-style-type: none"> · Surveillance and monitoring the cleaning of farm fields (“machambas”) to ensure no uncontrolled cutting of trees (preservation deforestation, protect red list species and some forest species at risk e.g. umbila/wild teak - <i>Pterocarpus angolensis</i>, black wood- <i>Dalbergia melanoxylon</i>). 	
<p>Forestry Plantations (EFF) Small nurseries produced in 2021/2022 a total of about 1332.98 ha of commercial plantations, 202.05 ha of restoration through enrichment and 157 ha of regeneration management were planted, benefiting about 196 plantation owners being [5 SMEs, 77 individuals, 19 associations being (3 with collective DUAT) and (16 which held individual DUAT and with a total of 108 members), 3 educational institutions).</p> <p>During the process of plantations establishment, a total of 1,533,143 seedlings were distributed and planted. Of these, 1,463,069 seedlings were from exotic species of which 1,314,717 <i>eucalyptus spp</i> were supplied by Portucel and 139,352, <i>Eucalyptus spp</i>, and 9,000 of <i>Acacia mangium</i> by TFCA.</p>	<ul style="list-style-type: none"> · Mapping, protection/recovery of about 10 hectares of critical habitats (riverside forest, water bodies, distinct erosion; native forest plots with some status of degradation...) and 90 hectares for the EFF (continuous process). To this end, restoration by stakes has been tested, with native species of umbila (<i>Pterocarpus angolensis</i>) and chanfuta (<i>Afzelia quanzensis</i>); · Quarterly monitoring visits to plantation areas for supervising and assessing the performance of safeguard aspects linked to grant payments 	In progress
<p>Sustainable Agriculture (113 Emerging Small Commercial Farmers - PACE's and 25 Youth PACE's with 14.080 Small farmers - PA's supported)</p> <p>75.000 seedlings were distributed in the districts of Alto Molocue, Gilé, Gurué and Mocuba, to restore the area of 97 PACEs.</p>	<ul style="list-style-type: none"> · Training and raising awareness on environmental and social good practices to beneficiaries; 	In progress
	<ul style="list-style-type: none"> · Training on pesticide use and awareness for the correct disposal of out of date pesticides including support for final disposal of pesticide packaging from suppliers.). The training includes Integrated Pest Management (IPM) techniques: (i) Biological Control (ii) Cultural and Crop Sanitation Practices (iii) Chemical Control 	
<p>Community projects (inputs distributed and technical assistance provided to about 1,300 community members in the buffer zone of the Gilé National Park – tree owners, beekeepers, charcoal makers and small farmers</p>	<ul style="list-style-type: none"> · Raising awareness and environmental education programmers carried out on sustainable natural resource management involving more than 10,000 community members; 	In progress

	<ul style="list-style-type: none"> · Production and dissemination of documentaries and songs on conservation of forest resources · Identification and preservation of 6,000 trees, in particular <i>Fabacea</i>, 	
<p>Community organization: Community Creation and legalization of 14 CBNRM organizations in the buffer zone of the Gilé National Park was supported and 2 forest management plans were elaborated which includes the identification of the potential of non-timber forest products for communities in Gilé and Mulevala districts;</p> <p>5 CBNRM supported to elaborate Project Business Plans for community natural resources management projects in Mulevala, Gilé, Mocubela, Mocuba and Maganja da Costa.</p>	<ul style="list-style-type: none"> · Training and raising awareness on environmental and social aspects; · Reinforcement of gender mainstreaming and participatory governance · Training on the operationalization of the Dialogue and Grievance Mechanism and dissemination of spots through community radios of Mocuba, Gilé and Maganja da Costa and placement of giant posters. 	In progress
<p>Land Tenure Regularization (187 communities were delimited; 162 Community Certificate issued; 144 Community Land Use Plans elaborated; and 153 Community Development Plans (Agenda) elaborated; 17189 titles in 2021, and 83353 cumulative titles issued until December 2021)</p>	<ul style="list-style-type: none"> · Field visits and interviews to monitor social community preparedness and the creation of community committees; 	
	<ul style="list-style-type: none"> · Site observation of the level of community engagement in the elaboration of community maps through Participatory Rural Approach (PRA); 	In progress (finalize and hand over the land titles and community certificates)

During the period corresponding to this report there was no record of new environmental and social risks.

Grievance Redress Mechanism in place and fully operational

A Grievance Redress Mechanism called the “Dialogue and Grievance Mechanism” that is a common mechanism for the projects that make up the World Bank Integrated Landscape Management Portfolio in Mozambique (MozFIP and Sustenta). The system suffered a breakdown in the first quarter and was improved, and introduced three green lines, one for each regional zone of the country (South, Center and North), and obtained 15 lines to operationalize the mechanism. The Annex-Table 4 below summarize the registered cases from January to December 2021 and January to December 2022 in the Zambézia landscape.

Annex-Table 4: Summary of conflicts in the Zambezia Landscape, 2021-2022

Event summary	2021	2022
Registered cases	65	43
Claims	42	33

Suggestions	19	10
Consultations	19	10
Land conflicts	6	0
Not proceeding	0	0
Need clarification	4	0
Resolved	28	20
Ongoing	14	1
MozFIP	43	8
SUSTENTA	22	35
Project Performance	39	18
Social	14	14
Environmental	2	11
Not Classified	4	0

Additional information is available on the website: <http://196.22.53.245:8080/fnds/estadistica>. The Complaints were about land conflicts, in particular about the land tenure regularization process (individual titles and community certificates). Definition of limits was the main issue raised by project beneficiaries. Other complaints are related to delays in the delivery of agricultural inputs to SECF (Small Emerging Commercial Farmer) and AFS (agro forestry systems). Complaints about environmental issues are related to poor protection of sensitive habitats (riparian areas) and lack of knowledge about pesticide handling.

Regarding the management of land conflicts, in 2021-2022, 6 conflicts were registered in the conflict management system of the Service Provider for Land Regularization in the Zambézia Landscape of which all of them were resolved. Six conflicts were presented by Verde Azul (Service Provider) for registration in the MDR, out of which 6 corresponded to the Zambézia Landscape, that required the attention of the safeguards team and were all resolved.

The conflict resolution system considers 3 levels of resolution: (i) between the parties involved with support from the local community leadership, local government, service provider and landscape safeguards (ii) between the parties involved with the support of FNDS safeguard specialist at central level (iii) between the parties involved with support of independent mediator Specifically, land conflict issues, when not resolved at community level with support of the FNDS, the SPGC and the conflict management office of the national land directorate are invited to support.

Until December 2022 almost 100% of the conflicts were resolved at local level with the support of community leaders, Landscape Safeguards and service providers.

To add to the above, there are traditional mechanisms for resolving complaints through local authorities (Community leaders). The MDR only responds to complaints about projects and community leaders are the main collaborators in resolving these conflicts.

Efficient and well-trained safeguard team

FNDS is responsible for overall strategic guidance and implementation of three of the four WB-funded projects (MozFIP and Sustenta). The World Wildlife Fund for Nature (WWF) implements MozDGM.

Capacity building and training for Project Implementation Unit (PIU) was crucial in order to identify potential impacts of the project and determine appropriate environmental and social category of the subproject during screening phase. In this regard, in December 2021, in Nampula, the training of 20 PIUZ technicians took place, by the FNDS Central team, in induction on safeguard policies and instruments. Also in 2022, the PIU technical team were trained to screening for BSP project.

FNDS has been accumulating experience and expertise in managing World Bank funded operations, particularly in climate change and natural resources management sectors. FNDS has the human resources - a team of four safeguards specialists at national level and one provincial safeguards specialist in the PIU in Zambézia province assisted by 1 community officer from Gilé National Park, 30 extensionists (agricultural value chains) and it is in the process of hiring 20 extensionists (15 to AFS and 5 to Forest plantation) - to oversee the MozFIP and Sustenta – and sufficient project funds to adequately implement the safeguards instruments, including the GRM, and the monitoring and reporting framework.

MozDGM aims to increase the participation of communities and community-based organizations in landscape management, and to build their capacity to prepare and manage their own projects. Communities are receiving trainings to improve the quality and frequency of their participation in local decision-making bodies, as well as in technical areas to support sustainable resource management practices. WWF manages all project activities, including safeguards screening, implementation and monitoring. Their safeguards specialist is receiving safeguards training from the World Bank. FNDS provides support to project oversight as a member of the DGM National Steering Committee.

The safeguards team with the support of the WB safeguards specialist has been preparing safeguards training program for the PIU staff, extensionists, and service providers to ensure that all the stakeholders involved in the subprojects received safeguards training before starting the implementation of project activities. In addition to these trainings, project beneficiaries participated in specific trainings on environmental and social good practices and the GRM.

The training plan is semi-annual; however, trainings are also carried out when individuals are hired to collaborate with the projects. All capacity building activities include, but are not limited to, the following: (i) training on the Environmental and Social Standards; (ii) GRM; (iii) Environmental and Social screening; (iv) Health and Safety; and (v) Pest Management Plan.

The beneficiaries of the training were NIRAS - Service Provider for the promotion of agroforestry systems and forest plantations; the community associations of Uapé and Nipiode; the beneficiary communities of the forest promotion scheme and agroforestry systems; the government representatives through District Permanent Secretaries, SPA, DPADT and SDAE's technicians.

Specifically, in 2021, the training of FNDS technicians in the context of strengthening institutional capacity stood out:

One (1) capacity building session carried out in collaboration with the WB which involved, in addition to the FNDS safeguards team, community development technicians from the CAs and 1 technician from MozDGM;

Two (2) training courses led by the BM with the objective of ensuring the knowledge of the project implementing partners on the new safeguards' standards adopted by the BM, specifically the Environmental and Social Standards, designated as "NAS", from 1-20 April;

Several training and awareness-raising sessions for the project's beneficiaries regarding COVID-19;

Introduction to the GALS methodology - Gender Action Learning System. In terms of the results achieved, we are currently working with 42 CGRN, 6 Associations and a total of 2,310 beneficiaries, of which 47% (1,093) are women. The GALS is being developed through 3 officers from the GALs, 2 community development officers from the PNAG and 1 from the NIRAS-EFF. And it is used for several levels.

20 CBOs (16 CGRN and 4 community associations) were mapped, identified and selected to benefit training with a CESC. And on March 31, 2021, the Zambezia Landscape CBOs Training Program was launched in Mulevala District, with the presence of the District Administrator;

Sensitized 390 members of the 20 beneficiary CBOs about the rights and duties of communities in relation to land and other natural resources. And 588 members of the CBOs were trained in terms of community

governance, a package led by the representative of the Community Governance Network of Natural Resources (R-GCRN).

Safeguards are a crosscutting issue and aspects of safeguards have been included in training on charcoal production, conservation agriculture and Non-timber forest products. These trainings benefited more than 1900 direct beneficiaries mainly in the buffer zone of the Gilé National Park and in the districts of Pebane, Mocubela and Maganja da Costa. The awareness on safeguard aspects included 12 schools in the buffer zone of Gilé National Park.

Particularly for the IPM, the focus is based on the increase capacity to extensionists and farmers, encouraged to work together to make experiments and come up with combinations that are suitable for the area.

The IPM considers: (i) Biological Control that involves the use of biological agents and predators to control pests and diseases. The method was successful in crops like cassava. (ii) Cultural and Crop Sanitation Practices was improved, the farmers usually implement crop rotation and mulching technics; (iii) Chemical Control - These measures involved the use of insecticides to manage weeds, pests and diseases. For chemical control, systemic and contact insecticide - Belt (Flubendiamide 480g / L) 10 ml ampoules (48% Concentrated Suspension) has been used for an area of ½ ha, and for sesame the project has used/ Cyperimetrina 5-10% (250 ml bottle), also for an area of ½ ha.

In the forest plantation activity, there are some termites and the termicides fipronil and vega have been used to protect the plantations.

The IPM includes a control of birds, mainly done by using the traditional way of scaring (the use of scarecrows is very common especially in cereal production areas), chasing and guarding of animals.

SIS on the right track

The online platform for the Safeguards Information System (SIS) was developed and is operational. It is in the public domain and provides general information on safeguards, specifically, the Dialogue and Complaints Mechanism (number of registrations, stage of attendance, category, gender discrimination); existing operational instruments and information on the 7 Cancun safeguards in the Mozambican context for the forest program by providing information on how safeguards are being addressed and respected throughout the implementation of REDD+ activities. The SIS is simple, accessible, auditable, comprehensive and in line with national legislation and applicable Word Bank and donor requirements; and oriented towards transparency and inclusion. The implementation of the SIS is taking a gradual and participatory approach. It is still an incipient process in Mozambique that requires well-structured coordination to enable the full participation of stakeholders (community, private sector, government, civil society).

The online platform for safeguards information in line with national legislation and the World Bank and donor requirements presents the following.

“Complementarity or consistency with national forestry programs and relevant international agreements” – MozBio, MozFIP and Sustenta subprojects aiming at the integration of rural households into sustainable agriculture and forest-based value chains, restoration of degraded areas, establishment of nurseries and new planted forests and promotion of agro-forestry systems in the Gilé National Park and its buffer zone are in line with the Sustainable Development National Program (2015-2030). It should also be noted that the implementation of AFS and the establishment of new-planted areas are key elements in the Action Plan of the REDD+ National Strategy, Reforestation National Strategy (2006-2026) and the Policy of Forestry and Wildlife Development (Resolution n.8/97 of April 1st). The operationalization of the protocol for preventing critical habitat conversion is an approach to comply with the national policy of biodiversity conservation and the convention of biological diversity of which Mozambique is signatory.

Transparent and efficient national forestry governance structures - FNDS is supporting the biannual assessment of forest operators to ensure their activities comply with national logging rules and propose correction measures when necessary. In addition, the four WB projects are supporting the establishment and strengthening of the Communities Based in Natural Resource Management (CBNRM) committees. These committees are community organizational structures aiming at the sustainable management of natural resources. MozFIP started strengthening capacity building for sustainable forest management in two community forest concessions in Gilé

and Mulevala districts. MozFIP and Sustenta projects are supporting the CBNRM committees in the delimitation process and MozDGM plans a capacity-building program for communities in the ERPA area.

Respect for the knowledge and rights of local communities - All initiatives implemented consider respect for local customary norms and habits. National legislation through Land Law and its regulation (Law 19/1997 of 1 October and Decree 66/99) and Forest and Wildlife Law (Law 10/99 of 7 July and Decree 12/2002 of 6 June) mention aspects to be considered in order to ensure respect and rights of local communities throughout the process of forest exploration and utilization particularly when referring to private sector investments for forest concessions or other investments that require the right to use and benefit from land. In the induction training to extensionists, service providers and all technical staff who have direct contact with the local communities, special attention is given to “community relations” including aspects to be respected in the communication with this target group, including best places to hold meetings, norms and customary habits.

Full and effective participation of stakeholders, in particular local communities: Effective participation in REDD + initiatives needs strong information and awareness raising campaigns, public consultations for subprojects and specific training programs. The landscape has an Integrated Development Platform where sustainable and integrated landscape development models and initiatives are discussed and harmonized among different stakeholders (representatives of local communities, civil society, local NGOs, provincial and district government) in regular meetings²⁰. The GRM is also an instrument that contributes to the full and effective participation of program participants.

Consistency with natural forests and biological diversity governance: The forest policy is currently being updated to improve the sustainable management of forest. Conversion of native forests to any activity, whether forest plantations, agriculture or other initiatives, is not eligible for receiving funding or grants by the projects in this landscape and the conversion of any natural habitat is strictly prohibited. The critical habitat conversion prevention protocol was developed to guide this particular aspect.

Actions to address risks of reversals: The adoption of new conservation agriculture techniques and the implementation of agroforestry systems allow the farmers to remain in the same area for several years. It is expected that, the landscape approach to integrated development contribute to the harmonization of sustainable practices of different landscape initiatives coordinated by different stakeholders.

Actions to reduce emissions displacement”- Monitoring and follow-up of activities is carried out with the support of the MRV unit which produces maps of landscape deforestation leading the different actors and stakeholders to know the deforestation stage particularly in their jurisdictional area and buffer zone. In addition, extensionists and the technical team of specialists from the project implementation unit at local level to ensure technical assistance and consequently the productivity of both small farmers and SAF producers carry out regular field visits. Land tenure regularization through DUATs titling is a model adopted in the landscape to ensure farmer presence in the same area. All initiatives are carried out with the support of the district government, which contributes to raising awareness and other aspects at community level. According to the information produced by the MRV unit, it can be seen that between 2017, 2018, 2019-2020 deforestation both in the area of implementation of the project and in the surrounding area has reduced. (Maps and additional information can be accessed through the link <https://storymaps.arcgis.com/stories/13866c3247764d47975e411663eaa810>)

²⁰ The BSP team undertakes to submit this information within the monitoring report for the year 2021

ANNEX 2: INFORMATION ON THE IMPLEMENTATION OF THE BENEFIT-SHARING PLAN

I. Requirements of FCPF on Benefit Sharing Plans

In 2018, the advance draft of the BSP was prepared but no payment was received to finance the implementation of BSP activities. The BSP was approved in February 2020. Between 2021 and 2022 individuals participated in consultation events at national, provincial and district level. The main objectives of these consultations were to discuss the program approach, the percentages of benefit allocation to each group of beneficiaries, allocation models/processes, priorities areas and benefits sharing challenges of the BSP. For more details, on how the activities are organized, please refer to the action plan at the end of the Annex 2.

The main modifications and updates in the approved version include:

- *Budget*: an increase in the operational costs to strengthen the project management team and include funds for supporting the implementation and monitoring of field activities.
- *Beneficiaries*: a modification regarding communities' eligibility was considered necessary. In the advance draft, only delimited local communities with forest cover were eligible. This criterion was considered too restrictive as less than 200 communities out of 1,700 would fulfil this condition by the end of the program. It was then decided that all communities within the ER project area were eligible. Concerning the forest cover, the 9 target districts have forest cover to a greater or lesser extent.
- *Flow of funds to beneficiaries*: this section was improved providing more details on how the funds will be transferred to the beneficiaries including the channels, allocation modalities and responsible entities.
- *Eligibility criteria and distribution modalities*: two main modifications were made in the allocation to communities. First, the allocation to communities continues to be performance-based but the approach has changed. In view of the large number of communities within the ER project area, it was not considered feasible to allocate funds to all the communities in the districts that achieve ER. The best option discussed during the consultations to benefit communities with sufficient funds to implement economic and social activities was the call for proposals. Second, the allocation of 10% of the community share to community capacity building to ensure communities acquire the necessary capacities to change their behavior and implement adequate land use practices to reduce deforestation and achieve targeted emission reduction. The approved BSP provides guidelines to implement these two new implementation modalities.
- *Monitoring*: a section on monitoring of ER program activities was added which includes responsibilities, reporting and financial audits.
- *Consultations*: this section was updated to include the consultations that took place in 2018 and 2019 and a table with the syntheses of the discussions.
- *Implementation guidelines*: an Annex was added with the implementation guidelines for the call for proposals component.

https://drive.google.com/file/d/1wEEo1LpE-JOTT68IRTI61ysoU8KfNVdR/view?usp=share_link

https://drive.google.com/file/d/1b2KRHivVnQXxyb5CRy_-Vntak5JIZGs1/view?usp=share_link

According to the detailed communication plan that was elaborated to ensure that the BSP implementation is widely disseminated for local communities in the field, several actions²¹ will be carried out, such as spots and radio plays; Conducting interviews with program technicians, in local languages; production of booklets with figures; Interpersonal approach carried out by technicians from FNDS and SDAEs. Production of short videos with testimonials from beneficiaries. These videos will be disseminated through digital platforms, such as the FNDS website, social networks and at community meetings, in order to share examples of neighboring beneficiaries. The latest version of BSP booklet is available on the FNDS website.

II. Monitoring and Reporting Requirements

²¹ All materials will be translated into local languages!

1. Benefit Sharing Plan Readiness

- 1.1 The final version of the BSP was approved in February 2020. Between 2018 and 2020, draft versions were discussed with relevant stakeholders in several forums at national, provincial and district level. In 2018, a total of four consultations were carried out in the framework of the preparation of the BSP, respectively: (i) at the multi-stakeholder landscape forum (MSLF) of Zambézia, two meetings involving representatives of local government, communities, private sector, NGOs; (ii) at the central level one meeting with government representatives in the sectors of Land, Conservation Areas, Agriculture, Forestry, Mineral Resources and Energy, Rural Development, State Administration and representatives of the World Bank and; presentation during the V National Conference of Community Management. In the events 445 individuals participated, 30% of them were women, for more details, please see the link below²². The main objective of the meetings was to discuss the percentages of benefit allocation, allocation models/processes, priorities areas and benefits sharing challenges of the BSP. In 2019, two consultations took place. In November, the BSP was presented to civil society organizations with the following objectives: 1) To reach consensus on improving mechanisms for channeling and utilizing community benefits through access to and exploitation of natural resources; 2) To consider the impact of benefits already channeled to local communities and approaches to their improvement and 3) To share experiences among different actors in the local development process with the involvement of local communities and existing potentials. The allocation of 70% to local communities was highly appreciated by the participants. In general, civil society members showed interest and availability for building the capacities of local communities. In December, the approved version of the BSP was discussed with MSLF in Zambézia Province. The main issues discussed were related to i) the selection of proposal based on good practices in terms of governance and the sustainable use of natural resources by communities and CBOs, ii) starting date for ER payments and iii) sustainability of the ER Program. The approved version of the BSP was posted on [FNDS website](https://www.fnds.gov.mz/index.php/pt/component/edocman/plano-de-partilha-de-beneficios-erpa/download) and the FCPF website: https://www.forestcarbonpartnership.org/system/files/documents/bsp_final_jan_2020_clean.pdf. The Portuguese version is available at the FNDS site. Brochures in Portuguese have been provided since June 2022, and are available on <https://www.fnds.gov.mz/index.php/pt/component/edocman/plano-de-partilha-de-beneficios-erpa/download> in order to ensure that all beneficiaries in particular local communities will be informed.
- 1.2 In addition to consultation meetings, where there was always an introduction to the meaning of REDD+ initiatives and what they represented for maintaining the role of forests in the conservation of soil, water and NTFPs and in improving the living conditions of communities, there were also 5 seminars, 2 in 2018 and 3 in 2019, with the aim of working on the annual plans with the provincial and district government. More details, can be seen in the link below²². Capacity building is a continuous process and therefore the main stakeholders will continue to be trained in order to improve the effectiveness of the BSP implementation. Furthermore, the BSP recognizes that communities need to acquire the capacities to change their behavior and implement adequate land use practices to reduce deforestation and achieve targeted emission reduction. The training will be based on issues related to the consolidation of governance systems, including own organization and strengthening in the sustainable management of natural resources and local business development. In order to reinforce the Communities in strengthening their governance and business development capacities. On the BSP action plan at the end of the Annex 2, several activities related with capacity building are previewed and Mulevala District was the first where CBOs were trained in 2020 and 2021. To this end, 10% of the community share (70% of the ER net payment) will be allocated to community

22

<https://docs.google.com/spreadsheets/d/1LOo1dvQyUOXMHOU20Djg61E3ECM7OgDbJEhf68jPZ5c/edit?usp=sharing>

capacity building and a Service Providers are in process of contracting for building the capacities of local communities in the program area.

1.3 During implementation until 2022, it was necessary to update the BSP document. This update was carried out in 2023 and is awaiting comments and subsequent approval.

2 Institutional Arrangements

2.1 The institutional arrangements for the implementation and coordination of the BSP builds on existing mechanisms and respects the role and function of all institutions involved in REDD+ and forest management in Mozambique. The overall governance of the BSP includes: the Ministry of Economy and Finance (MEF); the Bank of Mozambique; FNDS (currently under the Ministry of Agriculture and Rural Development - MADER), including the Matching Grant Unit composed by a team of 5 members at central level supported by the Sustenta Project. The Investment Committee is an independent body constituted and operational to respond to all projects being implemented in the FNDS. Its main function is to deliberate on the submitted proposals. It consists of 6 members that include: a) Matching Grant Scheme (MGS) manager (no voting rights); b) academy representative; c) representative of SMEs; d) representative of the business community; e) Government representative; and representative of the financial sector. At Province level FNDS has the Projects Implementation Unit in Zambézia province, until 2022 was composed by 33 members. At Provincial Directorate of Land Development and Environment of Zambézia (*Direcção Provincial de Desenvolvimento Territorial e Ambiente da Zambézia – DPDTAZ*), we work with 2 technicians, 1 from the planning sector and 1 from the Forest Department and at the Provincial Service for the Environment (*Serviço Provincial do Ambiente – SPA*) level, we work with 2 technicians. In addition, 18 planning technicians of the SDAE and SDPI of each district and 2 from Gilé National Park are also involved. Finally, the Zambézia Multi-Stakeholders Landscape Forum (MSLF) composed by 70 members that belong to several public, private, NGOs, Academia, Social communication and community institutions. These implementing entities are resourced by several projects to carry out their respective responsibilities. Despite the activities being conducted by the above-mentioned institutions, it is necessary to strengthen their skills in order to support the BSP implementation as well as to reinforce the ground team. More details can be seen on the BSP action plan at the end of Annex 2.

2.2 No regulatory or administrative approvals is required for implementing the BSP.

2.3 BSP stakeholders understand their obligations, roles and responsibilities. FNDS held at least 2 missions per year for each of the projects that contribute to the generation of ERs since 2018, and kept all relevant stakeholders aware of the Program and the BSP. In fact, there is great expectation of all stakeholders to receive the payment, since they know that deforestation rates have reduced in the program area. In addition, FNDS prepared the communication materials (brochures, posters, radio spots) and disseminated the BSP implementation modalities. The timeline of the communication steps is indicated on the BSP action plan at the end of Annex 2. In order to ensure the incorporation of messages about BSP in the activities of projects implemented in the landscape, several materials are ready (Alignment with technicians from PIUs (including extension technicians), SDAEs and Service Providers). The detailed communication plan was reviewed and approved, different communication forms are envisaged for the different stakeholders²³.

2.4 A Financial management assessment was undertaken by the World Bank, to evaluate the adequacy of the proposed project financial management arrangements, due to nature of this operation, to in particular determine whether FNDS – the Recipient’s - Dedicated Payment System, is acceptable to the Bank and to determine whether it has the adequate internal controls and oversight mechanism (e.g. audit) in place. The

²³ <https://www.fnds.gov.mz/index.php/en/our-projects/project-list/redd>

assessment revealed that there are adequate financial management arrangements at FNDS for the implementation of the project, specifically the management of the Dedicated Payment System. The assessment concluded that there is adequate capacity at FNDS which was established over the time on implementation of the Bank-financed operations. FNDS has experience in managing Bank-financed operations, and it is currently managing the following projects: Conservation Areas for Biodiversity and Development – Phase 2 (168802), Mozambique Forest Investment Project (P160033), Mozambique Agriculture and Natural Resources Landscape Management (P149620). However, the entity's capacity will be reassessed during implementation support missions and strengthened as needed. No major Financial Management issues were raised under these projects. The overall Dedicated Payment System was assessed to be acceptable to the Bank, and the risk rating was assessed as Substantial due to country fiduciary risk, capacity issues in the country, in particular at provincial level, and decentralized nature of Dedicate Payment system. The first ER payment was received in August 2021. Funds started to be disbursed in 1st quarter of 2022. For the BSP funds, the State's Administration and Financial System is being used. The first BSP financial reports are available on the link!

https://drive.google.com/file/d/1PR7qNq1FdOg7TCM01mBbRWICBG-2QiaZ/view?usp=share_link

- 2.5 The following accountability mechanisms are in place and functional: An internal financial audit still in progress. The report will be available for the next BSP report.
- 2.6 The Feedback and Grievance Redress Mechanisms²⁴ (FGRM) is operational. This system records and addresses feedback and grievances related to the WB projects, such as MozFip, that contributed to emissions reduction in the Zambézia Integrated Landscape Management Program (ZILMP). The registered cases were not related with BSP. For more details about registered cases between January 2021 to December 2022, please see <http://sismdr.fnds.gov.mz/fnds/estatistica>. The FGRM, is a system created to answer questions, clarify issues, and complaints from individuals or groups affected by the activities under the program. The FGRM also serves as an instrument for pursuit of harmonious relationship between the parties concerned and interested in the areas of implementation of projects and programs. The information about registration and complaints attendance is available in an online platform and is updated continuously. Individuals and communities who believe they are adversely affected by the community and private sector initiatives implemented by the BSP will also use this mechanism. In 2021 to 2022, please see above the Annex 1-Table 4 all figures related to WB project activities were received and all of them were satisfactorily resolved. This mechanism will be used for the BSP. Additional information on the FGRM²⁵.

In addition to the FGRM, there is also a continuous process of consultation and engagement with involvement of different actors through the Zambézia Integrated Development Platform. For more details, please see Annex 1.

- 2.7 FNDS has a team in place that is responsible for the ER program as well as for the BSP. At the national level will have a REDD+ focal point, MRV technicians, 1 safeguard specialists At provincial level, 1 BSP provincial coordinator, 1 safeguard specialist and 1 safeguard assistant, 1 administrative and financial assistant, 1 communication officer and 1 business development officer which are in process of contracting for the period of 2022/-2023. Operational costs for running the ER Program and its BSP are estimated to reach 500,000 USD per year. This amount is deducted from the Gross Payment that the country received.

3 Status of Benefit Distribution

- 3.1 The 1st ER payment, corresponding to the period of May to December 2018, was received in August 2021, the total amount was 6,430,245.00 USD, corresponding to 1,286,049 ERs.
- 3.2 According to the BSP, the MozFIP project covered (40%) has paid part of the operational costs. For that reason, amounted to 300,000.00 USD (60%) of the first payment, were allocated for operational costs, as well as, the 321,512.25 USD will be deducted for performance buffer. 5, 808,732.75 USD of the remaining funds have been allocated as follows:

²⁴ <http://sismdr.fnds.gov.mz/fnds/>

²⁵ <http://sismdr.fnds.gov.mz/fnds/>

Annex 2- Table 1 The funds allocation for all groups of the beneficiaries and the received funds on February 2023

DISTRIBUTION OF MONETARY BENEFITS FROM THE FIRST PAYMENT				
#	Description	Amount (USD)	Committed Value (USD)	Balance of 1 st Payment
1	Amount Received for the 1 st Payment	6,430,245.00		
1.1	Buffer (5%)	321,512.25	-	321,512.25
1.2	Annual Operational Costs	300,000.00	300,000.00	-
NET ER PAYMENT (5.808.732,75)				
1.3.1	Capacities of local Communities (10% of 70%)	406,611.29	406,611.29	-
1.3.2	Community projects (90% of 70%)	3,659,501.63	1,635,352.16	2,024,149.48
1.4	Private Sector ER initiatives (20%)	1,161,746.55	1,161,746.55	-
1.5	9 District Government (4%)	232,349.31	232,349.31	-
1.6	Gilé National Parque (4%)	232,349.31	232,349.31	-
1.7.1	Provincial Government - DPDTAZ (1%)	58,087.33	58,087.33	-
1.7.2	Provincial Government - SPAZ (1%)	58,087.33	58,087.33	-
	Total	6,430,245.00	4,084,583,27	2,345,661.73

3.3 Please find below the table with rubrics that were allocated to different beneficiaries

	ANAC/PNAG	Mocuba	ILE	GURUE	DPTAZ	M. da COSTA	Pebane	Molocue	Mulevala	Mocubela	GILE	SPAZ
Equipment and Machinery	-	-	-	516.86	-	-	3,069.62	-	-	-	4,050.12	3,164.56
In-Country Allowances	15,822.78	10,863.75	4,909.02	4,651.76	34,616.61	6,094.94	15,806.96	9,968.35	6,329.11	12,879.50	16,875.50	29,442.25
Salaries and remuneration	13,449.37	-	-	-	-	-	-	-	-	-	-	-
Civilian Personnel	-	-	-	-	-	-	-	-	-	-	-	-
Assets	18,148.73	6,518.25	-	1,033.73	481.33	2,289.90	5,367.25	2,982.59	2,689.87	2,575.90	2,565.08	8,702.53
Services	79,113.92	3,152.61	-	516.86	17,935.92	4,191.29	3,367.23	967.25	3,231.90	1,287.95	1,687.55	8,702.53
Communication	3,164.56	1,303.65	-	516.86	-	-	1,147.15	995.09	791.14	1,287.95	810.02	2,373.42
Fuels and Lubricants	39,556.96	10,863.75	2,877.98	2,584.31	5,192.96	2,689.87	9,572.78	4,970.71	3,164.56	6,439.75	7,762.73	6,329.11
Transport	63,291.14	10,863.75	-	-	-	-	-	-	2,373.42	1,287.95	-	-
Total	232,547.47	43,565.76	7,787.00	9,820.39	58,226.82	15,266.00	38,331.00	19,884.00	18,580.00	25,759.00	33,751.00	58,714.40

Considering the eligibility criteria, benefits were allocated as follows:

Eligibility criteria	Distribution Modalities	Beneficiaries
PESOPs	Direct allocation	Provincial Government (2)
Annual Plans	Direct allocation	Gilé National Park
PESODs	Performance based allocation/	District Government (9)
Call for proposals	Forest area	Communities (CBOs)
Call for proposals	Matching Grants	Private Sector

DIRECT ALLOCATION

Provincial Government

Due to the restructuring of the provincial government, new organizational structures were created at provincial level, namely the State Representation Council (CRE) and the Executive Council (CE). For this reason, of the 2% foreseen in the BSP, 1% was allocated to the Provincial Environment Service (SPA) at CRE level and 1% to the Provincial Directorate for Territorial Development and Environment (DPDTA) of the CE.

Gilé Nacional Park

According with BSP, the Gilé National Park through ANAC, received the value on the above table. https://drive.google.com/drive/folders/1KHHoAvOdeQSZqttoFgS_fkON8pSjwHQU?usp=share_link

PERFORMANCE BASED ALLOCATION/ FOREST AREA

District Governments

As for 9 district governments, each of them received the certain amount according with performance and forest cover criteria's, please see table 2. The link here:

https://drive.google.com/drive/folders/1jgXSQvm4CrhFz7fxg26INff85mtoYoAl?usp=share_link. The component of REDD+ activities of Economic & Social Plans (PESODs) of the districts, were prepared and discussed at district and provincial level, at different forum's and meetings, always with FNDS team. They can be seen here: https://drive.google.com/drive/folders/1BgFZmTwf7VXT1QYyVaj9HuwIjd0dzcuj?usp=share_link

Call for Proposals

Communities

According with BSP, 70% of the liquid amount is allocated to the Community Based Organizations (CBOs) projects. The 10% of the 70% was deducted on the budget for the communities per each district. It is shown in the following table:

Annex 2- Table 2 Funds allocation for Communities and District Government, according with performance and forest area criteria.

Districts	Net ERs (tCO2e)	ERs Contribution	Forest area	Forest Proportion	Value for Communities	Value for District Government
Alto Molócuè	118 238,87	9,2%	181 398	8,0%	\$313 180.20	\$19 936
Gilé	134 041,77	10,4%	424 689	18,7%	\$531 579.60	\$33 776
Gurúè	282 252,67	21,9%	78 383	3,4%	\$465 176.70	\$29 496
Ile	63 830,12	5,0%	39 448	1,7%	\$122 652.00	\$7 779
Maganja da Costa	112 803,61	8,8%	99 191	4,4%	\$240 442.20	\$15 251
Mocuba	193 431,09	15,0%	508 733	22,3%	\$684 416.63	\$43 428
Mocubela	119 210,79	9,3%	293 496	12,9%	\$405 704.70	\$25 743
Mulevala	141 351,96	11,0%	113 512	5,0%	\$292 629.60	\$18 560
Pebane	120 888,53	9,4%	538 254	23,6%	\$603 720.00	\$38 381
TOTAL	1 286 049	100%	2 277 103	100,0%	\$3 659 501.63	\$232 349

To calculate the amounts to be paid to communities and district governments, the sum of the percentage of performance and the percentage of forest in each district was considered, multiplied by the total amount to be allocated.

Annex 2- Table 3 List of the CBOs Projects approved by value chain

Value chain	Total of screenings
Agriculture	14
Bee keeping	22
Aquaculture	8
Forest	42
Infrastructure	6
Livestock	22
Processing	2
AFS	14
Tourism	1
Production and distribution of schools desks	2
TOTAL	133

CALL FOR PROPOSALS – MATCHING GRANTS

Private Sector

As for the application and decision process on financing, for private sector, the Matching Grants Unit will receive the result of the Letter of Interest proposals evaluation, made by Local Investment Committee. The final table and details will be available for the next BSP report. The pre-qualified list will be public announced for request of the business plan. Then, the complete proposals will be analyzed and verified if they are in accordance with the criteria defined by the product and submitted the decision of the competent body

(Investment Committee). The Investment Committee is an Independent body that will make the final financing decision.

Value Chain	Screening carried out	Failed companies/PIU	Selected Companies	Files under review at PIU	Files sent to WB
Agribusiness	5	2	2	1	1
Agro-Processing	1	1	0	0	0
General trade	1	0	1	1	0
Chicken Farming/Poultry	2	1	1	0	1
Hotels and tourism	8	1	6	4	2
Fish-farming	1	0	1	0	1
Processing	2	1	1	0	1
Tourism	2	1	2	0	2
TOTAL	21	7	14	6	8

- 3.4 The work is being carry out in the support and preparation of the teams that have assisted the process of convening and monitoring the reception of proposals, evaluating and assigning conditions for the projects to be implemented and their follow-up.

Due to the nature and organization of communities in Mozambique, five phases were identified so that benefits can reach to communities:

Five phases needed before a CBO can receive money:

1. Launching and Disclosure of the BSP call
 - a. Consultation of stakeholders
 - b. Mapping of CBOs;
 - c. Definition of terms and conditions
 - d. Formation of support teams (5 central, 8 provincial and 18 district);
 - e. Training of support teams;
 - f. Launching and Disclosure of the call
2. Elaboration and Submission of Proposals
 - a. window clarification
 - b. Mapping and Identification of priorities and value chains;
 - c. Completion of Forms and Investment Plan;
 - d. Submission of proposals
3. Selection and Validation of Proposals
 - a. Creation of the Local Evaluation Committee to evaluate the Proposals;
 - b. Selection of proposals
 - c. Validation of proposals (MGU);
4. Environmental and Social Screening and Procurement
 - a. Visit to Project Implementation Sites;
 - b. Preparation and submission of Environmental and Social Screening forms;
 - c. Acquisition of community projects.
5. Contracting and Disbursement of Funds
 - a. Signature of the Financing Agreement;
 - b. Endorsement of Administrative Court; (is the Contract final approval, but must appears as an annex the Bulletin of the Republic, which legalizes the OCB)
 - c. Disbursement of Funds;
 - d. Project implementation
 - e. Technical assistance and follow-up.

Communities need to be legalized to sign a financial agreement with the FNDS (Phase 5). At this stage, 106 out of the 133 OCBs are already legalized with the publication in the Bulletin of the Republic (BR), 08 paid for publication in the BR and 19 are still to pay for publication.

In the group of 106 OCB's already legalized, all of them have the environmental and social screening form approved, of which 95 were sent to the Administrative Court, where 43 already have the contract approved, and are able to receive the amount, and 52 are still awaiting approval.

Kindly refer to the table below to get the number of community initiatives and the associated amount of money by districts.

Districts	Number of BCO	First Instalment paid	Second instalment to be paid	TOTAL Total amount (MT)	
				Beekeeping value chain	Other Value chain
Alto Molócue	2	1.121.422,50	1.121.422,50	2.242.845,00	
Gilé	10	3.819.964,65	3.819.964,65	7.639.929,00	
Gurué	6	3.931.577,50	3.931.577,50	7.863.155,00	
Ile	5	1.828.240,00	1.828.240,00	3.656.480,00	
Maganja da Costa	1	771.450,00		771.450,00	1.542.900,00
Mocuba	5	1.499.525,50	1.499.525,50	2.999.051,00	
Mocubela	3	869.926,50	869.926,50	1.739.853,00	
Mulevala	2	581.600,00	581.600,00	1.163.200,00	
Pebane	6	2.891.805,00	2.891.805,00	5.783.610,00	
TOTAL	40	17.315.511,65	17.315.511,65	34.631.023,00	

Challenges during process of Benefit Sharing Plan (BSP) implementation and how these challenges have been addressed:

- Annex 2 Table 4 List of the Challenges and how they have been addressed
 In order to reduce time and bring efficiency to each step, the team adopted a different strategy, for example:
 Identification of the Focal Point in the National Press (Central and Provincial Level);
 Identification of Focal Point in the Tax Authority (Two at Provincial Level and one at Central Level);
 Request for Exceptional Visa for BSP OCB Contracts by the Administrative Court;
 Interaction with commercial banks to reactivate bank accounts;
 Technical assistance to OCBs in the processing of files
 Intern program to reinforce the technical team.

In conclusion, the strategy was so effective that the time consumed was reduced. As you can see on the table below:

Activity	2021 to June 2022	2023
----------	-------------------	------

BCO legalization	45 days	7 days
Bank account opening	18 Months	45 days
Visa from Administrative Court	60 days	25 days
Tax Number Identity Synchronization	90 days	20 days
Funds disbursement	20 days	10 days

3.5 Regarding the effectiveness of the mechanisms for ensuring transparency and accountability during the implementation of the BSP, a Feedback and Grievance Redress Mechanism (FGRM) system was created to answer questions, clarify issues, and complaints from individuals or groups affected by the activities under the program. The FGRM also serves as an instrument for the pursuit of harmonious relationship between the parties concerned and interested in the areas of implementation of projects and programs. The information about registration and complaints attendance is available online platform and updated continuously. Individuals and communities who believe they are adversely affected by the community and private sector initiatives are also using this mechanism. A safeguards Information System (SIS) has been designed at the national level and will also report on the progress of the ER Program. The SIS will provide information on how the safeguards will be treated and respected throughout the implementation of the ER Program. When the funds are allocated, all the transferred benefits will be disclosed in the following link: <https://www.fnds.gov.mz/mrv/>

3.6 Despite the disbursement not having reached the CBO's in 2023, main beneficiary, which are the communities, the benefit sharing distributions are still relevant to the core objectives and legitimacy of the RE Program beneficiaries, encourage the adoption of emission reduction measures, between others). However, to change to way of life, take longer period. The most important is provide more and diverse opportunities, and then it will make difference in order to achieve the ER Program, objectives. It is important that the beneficiaries know that the effort must be continuous, and the results depends only of their dedication.

3.7 The intention is to take advantage of the community's governance structure and strengthen the monitoring and follow-up mechanisms, as well as the development of capacities so that the communities can carry out the monitoring and evaluation as the process is being developed, in order to minimize the risks. There will be three monitoring channels:

- The first will periodic meetings with beneficiaries to monitor the status of implementation and satisfaction of beneficiaries;
- The second annual supervision activities will be carried out on the ground to check compliance with the contractual obligations relating to implementation of safeguards activities by service providers, private companies and other direct beneficiaries of the BSP; and
- The third is Feedback and Grievance Redress Mechanism (FGRM), described above.

3.8 To be expected that the beneficiaries continue to understand the relevance of ER program.

4 Implementation of the Environmental and Social Management Measures for the BSP

4.1 The measures that have been applied to manage environmental and social aspects resulting from the activities of implemented projects, such as safeguards instruments, protocols and the safeguards plan, can be seen in Annex 1 above. The community and private sector projects that will be implemented under the BSP will also follow the safeguards instruments, protocols and the safeguards plan.

5 Recommendations for BSP Improvement or Modifications.

5.1 Since benefit distribution has already commenced and completed for some beneficiaries and in progress for the remaining (communities and private sector projects), the status of benefit distribution to date, more updates if any on this aspect will be provided in the next BSP report.

5.2 Learning time for this pilot program was longer than expected, having delayed the start of making funds available for the first year. For example, to make payments to each OCB, there is a need for each organization to be legal and comply with all requirements. Great share of the OCBs do not meet these requirements, making this process complex and time-consuming in the first phase.

5.3 Due to the large number of communities and possible proposals presented, there may be constraints in the analysis, approval and subsequent disbursement. Other options have been discussed, in terms of procedures to make payments more accessible, such as funds to be made available “on CUT” (single treasury account) or “off CUT”. On the other hand, the possibility of contract a service provider that can handle payments, technical assistance and equipment acquisition. Also to maximize the opportunity of EnABLE initiative to support the process of organizing the necessary requirements, that is, legalization and training. So that OCBs can compete and develop programs that diversify their activities in order to reduce the need to open forest areas for shifting cultivation.

5.4 Please refer to the below mention BSP action plan.

Annex 2-Table 5: Benefit Sharing Plan Action Plan.

Activity	Deadline	Status	Observation
1. Inter-Institutional Interaction (WB, MEF, MTA and Provincial Government)			
3° ER Monitoring Report (ER-MR)	30 th of November	Done	
Submission of the BSP and REDD+ Status Report	Continuous	In progress	
EnABLE support	Continuous	In progress	
Regular meetings with MTA, MEF, SdEZ and GPZ	Continuous	In progress	
2. Assistance to the districts			
Support the inclusion of REDD+ activities in annual provincial and Districts Economic and Social Plans	Continuous	In progress	The PESOD for 2021 and 2022 can be seen at the following link https://drive.google.com/drive/folders/1BgFZmTwf7VXT1QYyVaj9HuwIjd0dzcuJ?usp=share_link
Monitoring the implementation of district plans	Continuous	In progress	Monitoring the implementation of PESODs 2021 and 2022
Support the inclusion of REDD+ activities in annual Gilé National Park	Continuous	In progress	Elaborated de Annual GNP for REDD+ 2021 and 2022
3. BSP Communication			
Develop key messages to be harmonized and disseminated to different stakeholders	August/September 2020	Done	Several materials are ready
Ensure the incorporation of messages about BSP in the activities of projects implemented in Landscape	Continuous	In progress	Several materials are ready (Alignment with technicians from PIUs (including extension workers), SDAEs, SDPIs and Service Providers)
Produce and publicize radio plays	December 2021	Done	Produced 7 radio program to publicize in community radio for Local Communities.

Organize a symbolic ceremony for the payment of the first tranche (with massive press coverage)	October 2021	Done	The first ceremony took place at 10.02.2021, For central level in Maputo and the Second at 22.10.2021, in Mocuba at Provincial level. In the both ceremony was attended by journalists and general public
Produce newsletter (Portuguese)	2022	Done and in progress	For donors and other audiences
Produce short videos with English subtitles (about success stories)	November 2021	Not started	For donors and other audiences
4. Financing the CBOs			
Contracting the CBOs	September 2022	In progress	The Terms of reference was developed and approved by WB
Disbursement of Funds	October 2022	In progress	
Project implementation	November 2022	In progress	
Capacity Building	November 2022	In Progress	In 2023
Provision of accounts	November 2022	In progress	In 2023
Monitoring of the activities	Continuous	In progress	
5. Capacity building of communities/CBOs			
Develop ToRs and hire a Community Based Organizations trainer in the Mulevala district in matters related to sustainable management of natural resources and local business development	May 2020 and August/2021	Done	The received funds will be used for implementation of community development projects. This was a pilot program, paid by MozFIP project. The actual schedule will be different. The Service Providers will be training the CBOs under project implementation to all districts
Monitor training activities in collaboration with Community Based Natural Resources Management Network	Continuous by December 2021		Monitored the 2 training activities in collaboration with member of Community Based Natural Resources Management Network Finalized and submitted the final report of capacity the community, with the annex

Contracting of Forestry and Agriculture Service Providers	May 2023	In progress	Ongoing the process of hiring the service provider to train and capacity the CBOs of the landscape. The process to contract the Service Provider is in progress. Proposals for Expression of Interest have already been evaluated. It is expected to proceed to the next step, which will be requesting the technical and financial proposal from the pre-qualified companies. Please see the ToRs for the SP https://www.dropbox.com/s/r3y7vv4m2c9rvci/TdR%20Procurement%20Asistencia%20Tecnica_BSP.docx?dl=0
Contracting of Infrastructure Service Providers	May 2023	In progress	
Contracting of Technical Assistance and Monitoring in the field of beekeeping	May 2023	In progress	This Service Provider will assist 22 community beekeeping projects and guarantee the market for the honey produced.
6. Financing to the private sector/Matching grants			
Review of the Minutes and the CAL report	March	Done	
Screening and licensing	April	Done	
Selection and Announcement of pre-selected candidates	April	Done	
Request for complete proposals from pre-selected candidates	April	Done	
Submission of complete proposals by pre-selected candidates	May	Done	
Analysis and Consultations in relation to the proposals submitted	May	Done	
Final Decision by the Investment Committee	June	In progress	
Negotiation of the Grant Agreement with Approved candidates	June/July	Not started	

Execution of the activities	September	Not started	
Activity monitoring	September	Not started	
7. Hiring of the BSP reinforcement team			
Develop ToRs for technical safeguards assistant	October 2021	Done	
To develop ToRs and hire technical communication assistant	April 2023	Done	The contract is already signed
To Develop ToRs and hire technical business unit assistant	April 2023	Done	The contract is already signed
To develop ToRs and hire technical safeguards specialist	June 2023	Done	The contract is already signed
8. 2 nd Call for CBO's projects launched			
Technical sheet for call of community projects	September 2023	Done	
Disclosure of the Community Projects Financing Window	January 2024		
Incorporation of the Eco Community and MozDGM projects	September 2023	Done	
Produce registration / registration form for applications	October2023	Done	
Disseminate the finance window	January 2024		
Call launch	January 2024		
Support the logistics of the selection process for potential beneficiaries	February2024		
Support in the process of preparing investment plans	January 2024		
Perform screening and licensing of selected subprojects	February/March2024		

Development of communication strategy and dissemination materials (pamphlets, etc.)	December/January 2023/24		
9. Financing the CBOs			
Contracting the CBOs	April 2024		
Disbursement of Funds	May 2024		
Project implementation	May/June 2024		
Capacity Building	May/June 2024		
Provision of accounts	May 2024		
Monitoring of the activities	Continuous		
10. Monitoring under the Emissions Reductions Program			
Submission of approved proposals to SURVEY123	April 2023	Done	
Dashboard production (Grants, Approved, Value chain)	April 2023	Done	
Preparation of Monitoring Sheets (Finance, Procurement and Execution)	April 2023	Done	
Alignment meeting to approve the forms	April 2023	Done	
Preparation of forms in SURVEY123	April 2023	Done	
Design of the Monitoring Dashboard	April 2023	Done	
Data collection and input	Continuous	In progress	
System Implementation	Continuous	In progress	

ANNEX 3: INFORMATION ON THE GENERATION AND/OR ENHANCEMENT OF PRIORITY NON-CARBON BENEFITS

Priority Non-Carbon benefits

1. In the area on the ER program is taking place, several activities from the World Bank-financed investment projects, such as the Agriculture and Natural Resources Landscape (Sustenta) Project; Mozambique Sustainable Rural Economy Project (MozRural) Project; Mozambique Forest Investment (MozFIP) Project and the Dedicated Grant Mechanism (MozDGM) project, that provide non-carbon benefits are taking place, mainly within the scope of the implementation of the MozFIP, SUSTENTA and MozRural projects. The list can be seen in the table below.

Annex 3 -Table 1: List of the identified set of priority Non-Carbon benefits

Priority Non-Carbon Benefit	Details on activities for generation and enhancement Approach (as defined in ERPD including relevant indicators)
Improvement of local livelihood through securing long-term access to forest resources and environmental benefits	
Increase of land areas under sustainable landscape management practices	<p>Areas of multiple-use forests planted established under the Planted Forest Grant Scheme;</p> <p>Restoration of degraded areas and protection of fragile habitats, included waters spring;</p> <p>Areas of agroforestry systems established;</p> <p>Gilé National Park under improved management plans and law enforcement, as measured by the Management Effectiveness Tracking Tool.</p>
Long term adoption of sustainable land use practices	<p>Forest Plantations under the Planted Forest Grants Scheme;</p> <p>Areas of agroforestry systems established;</p> <p>Sustainable production of charcoal;</p> <p>Value chain development of non-timber forest products (NTFP)</p>
Clarified land tenure	Community land delimitations and registered parcels for Right to Use and Benefit from Land (DUAT) emissions, and DUAT emissions for families
Strengthening of forest governance and forest resources management	
Improved forest governance and transparency	<p>The approval of the National Land Use Plan, which is expected to foster proper land use planning, with land use decisions being based on transparent information and a consultative process on land use priorities;</p> <p>Establishment of national Monitoring, Reporting and Verification (MRV) system;</p> <p>Patrol and inspection, prevention and detection of the forestry sector;</p> <p>The operationalization of the National Forest Information System, which is expected to improve information availability, accessibility and transparency, contributing to an effective forest monitoring and control;</p> <p>Biannual evaluation of forest concessions to check the compliance of forest operators with management plans and other legal and basic sustainability requirements (fiscal obligations, social security, qualified rangers, concession contract, availability of statistical information, industrial plans, technical capacity, delimitation of area and harvesting blocks, etc.).</p>

Enhanced participatory forest and land use management	Consist of improved overall governance and access to information. Through the well-functioning of the Zambézia Integrated Development Platform (ZIDP), it includes enhanced landscape-level dialogue and multi-stakeholder decision-making on the use of natural resources, contributing to integrated landscape management. Eventually, their ability to participate in decisions over natural resources can empower stakeholders and bring additional long-term benefits for resource management.
---	---

There is information available on the achievement of goals for the generation and / or appreciation of Non-Carbon Benefits in the progress reports of projects that took place at ZILMP, mainly from MozFIP²⁶.

Annex 3-Table 2: The achievement of goals of the generation of Non-Carbon Benefits

Priority Non-Carbon Benefit	Details on activities for generation and enhancement Achievement of goals																																																																																						
Improvement of local livelihood through securing long-term access to forest resources and environmental benefits																																																																																							
Increase of land areas under sustainable landscape management practices	<p>Most of the activities under this Non-Carbon Benefits priority are part of component 1 of the MozFIP project. The fulfillment of the activities of this component can be seen in its 2021 e 2022 annual report, approved by the world bank. For more details, see the Link</p> <p>https://drive.google.com/file/d/14VL3QaGDGWcHO6gGov0AaxHMUeJWp5E/view?usp=share_link</p> <table border="1"> <thead> <tr> <th rowspan="2">District</th> <th rowspan="2">Planted area (ha)</th> <th colspan="2">Restoration Program (ha)</th> <th rowspan="2">SAF (ha)</th> <th rowspan="2">Total (ha)</th> </tr> <tr> <th>Enrichment planting* (ha)</th> <th>Management of natural regeneration* (ha)</th> </tr> </thead> <tbody> <tr> <td>Alto Molócué</td> <td>86.49</td> <td>42</td> <td>63</td> <td></td> <td>191.49</td> </tr> <tr> <td>Gilé</td> <td></td> <td></td> <td></td> <td></td> <td></td> </tr> <tr> <td>Gurué</td> <td>148.32</td> <td>61</td> <td>6</td> <td>50</td> <td>265.32</td> </tr> <tr> <td>Ile</td> <td>44.85</td> <td>7</td> <td></td> <td></td> <td>51.85</td> </tr> <tr> <td>Lugela</td> <td></td> <td>9</td> <td>8</td> <td></td> <td>17</td> </tr> <tr> <td>Maganja da Costa</td> <td>43</td> <td>18</td> <td>61</td> <td></td> <td>122</td> </tr> <tr> <td>Mocuba</td> <td>72.79</td> <td>36</td> <td>15</td> <td>10</td> <td>133.79</td> </tr> <tr> <td>Mocubela</td> <td></td> <td></td> <td></td> <td></td> <td></td> </tr> <tr> <td>Milange</td> <td>37.12</td> <td>29</td> <td>51</td> <td></td> <td>117.12</td> </tr> <tr> <td>Mulevala</td> <td></td> <td></td> <td></td> <td>10</td> <td>10</td> </tr> <tr> <td>Pebane</td> <td></td> <td></td> <td></td> <td></td> <td></td> </tr> <tr> <td>Total</td> <td>433</td> <td>202</td> <td>204</td> <td>70</td> <td>909</td> </tr> <tr> <td>* 2021_2022</td> <td></td> <td></td> <td></td> <td></td> <td></td> </tr> </tbody> </table> <p>The area under sustainable landscape management practices achieved at the end of project in Zambézia is 39,949.00ha</p>	District	Planted area (ha)	Restoration Program (ha)		SAF (ha)	Total (ha)	Enrichment planting* (ha)	Management of natural regeneration* (ha)	Alto Molócué	86.49	42	63		191.49	Gilé						Gurué	148.32	61	6	50	265.32	Ile	44.85	7			51.85	Lugela		9	8		17	Maganja da Costa	43	18	61		122	Mocuba	72.79	36	15	10	133.79	Mocubela						Milange	37.12	29	51		117.12	Mulevala				10	10	Pebane						Total	433	202	204	70	909	* 2021_2022					
District	Planted area (ha)			Restoration Program (ha)				SAF (ha)	Total (ha)																																																																														
		Enrichment planting* (ha)	Management of natural regeneration* (ha)																																																																																				
Alto Molócué	86.49	42	63		191.49																																																																																		
Gilé																																																																																							
Gurué	148.32	61	6	50	265.32																																																																																		
Ile	44.85	7			51.85																																																																																		
Lugela		9	8		17																																																																																		
Maganja da Costa	43	18	61		122																																																																																		
Mocuba	72.79	36	15	10	133.79																																																																																		
Mocubela																																																																																							
Milange	37.12	29	51		117.12																																																																																		
Mulevala				10	10																																																																																		
Pebane																																																																																							
Total	433	202	204	70	909																																																																																		
* 2021_2022																																																																																							
	Most of the activities under this Non-Carbon Benefits priority are part of component 2 of the MozFIP project. The fulfillment of the activities of this																																																																																						

26 <https://www.fnds.gov.mz/index.php/en/our-projects/project-list/MozFIP>
<https://www.fnds.gov.mz/index.php/en/our-projects/project-list/MozFIP>

Long term adoption of sustainable land use practices	component can be seen in 2021 annual report, approved by the world bank. For more details, see the Link: https://docs.google.com/document/d/1I0fkwGinfrzVAYVvHIh7gXKDQJmKiQvs/edit?usp=drive_link&oid=115860222606486742127&rtpof=true&sd=true																												
Clarified land tenure	This is also part of MozFIP component 1. 187 communities were delimited and land titles number emitted increased to 17.189. <table border="1"> <thead> <tr> <th></th> <th>2021</th> <th>2022</th> <th>Total</th> </tr> </thead> <tbody> <tr> <td>Communities delimited</td> <td>18783</td> <td>187</td> <td>270</td> </tr> <tr> <td>PCUT</td> <td>14468</td> <td>144107</td> <td>175</td> </tr> <tr> <td>Unofficial certificates</td> <td>16247</td> <td>162106</td> <td>153</td> </tr> <tr> <td>PDutRDuat</td> <td>15318 344</td> <td>15326 881</td> <td>45 225</td> </tr> <tr> <td>DUAT</td> <td>16 529</td> <td>19 880</td> <td>36 409</td> </tr> <tr> <td>Titles</td> <td>17189</td> <td>1718917 189</td> <td>17 189</td> </tr> </tbody> </table>		2021	2022	Total	Communities delimited	18783	187	270	PCUT	14468	144107	175	Unofficial certificates	16247	162106	153	PDutRDuat	15318 344	15326 881	45 225	DUAT	16 529	19 880	36 409	Titles	17189	1718917 189	17 189
	2021	2022	Total																										
Communities delimited	18783	187	270																										
PCUT	14468	144107	175																										
Unofficial certificates	16247	162106	153																										
PDutRDuat	15318 344	15326 881	45 225																										
DUAT	16 529	19 880	36 409																										
Titles	17189	1718917 189	17 189																										
Strengthening of forest governance and forest resources management																													
Improved forest governance and transparency	The activities also are part of component 2. Please see above the progress in achieving Non-Carbon Benefits																												
Enhanced participatory forest and land use management	This activity reached 33% and 32% in relation to what was planned for 2019 and 2020. For more details, see the Link: (https://docs.google.com/document/d/1I0fkwGinfrzVAYVvHIh7gXKDQJmKiQvs/edit?usp=drive_link&oid=115860222606486742127&rtpof=true&sd=true)																												

Adapted from table 2, annex 1 of 2021 and 2022 annual MozFIP Report.

https://www.dropbox.com/s/qibzags2m4xjmjz/Relat%C3%B3rio%20Balan%C3%A7o%20Anual%20de%202019_APROVADO.docx?dl=0 and https://www.dropbox.com/s/n49lq01rpouw836/Relat%C3%B3rio%20Anual%202020_FINAL.docx?dl=0

Other Non-Carbon benefits and additional information as linked to Monitoring and Evaluation Framework

Other (non-priority identified) Non-Carbon benefits

The activities that provided non-carbon benefits are activities that the beneficiaries were already developing, but in a more organized and technically oriented way. In this way, all of them are in line with their culture. These activities, although not presented by gender, the beneficiaries are the families where the whole household will benefit. Such as:

Livelihood enhancement and sustainability

Within the ER Landscape of Zambézia program, with to sustain and improve livelihoods is one of the main objectives. With the priority non-carbon benefits listed above, the following non-carbon benefits have been provided in the ER MR period 2021 and 2022. As a result of the four World Bank investment projects, such as the Agriculture and Natural Resources Landscape (Sustenta) Project; (SREP) project; Mozambique Forest Investment (MozFIP) Project and the Dedicated Grant Mechanism (MozDGM) project that were implemented in the ER Program2020. Due to the areas reached and the number of people involved, these medium and long-term benefits will make a difference in local communities' livelihood.

1.1. The ER program achieved²⁷:

- 2.1.1 453,35 ha with agroforestry systems established, including 1610 farmers from 42 communities were involved; and 1332.98 ha of restored area.
- 2.1.2 268 Charcoal producers, organized in 4 associations in Pebane, Mocubela, Maganja da Costa and Ile;
- 2.1.3 Inputs distributed and technical assistance provided to about 1,300 community members in the buffer zone of the Gilé National Park – tree owners, beekeepers, charcoal makers and small farmers;
- 2.1.4 187 communities were delimited in 2021 and 2022, in an area of 426.477 ha in 2021 to 2022 and 19,860 individual land uses rights issued respectively.
- 2.1.5 Small nurseries produced in 2020/2021 a total of about 120,000 seedlings of different species. Out of the total, Chazeira de Moçambique in Gurue produced 80,000 of exotics spp and 40,000 of native spp were produced by 13 different small nurseries including, women and youth, mainly in Gurue, A.Molocue and Mocuba.
- 2.1.6 In 2021/2022 season, a total of 148,000 seedlings of native and exotic species, Chanfuta, Umbaua, Eucalyptus, Acácia, were produced by 9 (nine) small nurseries of which 2 (two) involved 16 women.
- 2.1.7 In partnership with the SPA, environmental screenings and monitoring of good practices were performed as well as identification of sensitive habitats, and Environmental Licenses were issued

Biodiversity

2.2 In the ER program, two projects are testing ways to conserve biodiversity in the Gilé National Park: i) the Mozambique Conservation Areas for Biodiversity and Development Project (MozBio); and ii) the Forest Investment Project (MozFIP). The MozBio project contributed with financial and technical support the Gilé National Park and its buffer zone; while MozFIP has provided financial and technical support to the law enforcement activities in the Gilé National Park. The Gilé NP is a key biodiversity area in the landscape. In addition, the MozFIP and Sustenta projects are restoring degraded areas, which also contribute to the biodiversity conservation through the connection of forest fragments.

Protected/conserved areas

2.3 The ER Program area included the Gilé National Park (GNP),, that covers an area of 2861 km² and 1671 km²a buffer zone. with 1671 km². The area has not changed.

Re/afforestation and restoration

2.4 Total reforested and restored area: Beneficiaries contracted, restored, households

Until 2022, the final year of the project, 3,000 Ha were planted, benefiting a total of 223 plantation owners or Beneficiaries, of which 36% were Women, and 670ha of degraded areas restored. 2014 Permanent and Seasonal jobs created. Women were involved in almost all forestry activities namely field preparation, planting, plantations maintenance, nurseries through associations and as employees. This positive involvement was achieved thanks to the great contribution of the women empowering program (GALS)

²⁷ Sources of information: 1- Etc Terra-Rongead / IGF, Sétimo relatório de progresso Mozbio-Gilé, Dezembro de 2018 (<https://www.fnds.gov.mz/index.php/pt/documentos/relatorios/mozbio-gile-se-timo-relato-rio-de-progresso-final>); 2- MozFIP 2018 annual report (<https://www.fnds.gov.mz/index.php/pt/documentos/relatorios/mozbio-gile-se-timo-relato-rio-de-progresso-final>).

At the same time, 1,035 ha of Agroforestry Systems were established in the Districts of Gurue, Mocubela and Mulevala, benefiting a total of 1,672 local producers, of which 34% are women, see the table 3 below.

Annex 3-Table 3: Planted area 2021/2022

		2021	2022	Total
Beneficiaries contracted (n)		2013	104	2117
% of Women		24%	36%	
Planted area (ha)		1.332,98	432,57	1765,55
Restored (ha)		359	406	765
SAF	Households	1.610	1452	3062
	% of Women	(36%)	36%	
	Ha	1.035	942	1.977

Finance and Private Sector partnerships

2.5 The program budget has not changed.

We do not yet have evidence with private sector partnerships. The partnerships are at beginning stage

The program budget has not changed.

3 Other Non-Carbon benefits and additional information

Policy development

3.1 Under CF Program, through MozFIP project, the National Forest Policy, was developed and approved in early 2020. The key changes in this legislation is the environmental domain main: to ensure the protection, conservation, creation, enhancement, restoration and sustainable use of the forest heritage. In this domain, priority is given to the maintenance of forest ecosystems, threatened or endangered species, preservation and enhancement of their services through **payment for environmental services**, mitigation of climate change, maintenance of water services, soil stabilization and others. More details will be providing next BSP report

Capacity building

Small nurseries produced in 2020/2021 a total of about 120,000 seedlings of different species. Out of the total, Chazeira de Moçambique in Gurue produced 80,000 of exotics spp and 40,000 of native spp were produced by 13 different small nurseries including, women and youth, mainly in Gurue, A.Molocue and Mocuba.

In 2021/2022 season, a total of 148,000 seedlings of native and exotic species, Chanfuta, Umbaua, Eucalyptus, Acácia, were produced by 9 (nine) small nurseries of which 2 (two) involved 16 women.

In partnership with the SPA, environmental screenings and monitoring of good practices were performed as well as identification of sensitive habitats, and Environmental Licenses were issued.

GALS was a tool to reach and train beneficiaries:

- a) In Gurue, 85 participants in GALS, including women, were trained with the objective to support beneficiaries and their households to remove gender barriers that prevent them plan and use resources for family socio-economic progression
- b) Training was performed on finances (credito rotativo), health and empowerment of women
- c) Beneficiaries were trained on environmental good practices including use of agro-chemicals and grievances mechanism

In the ER Program, three main beneficiaries were identified under the four WB projects (MozFIP, MozBio, SUSTENTA and MozDGM), namely, local communities, private sector and Zambézia provincial and 9 districts government. In 2021 and 2022, the beneficiaries involved were mainly, the local communities and Zambézia government:

During the report period, several training and capacity interventions involved local communities.

In the Gilé National Park and its buffer zone²⁸, in the subjects of charcoal production, conservation agriculture and non-wood forest products, some training actions were made. Such as recycling and training actions for the construction of improved charcoal works. To obtain greater efficiency in charcoal production and to reduce deforestation, 75 charcoal workers were involved, of which of them 12 were women; including 2 theater groups to present plays about the impact of uncontrolled fires and methods to control them, with 884 participants, of 4 areas, namely Namala, Naheche, Moneia and Mulela²⁹. On the same theme, 920 people were sensitized, of which 300 were women. On Conservation Agriculture³⁰, 327 people were trained on improved dryers, of which 64 were women; 2 producers from the community nurseries supported by the INCAJU nursery team were trained in carrying out the grafting of cashew trees and mango trees. 1061 beneficiaries during the 2018/19 agricultural season (78% men and 22% women) were trained in the practice of conservation agriculture and followed by project technicians in their own agricultural field, with dissemination to 40 volunteer producers and receiving support technician

Training was carried out on: (i) Planting Techniques, including hydrogel (ii) protection against pest and disease attacks; (tree species and preparation of organic pesticides), application of inorganic pesticides (iii) training in georeferenced data collection tools, survey123 and Kobocollect (iv) Yield measurement; (v) post-harvest and commercialization. These training sessions were delivered by FNDS specialists, PIU team and PS specialists, and covered a total of 124 individuals being (15) extension workers, (43) community facilitators. and 21 heads of village and production. The reply was made to other extensionists, community facilitators including local producers.

In the question period, the process of issuing licenses for the start of activities was late, conditional on the regularization of the infractions of the previous year for access to licensing in 2021 all operators who were in an irregular situation, as a result of the articulation with the licensing institution. In order to safeguard the performance and improve the technical skills of inspectors during the forest inspection process, training was carried out for inspectors at all fixed posts and mobile brigades regarding the procedures during the action. This initiative appears as a way to standardize individual perceptions and clear up eminent doubts during the process of issuing notices of fines, as well as in the case of news reports. To this end, the delegation carried out staggered nine (9) rounds of all posts where they were taught based on the Forestry and Wildlife Legislation Manual, covering about 37 inspectors.

²⁸ Source: Etc Terra-Rongead/IGF, Relatório final do projecto de progresso Mozbio-Gilé, 2016-2019, Outubro de 2019 (click [here](#) for more details).

²⁹ Etc Terra-Rongead / IGF, Nono relatório de progresso Mozbio-Gilé, Julho de 2019 (click [here](#) for more details).

³⁰ Etc Terra-Rongead / IGF, Oitavo relatório de progresso Mozbio-Gilé, Abril de 2019 (click [here](#) for more details)

Consultation seminars were held on the Forest Surveillance Strategy and the recommendations obtained during the sessions were incorporated. The strategy is now complete and the document is modeled. Several impressions were made and distributed among the various key stakeholders. The implementation of joint surveillance in Gilé started with a delay, with nine (9) joint surveillance missions being carried out within the conservation area and buffer zone, including the multiple use areas of the landscape around the Park, with the primary objective of controlling the buffer zone and contiguous or multiple-use areas around the PNGile within the landscape area in the districts of Mulevala, Pebane, Mocubela and Gilé.

As a way of minimizing the problems encountered in the scope of the actions of the actors involved in forest inspection, including the instruction of the process, training was carried out with the participation of 50 people, including prosecutors and judges from 8 districts, PPRNMA, Customs, DPAQUA and PNG from all districts of Zambézia Province, which culminated in a Declaration/Commitment in an effort to strengthen the alignment and articulation of actors and improve intra and inter-institutional collaboration. Additionally, the procedures for the destination of goods and mechanisms of coercive collection were clarified.

During 2021, training was carried out for key actors on procedures for issuing notices of fines and on matters of Legal procedures in forest inspection, with a debate on the topic being planned during the II National Meeting of Environmental Inspection. The workshop with UNODC was also held, with the preparation of the corruption risk assessment plan and risk mitigation plan by the 1st quarter of 2022.

A regional training center on Wood identification was carried out between January and June 2021. It covered an average of 30 to 35 participants from the 4 provinces, namely: Zambézia, Sofala, Manica and Tete. This training involved technicians from the National Forestry Directorate (DINAF), Provincial Environmental Services (SPA), Provincial Delegations of the Agency for Environmental Quality (AQUA), Customs / Tax Authority (AT) and Police of the Republic of Mozambique (PRM).

The service provider (CESC) was hired to train CBOs in the district of Mulevala, having identified 29 communities and selected 20 CBOs, which benefited from this pilot training. The following activities were carried out: (i) launch of the Capacity Building Project for the Zambézia Landscape CBOs, in the District of Mulevala, (ii) identification of 40 community agents in the 20 CBOs; (iii) training of 37 community agents from the 20 beneficiary OCBs; (iv) Sensitization and Capacity Building of the 20 beneficiary CBOs in terms of community governance, a package led by the Community Governance Network of Natural Resources (R-GCRN); (v) Identification of partnerships between CBOs and the private sector; (vi) sensitized 390 members of the 20 beneficiary CBOs about the rights and duties of communities in relation to land and other natural resources; (vii) 588 members of the CBOs were trained in terms of community governance, a package led by the representative of the Community Governance Network of Natural Resources (R-GCRN).

Realized on October 15, 2021, at the Radisson Blu Hotel, in Maputo, a ceremony for the delivery to Mozambique of the First Payment for Emission Reduction. The ceremony was led by Her Excellency the Minister of Land and Environment, Ivete Maibaze, and was attended by the Chairman of the FNDS Board of Directors, Cláudio Borges, the Global Director of Climate Change, Bernice K. Van Bronkhorst, and the representative of World Bank Director in Mozambique, Michael Matera.

In turn, on October 21, 2021, at the SUMEIA Hotel, in Mocuba, a replica of the ceremony for the disclosure of the first payment for the reduction of emissions from Deforestation and Forest Degradation, in Zambézia, which culminated in the delivery of the Certificate of Merit to the districts covered by the Program, in recognition of their efforts in Reducing Emissions registered in 2018. The ceremony was led by Her Excellency the Secretary of State of the Province of Zambézia and was attended by the Inspector General of the Ministry of Land and Environment (MTA), District Administrators of the 9 Districts, President of the Municipal Council of Mocuba,

Director of the Provincial Environment Service and of the Provincial Directorate of Territorial Development of Zambezia, sector technicians, Representatives of the Private Sector, Civil Society, Academies, Local Communities, among others.

Launched on February 14, 2021 the Call for Manifestation of Interest (Mol) Eco Community in Mocuba (Zambézia), with district replicas with the participation of government institutions. The provincial launch event was well attended with the presence of CEPs, DPTA, SPAEs, OCBs, companies and media that were invited to cover the event. This window targets OCBs and OCBs in partnership with companies. Mol aims to contribute to reducing deforestation, increasing community awareness of conserving forest resources and promoting non-wood forest products.

During the application submission phase, 139 CBOs that applied were registered. Also for Manifestation of Interest (Mol) CBOs in partnership with companies applied for a total of 9 companies. After the evaluations were carried out by the local Evaluation Committee, the following results were obtained: 26 approved CBOs and 04 companies approved for Zambézia. The official disclosure of the results of the Call for companies was made, unfortunately due to the negative effect of the covid-19 pandemic and the restrictions imposed by the Government, the event was virtual. Disclosure to CBOs followed direct communication with approved CBOs and through official letters sent to SDAEs and district headquarters. Environmental and social screening was also carried out with the involvement of SPA technicians. The team made visits to the places where the proponents intend to implement their initiatives and in the case of small areas there is no need to license the activities as the impacts are negligible.

Launched in November 23, 2021 the Call for Manifestation of Interest (Mol) for Benefit Sharing Plan (BSP) in Quelimane (Zambézia), the first for access to funding for Community-Based Organizations (OCB's) and for access to subsidized subsidies for companies in Zambézia. The end of the two calls was scheduled for January and February 2022 respectively. The provincial launch event was well attended with the presence of CEPs, DPTA, SPAEs, OCBs, companies and media that were invited to cover the event. This window targets OCBs to contribute to reducing deforestation, increasing community awareness of conserving forest resources and promoting non-wood forest products.

The multidisciplinary team of the BSP of the FNDS at the level of Zambézia and Maputo, dedicated much of its time in dissemination of the Call within the scope of the BSP in the Zambézia landscape, with district replicas with the participation of government institutions. 2019³¹ and 2020

Strengthen capacity building for sustainable forest management in two community forest concessions in the districts of Gilé and Mulevala. 135 members were trained, of which 45% were women.³²

During the ER MR period, the following training actions were carried out for members of the provincial government of Zambézia, 9 districts and Platform of Zambézia:

The first ordinary meeting of the Zambézia platform coordination group took place on January 22, 2022, in the INSS meeting room, with the participation of 14 members of this body, 4 women and 10 Men, due to the guidelines decreed in the state of emergency, and on the other hand, due to the overlapping agendas of some members, this aimed to discuss the following agenda items: (i) Presentation and appreciation of the activities developed by each thematic group of the platform; (ii) Consideration and Approval of the Platform's Activities Plan for the Year 2021; (iii) Setting the date and agenda for the 4th Plenary Session; and (iv) Consideration of the TOR's for hiring the Executive Secretary of the platform.

And on July 1, 2022, in the RADEZA meeting room, Quelimane city, the second ordinary meeting of the same Coordination Group of the Zambézia platform took place, with the participation of 18 members of this body, 7

³¹ NEA-WWF, Relatório de progresso narrativo e financeiro do Mecanismo de Doação Dedicado às Comunidades Locais no ano de 2019, Março de 2020 and NEA-WWF, Relatório de progresso narrativo e financeiro do Mecanismo de Doação Dedicado às Comunidades Locais no ano de 2020, Março de 2021

³² Source: MozFIP 2019 annual progress report, MozFIP 2020 annual progress report

of which were women. and 11 Men, having discussed the following agenda items: (1) Presentation and discussion of the current stage of the platform's functioning; (ii) Perspectives on the future of the Zambézia Platform; (iii) Approval of the Semiannual Activity Plan (July to December 2021); and (iv) Scheduling the Plenary Session of the Platform, date, place, modality, format and topics.

During 2021, “on the job trainings” were held in the 9 MozFIP districts in Zambézia, as well as the seminar on Geospatial planning tools in December, for district service technicians.

In the introduction training, service providers and all technical staff who have direct contact with the local communities, special attention was given to “community relations” including aspects to be respected in the communication with this target group, including best places to hold meetings, norms and customary habits.

To date, all activities of the NCB have been carried out in compliance with the safeguards requirements and within the scope of the BSP should also continue to comply with the safeguards instruments, protocols and plans.

Other

3.3 Institutional Strengthening

With the implementation of the Safeguards Information System and the Dialogue and Complaint Mechanism, it allows for greater transparency in the process of implementing the emission reduction program and projects.

Annex 4: CARBON ACCOUNTING - ADDENDUM TO THE ERPD

"Intentionally left blank"

ANNEX 5: DETAILED DESCRIPTION OF THE APPLICATION OF THE REVERSAL RISK ASSESSMENT TOOL

Detailed Description of the Assessment of Reversal Risk

The assessment of the reversal risk was conducted based on the risk categories defined by the FCPF applying risk assessment tool. For each risk factor, available evidence was compiled from program documentation, consultations, institutional arrangements, and field implementation.

1. Lack of Broad and Sustained Stakeholder Support

- Stakeholder consultation has been systematically conducted and documented, with clear reporting in safeguards reports.
- A Benefit Sharing Mechanism has been established and validated with communities.
- Memoranda of Understanding (MoUs) were signed with implementing partners to formalize collaboration.
- A Feedback and Grievance Redress Mechanism is operational, ensuring that concerns from stakeholders are addressed.
- These mechanisms ensure that local communities and partners remain engaged over the long term.

Conclusion: Risk is considered **low** due to strong participation and long-term stakeholder commitment.

2. Lack of Institutional Capacities and/or Ineffective Vertical/Cross-Sectorial Coordination

- A dedicated national structure is designated and empowered to implement the ER Program.
- Government entities have prior experience in implementing multi-sectoral initiatives.
- Collaboration between central and provincial levels has been formalized and is functional.
- Inter-sectorial coordination mechanisms are in place, supported by complementary projects.
- Staff have been deployed at the local level, with continuous capacity-building and training.

Conclusion: Risk is considered **medium**; as institutional capacity is in place but continued strengthening is needed.

3. Lack of Long-Term Effectiveness in Addressing Underlying Drivers

- The ER Program is complemented by other initiatives targeting deforestation and degradation drivers.
- Legal and regulatory frameworks already integrate REDD+ objectives.
- Incentives for sustainable agriculture and land use practices are being implemented, ensuring continuity beyond the ERPA.
- Stakeholders perceive tangible non-carbon benefits, such as improved land tenure and alternative livelihoods.
- Extension services and local institutions are working to ensure adoption of sustainable practices in the long term.
- **Conclusion:** Risk is considered **low**, given the legal frameworks and ongoing complementary initiatives.

4. Exposure and Vulnerability to Natural Disturbances

- Mozambique is exposed to risks such as fires, droughts, and pests.
- Preventive measures have been promoted, including climate-smart agriculture and fire management activities.

- Although mitigation measures exist, the country remains vulnerable to climate-related events beyond the Program's control.
- **Conclusion:** Risk is considered **medium**; as natural events cannot be fully prevented despite mitigation actions.

ANNEX 6: ER MONITORING REPORT (ER-MR) ON THE AREA OUTSIDE THE SCOPE OF ZAMBÉZIA INTEGRATED LANDSCAPE MANAGEMENT PROGRAM (ZILMP)

This annex was prepared as part of the Government's commitment to monitor and report in parallel the annual emissions reduction in the area outside the scope of Zambézia Integrated Landscapes Management Program (ZILMP) within the Zambézia province under the Emission Reductions Purchase Agreement (ERPA) signature.

5.1 CARBON POOLS, SOURCES AND SINKS

5.1.1 Description of Sources and Sinks selected

Sources/Sinks	Included?
Emissions from deforestation	Yes
Emissions from forest degradation	No
Enhancement of carbon stocks	No
Sustainable management of forests	No
Conservation of carbon stocks	No

5.1.2 Description of carbon pools and greenhouse gases selected

Carbon Pools	Selected?
Above Ground Biomass (AGB)	Yes
Below Ground Biomass (BGB)	Yes
Biomass in non-woody vegetation	No
Dead organic matter	No
Soil Organic Carbon (SOC)	No

GHG	Selected?
CO ₂	Yes
CH ₄	No
N ₂ O	No

5.2 REFERENCE LEVEL

5.2.1 Reference Period

The reference period is from 2005 – 2015 (11 years).

5.2.2 Forest definition used in the construction of the Reference Level

According to the national REDD+ strategy and to the Final Report on Forest Definition (Falcão and Noa, 2016) approved by MITADER in November 2016, forest in Mozambique is defined as followed: **minimum area of 1 ha, minimum height at maturity of 3 m and minimum tree cover of 30%.**

The previous GHG inventories used the previous forest definition of Mozambique (minimum area of 0.5 ha, minimum height of 5m and minimum tree cover of 10%). However, future GHG inventories will use the updated forest definition.

5.2.3 Average annual historical emissions over the Reference Period

1.1.3.1.1.1 Description of method used for calculating the average annual historical emissions over the Reference Period

The UNFCCC does not give any directives with regards to the reference period for the RL. However, the Forest Carbon Partnership Facility (FCPF) have specific guidelines, setting a minimum of 10 years and a maximum of 15 years. The chosen period for the construction of the RL is from 2005 to 2015, 11 years.

In accordance with the UNFCCC decisions, the method used to assess emissions is the one described in IPCC (2006) for Land (Forest land in the present case) converted to other land use (e.g., croplands, grasslands, etc.) consisting on the multiplication of activity data – area of land converted from forest land to other land (e.g., cropland or grassland in the present case) – by emission factors – difference of carbon stocks before and after deforestation – as presented on the following equations. The data used for the present document are Tier 2 (country specific data or country level estimates) or Tier 3 (data specifically produced for the ER Program) when possible. Activity data are produced on the reference period with spatially explicit method based on available satellites images. Emissions factors are derived from literature or forest inventory in the accounting area.

In compliance with criterion 13 of FCPF MF (FCPF, 2016) that specifies that RL should not exceed the average annual historical emissions, different activity data of the reference period will be averaged to produce annual deforestation areas over the whole period.

As analysis is done over the reference period, long term (11 years) changes (increase or decrease) of carbon stocks on deforested areas (land converted to another land use) are considered instead of annual increase or decrease - see the **Equation 16**.

Gross emissions of the RL from deforestation over the Reference Period (RL_{RP}) are estimated as the sum of annual change in total biomass carbon stocks (ΔC_{B_t}) during the reference period as shown in the equation below.

$$RL_{RP} = \frac{\sum_t^{RP} \Delta C_{B_t}}{RP} \quad \text{Equation 15}$$

Where:

$$\begin{aligned} \Delta C_{B_t} &= \text{Annual change in total biomass carbon stocks at year } t; \text{ } tC \cdot \text{year}^{-1}; \\ RP &= \text{Reference period, years.} \end{aligned}$$

Following the 2006 IPCC Guidelines, the annual change in total biomass carbon stocks forest land converted to other land-use category (ΔC_{B_t}) would be estimated through the following equation:

$$\Delta C_{B_t} = \Delta C_G + \Delta C_{CONVERSION} - \Delta C_L \quad \text{Equation 16}$$

Where:

$$\begin{aligned} \Delta C_{B_t} & \text{ Annual change of total biomass carbon stocks during the period, in } tC \text{ per year;} \\ \Delta C_G & \text{ Annual increase in carbon stocks in biomass due to growth on land converted to another land-use category, in } tC \text{ per hectare and year;} \end{aligned}$$

$\Delta C_{CONVERSION}$ Initial change in carbon stocks in biomass on land converted to other land-use category, in tC per hectare and year; and

ΔC_L Annual decrease in biomass carbon stocks due to losses from harvesting, fuel wood gathering and disturbances on land converted to other land-use category, in tC per hectare and year.

Following the recommendations set in chapter 2.2.1 of the GFOI Methods Guidance Document for applying IPCC Guidelines and guidance in the context of REDD+³³, the above equation will be simplified and it will be assumed that:

- The annual change in total biomass carbon stocks (ΔC_B) is equal to the initial change in carbon stocks ($\Delta C_{CONVERSION}$);

Considering equation 2.16 of the 2006 IPCC GL for estimating ($\Delta C_{CONVERSION}$) the change of biomass carbon stocks could be expressed with the following equation:

$$\Delta C_{B_t} = \sum_{j,i} (B_{Before,j} - B_{After,i}) \times CF \times \frac{44}{12} \times A(j,i)_{RP} \quad \text{Equation 17}$$

Where:

$A(j,i)_{RP}$ Area converted/transited from forest type j to non-forest type i during the Reference Period, in hectares per year. In this case, three forest land conversions are possible:

- (Semi-)deciduous forest to Non-forest type i ;
- (Semi-)evergreen forest to Non-forest type i ; and
- Mangrove forest to Non-forest type i .

Five types of non-forest land are considered:

- Cropland (C);
- Grassland (P);
- Wetland (A);
- Settlement (U); and
- Other lands (O).
-

$B_{Before,j}$ Total biomass of forest type j before conversion/transition, in tons of dry matter per ha. This is equal to the sum of aboveground ($AGB_{Before,j}$) and belowground biomass ($BGB_{Before,j}$) and it is defined for each forest type.

$B_{After,i}$ Total biomass of non-forest type i after conversion, in tons dry matter per ha. This is equal to the sum of aboveground ($AGB_{After,i}$) and belowground biomass ($BGB_{After,i}$) and it is defined for each of the five non-forest IPCC Land Use categories.

CF Carbon fraction of dry matter in tC per ton dry matter. The value used is:

- **0.47** is the default for (sub)tropical forest as per IPCC AFOLU guidelines 2006, Table 4.3.


44/12 Conversion of C to CO₂

1.1.3.1.1.2 Activity data and emission factors used for calculating the average annual historical emissions over the Reference Period

Activity data

³³ https://www.reddcompass.org/documents/184/0/MGD2.0_English/c2061b53-79c0-4606-859f-cf6c8cc6a83

Parameter:	$A(j,i)_{RP}$
Description:	Area converted from forest type j to non-forest type i during the reference period.
Data unit:	hectare per year.
Source of data and description of measurement/calculation methods and procedures applied:	<p><i>i. Approach and source</i></p> <p>Activity data for deforestation were obtained from an annual historical time series analysis of land use, land-use change and forestry (LULUCF) carried out by five trained operators in approximately 98 effective working days (4.4 months), for the period of 2001 – 2016 across the country, using the Collect Earth Open tool.</p> <p>Activity data have been generated following IPCC Approach 3 for representing the activity data as described in the 2006 IPCC Guidelines for National Greenhouse Gas Inventories (Volume 4, Chapter 3, Section 3.13), i.e., using spatially-explicit observations of land-use categories and land-use conversions over time across the country, derived from sampling of geographically located points. The result was forest cover data for 2016 and forest cover change data for every year from 2001 to 2016.</p> <p>The period of AD analysis from 2005 to 2015 (11 years) considered for the ER in the area outside the scope of ZILMP within the Zambézia province, could be adapted within the general period 2001 – 2016 with little effort, due to the operators collecting the date of the LULC change.</p> <p><i>ii. Sampling design</i></p> <p>A systematic 4 x 4 km grid consisting of a total of 48, 894 sampling points was established at a national level to generate the historical activity data for the entire area of the country using high and medium resolution imagery, which is the same grid used to allocate the NFI clusters from the Stratified Random Sampling design. At jurisdictional level, this corresponds to 2,984 points being interpreted. Each sampling point was visually assessed and its information was collected and entered in a complete database on LULC changes at the national level.</p> <p><i>iii. Response design</i></p> <p><u>Spatial sampling unit</u></p> <p>The spatial sampling unit from each point was defined as a point with a spatial support consisting of a 100m x 100m plot (1 ha), where an internal grid of 5 x 5 points (20m x 20m grid) is overlapped. Each point from the internal grid has a weight coverage of 4% (Annex-Figure 1).</p>




Annex-Figure 1: Spatial sampling unit

Source of reference data

The sampling approach for historical AD calculation based on the regular National 4 x 4 km grid has been designed and conducted using the high and medium resolution images repository available through Google Earth and Earth Engine as a visual assessment exercise. These imagery with digital forms (Annex-Figure 2) designed to collect the LULCC information on the points of the grid are automatically accessible through the Collect Earth tool (www.openforis.org) along with scripts accessible through Earth Engine code that facilitate vegetation type's interpretation (e.g. MODIS or Landsat NDVI time series). Each point of the grid is photo-interpreted thanks to Collect Earth tool and the year and type of changes are also collected.

The use of various scripts programmed on Earth Engine Code facilitates the interpretation of the vegetation type and the determination of LULC changes. Specifically, the MOD13Q1 (NDVI 16-day Global Modis 250 m) graphic from 2001-2016, most recent Sentinel-2 image, most recent Landsat-8 pan sharpened image, Landsat-7 pan sharpened image (2000, 2004, 2008, 2012), etc.


The completeness of the series is guaranteed using remote sensing products from medium resolution imagery repositories from 2001 (e.g. Annual TOA Reflectance Composite, Annual NDVI Composite, Annual EVI Composite, Annual Greenest-Pixel TOA Reflectance Composite, etc. from Landsat 5 TM) and the most recent Sentinel-2 image from 2016. In this way, a temporal analysis of LULC changes has been completed for each sampling point of the national 4 x 4 km grid (48,894 records).



Annex-Figure 2: LULCCF detection using Collect Earth Tool (www.openforis.org). Digital forms designed with Collect Tool.

Reference labelling protocol

The activity data was generated considering the national land use and land cover classification system, which reflects the six broad IPCC Land Use categories. A set of hierarchical rules were established and used to determine the LULCCF category based on a certain percentage and taking into account the national forest definition as well (Annex-Figure 3). A single land use class is easier to classify, but it becomes challenging when there is a combination of two or more land use classes within the area of interest. Thus, this is where the hierarchical rules are important to determine the land use. Any sampling unit that has 30% of tree canopy cover is considered a forest, according to the national forest definition, even if it has more than 20% of settlements, crops or other land use, the forest is priority. In the case the sampling unit was classified as forest land and different forest types were present in the sampling unit, a majority rule was used in this case, i.e. the largest forest cover class is considered as forest type within the plot.



Annex-Figure 3: Decision tree for the attribution of the LULCCF category based on the percentage cover of the elements present in the sampling unit of 1 ha.

iv. Analysis

The estimation of the areas corresponding to a certain category changes from a forest type to a non-forest type in the framework of this systematic sampling approach was based on assessments of area proportions. According to 2006 IPCC Guidelines for National Greenhouse Gas Inventories (Volume 4, Chapter 3, Section 3.33), the proportion of each land-use or land-use change category is calculated by dividing the number of points located in the specific category by the total number of points, and area estimates for each land-use or land-use change category are obtained by multiplying the proportion of each category by the total area of interest, in this case, the area outside the scope of ZILMP within the Zambézia province.

$$A_i = p_i \times A \tag{Equation 18}$$

Where:

- A_i Area estimate on forest type j converted to non-forest type i ; hectare
- p_i Proportion of points on forest type j converted to non-forest type i ; dimensionless
- A Total area of interest; hectare

$$p_i = \frac{n_i}{N} \tag{Equation 19}$$

Where:

- n_i Number of points on forest type j converted to non-forest type i ; number
- N Total number of points; number

The standard error (ha) of an area estimate was obtained as (2006 IPCC Guidelines for National Greenhouse Gas Inventories, Volume 4, Chapter 3, Section 3.33):

$$e_i = A_i \times \sqrt{\frac{p_i \times (1 - p_i)}{N - 1}} \tag{Equation 20}$$


Where:

- A Area of interest, ha.
- p_i Proportion of points on land use change category i , dimensionless.
- n Number of sampling units, number.

	The 90% confidence interval for A_i , the estimated area of land-use category i , was given approximately by ± 1.64 times the standard error.																						
Value applied	<table border="1"> <thead> <tr> <th>Type of change</th> <th>2005-2015</th> </tr> </thead> <tbody> <tr> <td>Semi-deciduous forest to cropland</td> <td>17,505.56</td> </tr> <tr> <td>Semi-deciduous forest to grassland</td> <td>2,435.56</td> </tr> <tr> <td>Semi-deciduous forest to other lands</td> <td>0.00</td> </tr> <tr> <td>Evergreen forest to cropland</td> <td>4,566.67</td> </tr> <tr> <td>Evergreen forest to grassland</td> <td>152.22</td> </tr> <tr> <td>Evergreen forest to other lands</td> <td>152.22</td> </tr> <tr> <td>Mangrove forest to cropland</td> <td>0.00</td> </tr> <tr> <td>Mangrove forest to grassland</td> <td>152.22</td> </tr> <tr> <td>Mangrove forest to other lands</td> <td>304.44</td> </tr> </tbody> </table>	Type of change	2005-2015	Semi-deciduous forest to cropland	17,505.56	Semi-deciduous forest to grassland	2,435.56	Semi-deciduous forest to other lands	0.00	Evergreen forest to cropland	4,566.67	Evergreen forest to grassland	152.22	Evergreen forest to other lands	152.22	Mangrove forest to cropland	0.00	Mangrove forest to grassland	152.22	Mangrove forest to other lands	304.44		
Type of change	2005-2015																						
Semi-deciduous forest to cropland	17,505.56																						
Semi-deciduous forest to grassland	2,435.56																						
Semi-deciduous forest to other lands	0.00																						
Evergreen forest to cropland	4,566.67																						
Evergreen forest to grassland	152.22																						
Evergreen forest to other lands	152.22																						
Mangrove forest to cropland	0.00																						
Mangrove forest to grassland	152.22																						
Mangrove forest to other lands	304.44																						
QA/QC procedures applied:	<p>Quality Control consisted in having a team of 5 technicians with experience in forests and remote sensing, all trained together by an MRV specialist. The team worked in the same office, and discussed any classification issues with each other.</p> <p>Quality Assurance was conducted using the SAIKU extension of Collect Earth. This tool allows the detection of whether:</p> <ul style="list-style-type: none"> i) Data point was not filled ii) The class assigned followed the classification hierarchy, based on the % of individual element cover iii) Year of the Old image/Change image was less than the current image iv) Change classes are consistent with previous and current classes v) Open and closed forest was correctly classified, based on the 30% (open) and 65% (closed) cover threshold <p>In the case of any error being detected, the ID of the data point was registered and the user performed the necessary corrections.</p>																						
Uncertainty associated with this parameter:	<table border="1"> <thead> <tr> <th>Category change</th> <th>Uncertainty estimate (confidence interval at 95%)</th> </tr> </thead> <tbody> <tr> <td>Semi-deciduous forest to cropland</td> <td>17.92%</td> </tr> <tr> <td>Semi-deciduous forest to grassland</td> <td>48.88%</td> </tr> <tr> <td>Semi-deciduous forest to other lands</td> <td>-</td> </tr> <tr> <td>Evergreen forest to cropland</td> <td>35.61%</td> </tr> <tr> <td>Evergreen forest to grassland</td> <td>196.00%</td> </tr> <tr> <td>Evergreen forest to other lands</td> <td>196.00%</td> </tr> <tr> <td>Mangrove forest to cropland</td> <td>-</td> </tr> <tr> <td>Mangrove forest to grassland</td> <td>196.00%</td> </tr> <tr> <td>Mangrove forest to other lands</td> <td>138.57%</td> </tr> <tr> <td></td> <td></td> </tr> </tbody> </table>	Category change	Uncertainty estimate (confidence interval at 95%)	Semi-deciduous forest to cropland	17.92%	Semi-deciduous forest to grassland	48.88%	Semi-deciduous forest to other lands	-	Evergreen forest to cropland	35.61%	Evergreen forest to grassland	196.00%	Evergreen forest to other lands	196.00%	Mangrove forest to cropland	-	Mangrove forest to grassland	196.00%	Mangrove forest to other lands	138.57%		
Category change	Uncertainty estimate (confidence interval at 95%)																						
Semi-deciduous forest to cropland	17.92%																						
Semi-deciduous forest to grassland	48.88%																						
Semi-deciduous forest to other lands	-																						
Evergreen forest to cropland	35.61%																						
Evergreen forest to grassland	196.00%																						
Evergreen forest to other lands	196.00%																						
Mangrove forest to cropland	-																						
Mangrove forest to grassland	196.00%																						
Mangrove forest to other lands	138.57%																						
Any comment:																							

Emission factors

Parameter:	AGB _{before,j}
Description:	Aboveground biomass of forest type <i>j</i> before conversion,
Data unit:	tons of dry matter per ha
Source of data or description of the method for developing the data including the spatial level of the data (local, regional, national, international):	<p>The data used for the present document are Tier 2 (country specific data or country level estimates or locally derived estimates) and they were sourced from the NFI (for deciduous and evergreen forests) or for Mangrove forests.</p> <p>For semi-deciduous and evergreen forest, data are from the Zambézia Forest Inventory. It includes data that was collected in Zambézia province during the NFI, in 2017 and 2018. Although the inventory covers the whole province of Zambézia this is still representative of the forests located in the ZILMP as forests across the province are homogenous (floristic and structural composition). Moreover, the higher sample size of the inventory covering the whole province will enable more precise estimates for emission factors.</p> <p><i>i. Sampling design</i></p> <p>Carbon stocks before conversion for deciduous and evergreen forests were estimated using data from the National Forest Inventory sample units that were located in Zambézia province. The sample units for surveying carbon stocks were allocated using restricted stratified random sampling, using 4 * 4 km systematic grid superimposed on the agro-ecological zoning map, and stratified among the 12 forest types. Was considered as the strata, the semi-deciduous forest “open and closed”, Miombo forest “open and closed”, semi-evergreen forest “open and closed”, semi-evergreen Mountain forest “open and closed”, Mopane forest “open and closed”, and Mecrusse forest “open and closed”, of which only the first eight types occur in Zambézia province.</p> <p>The total number of sample units was determined using the optimal allocation (assuming a maximum error of 10% for the total volume, and 5% of confidence level). Proportional allocation was used to determine the number of sample units per stratum (Husch, Beers, and Kershaw 2003). For Zambézia province, 128 clusters (512 plots) were distributed between the eight (8) forest types. The cluster was used as a sampling unit, and each cluster has 4 plots of 0.1 ha (20 * 50 m), where each plot was divided into 4 sub-plots of 0.025 ha (10 * 25 m) (Annex-Figure 4).</p>



Annex-Figure 4: Design of each cluster used in the National Forest Inventory.

For estimating emission factors, the eight strata were aggregated into 2, and the similarity of the strata was used for the aggregation. The aggregation was done with the purpose of harmonizing the forest strata of the activity data with the emission factor data. Below the aggregation table.

Allocation stratum	EF Strata for MR
semi-deciduous open forest	semi-deciduous forest
semi-deciduous closed forest	
miombo open forest	
miombo closed forest	
semi-evergreen Mountain open forest	semi-evergreen forest
semi-evergreen Mountain closed forest	
semi-evergreen open forest	
semi-evergreen closed forest	

ii. Data collection

The plots were used for data collection of adult trees ($dbh \geq 10cm$), and the subplots "A" were used for data collection of established regeneration trees ($10cm > dbh \geq 5 cm$), which were included in the calculation of the carbon stocks. Data collected in the plots and subplots included tree information (dbh, scientific name, total and commercial height, stem quality), soil, forest type (this information was used to validate the information from agro-ecological zoning map), and other important information. Tree data were used to estimate above ground biomass (AGB) and below ground biomass (BGB).

The NFI did not cover Mangrove forests, so, data from the literature was used. For other strata, data from literature were also used.

Details of data collection can be found at <https://www.fnfs.gov.mz/mrv/index.php/documentos/guioes/35-directrizes-do-inventario-florestal-nacional/file>.

iii. Prediction at plot level

Above ground biomass (AGB) and below ground biomass (BGB) were estimated using a series of allometric equations adjusted for ecosystems or tree species similar to those in the Zambézia province (Annex-Table 6), and this equation was applied at tree level.

The use of the equations meant, applying allometric equations of the specific species (*Millettia stuhlmannii* taub., *Pterocarpus angolensis* DC., *Afzelia quanzensis* Welw.) in all trees of these species to estimate AGB, regardless of forest types; The allometric equation of the semi-deciduous forest was applied for all trees of this forest type (except the species mentioned above), as well as in all trees of the species *Brachystegia spiciformis* Benth., and *Julbernardia globiflora* (Benth.) Troupin to estimate AGB and BGB, because they were the main species used to adjust this equation in this forest type. The equations of the semi-evergreen forest were applied in all remaining trees of this forest type to estimate AGB; and apply the semi-deciduous forest equation in all trees to estimate the BGB in this forest type (including species mentioned above in other forest type), and apply factor 0.275 (shoot:ratio) to estimate the BGB of the semi-evergreen forest.

Annex-Table 6: List of allometric equations used to estimate above and below biomass.

Stratum	Forest type or species	Above-ground biomass (AGB) [kg]	Below-ground biomass (BGB) [kg]
Semi-deciduous forest	Semi-deciduous forest (open and closed)	$\hat{Y} = 0.0763 * DAP^{2.2046} * H^{0.4918}$	$\hat{Y} = 0.1766 * DAP^{1.7844} * H^{0.3434}$
		Author: Mugasha <i>et al.</i> (2013)	Author: Mugasha <i>et al.</i> (2013)
	<i>Millettia stuhlmannii</i> taub.	$\hat{Y} = 5.7332 * DAP^{1.4567}$	$\hat{Y} = 0.1766 * DAP^{1.7844} * H^{0.3434}$
		Author: Mate <i>et al.</i> (2014)	Author: Mugasha <i>et al.</i> (2013)
	<i>Pterocarpus angolensis</i> DC.	$\hat{Y} = 0.2201 * DAP^{2.1574}$	$\hat{Y} = 0.1766 * DAP^{1.7844} * H^{0.3434}$
		Author: Mate <i>et al.</i> (2014)	Author: Mugasha <i>et al.</i> (2013)
<i>Afzelia quanzensis</i> Welw.	$\hat{Y} = 3.1256 * DAP^{1.5833}$	$\hat{Y} = 0.1766 * DAP^{1.7844} * H^{0.3434}$	
	Author: Mate <i>et al.</i> (2014)	Author: Mugasha <i>et al.</i> (2013)	
Evergreen forest	Evergreen forest (open and closed)	$\hat{Y} = \exp(-2.289 + 2.649 \ln(DAP) - 0.021(\ln(DAP))^2)$	$\hat{Y} = AGB * R/S; \quad R/S = 0.275$
		Author: IPCC (2003)	Author: Mokany <i>et al.</i> (2006)
		$\hat{Y} = 0.0613 * DAP^{2.7133}$	$\hat{Y} = AGB * R/S; \quad R/S = 0.275$

	Evergreen MoUtain forest (open and closed)	Author: Lisboa <i>et al.</i> (2018)	Author: Mokany <i>et al.</i> (2006)							
	<i>Millettia stuhlmannii</i> taub.	$\hat{Y} = 5.7332 * DAP^{1.4567}$	$\hat{Y} = 0.1766 * DAP^{1.7844} * H^{0.3434}$							
		Author: Mate <i>et al.</i> (2014)	Author: Mugasha <i>et al.</i> (2013)							
	<i>Pterocarpus angolensis</i> DC.	$\hat{Y} = 0.2201 * DAP^{2.1574}$	$\hat{Y} = 0.1766 * DAP^{1.7844} * H^{0.3434}$							
		Author: Mate <i>et al.</i> (2014)	Author: Mugasha <i>et al.</i> (2013)							
	<i>Afzelia quanzensis</i> Welw.	$\hat{Y} = 3.1256 * DAP^{1.5833}$	$\hat{Y} = 0.1766 * DAP^{1.7844} * H^{0.3434}$							
		Author: Mate <i>et al.</i> (2014)	Author: Mugasha <i>et al.</i> (2013)							
	<p>iv. Estimation</p> <p>The estimation of mean and their respective uncertainties (standard error, sampling error, and confidence interval) for the variables biomass, carbon and carbon dioxide equivalent (above and below ground) for the two strata (semi-deciduous forest and semi-evergreen forest), were done using the forest inventory data analysis approach proposed by Bechtold & Patterson (2005) chapter 4 of the book “The Enhanced Forest Inventory and Analysis Program-National Sampling Design and Estimation Procedures”. Details of this methodology are described in Zambézia inventory report, available at https://fnds.gov.mz/mrv/index.php/documentos/relatorios/41-relatorio-de-inventario-florestal-da-zambezia-actualizado/file. For mangrove forests, data are secondary, extracted from existing literature. Stringer <i>et al.</i> (2015)³⁴ made an inventory on this ecosystem in the Zambezi delta in Mozambique; we can easily assume that carbon stocks are comparable to those of mangroves in Zambézia province. They divided mangroves into 5 strata and estimated carbon stocks in above and belowground biomass. Since we do not have information on these specific strata for ZILMP, the mean and standard error of biomass (AGB and BGB) of mangrove forest, comes indirectly from table 1 of the article by Stringer <i>et al.</i> (2015). For its determination, first the mean of carbon was found for the two pools (sum of overstory and understory carbon) for each stratum (Height Class 1, ..., Height Class 5), followed by the calculation of the mean of the ecosystem (mean weighted according to the stratum areas). Finally, the carbon was converted to biomass using the conversion factor of 0.47 proposed in the IPCC good practice guide.</p> <p>Spatial level: Regional</p> <p>Spatial level: Regional</p>									
	Value applied:	<table border="1"> <tr> <td>Semi-deciduous forest (FSD)</td> <td>144.69</td> </tr> <tr> <td>Evergreen forest (FSSV)</td> <td>123.13</td> </tr> <tr> <td>Mangrove forest (FF)</td> <td>269.01</td> </tr> </table>			Semi-deciduous forest (FSD)	144.69	Evergreen forest (FSSV)	123.13	Mangrove forest (FF)	269.01
	Semi-deciduous forest (FSD)	144.69								
Evergreen forest (FSSV)	123.13									
Mangrove forest (FF)	269.01									

³⁴ Stringer, C. E.; Trettin, C. C.; Zarnoch, S. J. and Tang, W. 2015. Carbon stocks of mangroves within the Zambezi River Delta, Mozambique. *Forest Ecology Management* 354:139–148.

	The values above are estimated and extracted in the " Emission factor v.2 " workbook, and then they are recorded in the cells "B4", "B10" and "B16" respectively, of the "BIOMASS" worksheet tab in the " Outside ZILMP Emissions Calculations RL (2005 2015) 28 10 20 " workbook. These values are then applied in the range "C9:C26" of the "Reference Level" worksheet tab in the " Outside ZILMP Emissions Calculations RL (2005 2015) 28 10 20 " workbook for estimating emissions.								
QA/QC procedures applied	The QA/QC procedures consisted on the following: <ul style="list-style-type: none"> • SOPs were developed. • A training on the SOPs was conducted prior to the field work. This training lasted for 3 weeks, and consisted of training on the usage of all equipment and evaluating the specific skills of each participant, in order to determine the team and brigade leaders. On the start of the 2nd phase of the IFN (2017) an additional 1-week training was conducted, to refresh the participants and train any new members. • The supervisor of each inventory team conducted a remeasurement of 4 trees per plot which means 16 trees per cluster. This served to ensure that the SOPs were adequately implemented. • An independent measurement of 10% of the plots. This activity was conducted by technicians of the National Directorate of Forests, who had participated in the Provincial Inventories of Gaza and Cabo Delgado. Diameter below 10%. • The adequacy of the allometric models, including root-to-shoot ratios used was confirmed by experts of the Faculty of Agronomy and Forest Engineering (FAEF) and the Department of Biology Sciences (DCB) of the Eduardo Mondlane University (UEM). • The World Bank conducted two regular supervision missions of the National Forest Inventories to confirm the adequate implementation of the SOPs and suggest areas for improvement. The report can be found here. • An independent expert (Jim Alegria, ex-US Forestry Service) was hired in order to evaluate the methodology for the inventory and support in the estimation step. The report can be found here. 								
Uncertainty associated with this parameter:	<table border="1"> <thead> <tr> <th>Forest type</th> <th>Uncertainty estimate (confidence interval at 95%)</th> </tr> </thead> <tbody> <tr> <td>FSD</td> <td>19.72%</td> </tr> <tr> <td>FSSV</td> <td>18.33%</td> </tr> <tr> <td>FF</td> <td>8.00%</td> </tr> </tbody> </table>	Forest type	Uncertainty estimate (confidence interval at 95%)	FSD	19.72%	FSSV	18.33%	FF	8.00%
Forest type	Uncertainty estimate (confidence interval at 95%)								
FSD	19.72%								
FSSV	18.33%								
FF	8.00%								
Any comment:	-								

Parameter:	BGB _{before,j}
Description:	Belowground biomass of forest type <i>j</i> before conversion,
Data unit:	tons of dry matter per ha
Source of data or description of the method for developing the data including the spatial level of the data (local, regional,	<p>For semi-deciduous and evergreen forest, data are from the Zambézia Forest Inventory. It includes data that was collected in Zambézia province during the NFI, in 2017 and 2018. Please refer to parameter AGB_{before,j} for more information.</p> <p>For mangrove forests, please refer to parameter AGB_{before,j} for more information.</p> <p>Spatial level: Regional</p>

national, international):									
Value applied:	<table border="1"> <tr> <td>Semi-deciduous forest (FSD)</td> <td>49.95</td> </tr> <tr> <td>Evergreen forest (FSSV)</td> <td>42.06</td> </tr> <tr> <td>Mangrove forest (FF)</td> <td>85.43</td> </tr> </table> <p>The values above are estimated and extracted in the workbook "Emission factor v.2" and then they are recorded in the cells "B45", "B51" and "B57" respectively, of the "BIOMASS" worksheet tab in the "Outside ZILMP Emissions Calculations RL (2005 2015) 28 10 20" workbook. These values are then applied in the range "E9:E26" of the "Reference Level" worksheet tab in the "Outside ZILMP Emissions Calculations RL (2005 2015) 28 10 20" workbook for estimating emissions.</p>	Semi-deciduous forest (FSD)	49.95	Evergreen forest (FSSV)	42.06	Mangrove forest (FF)	85.43		
Semi-deciduous forest (FSD)	49.95								
Evergreen forest (FSSV)	42.06								
Mangrove forest (FF)	85.43								
QA/QC procedures applied	Please see section QA/QC procedures under parameter $AGB_{before,j}$.								
Uncertainty associated with this parameter:	<table border="1"> <thead> <tr> <th>Forest type</th> <th>Uncertainty estimate (confidence interval at 95%)</th> </tr> </thead> <tbody> <tr> <td>FSD</td> <td>16.58%</td> </tr> <tr> <td>FSSV</td> <td>16.71%</td> </tr> <tr> <td>FF</td> <td>10.00%</td> </tr> </tbody> </table>	Forest type	Uncertainty estimate (confidence interval at 95%)	FSD	16.58%	FSSV	16.71%	FF	10.00%
Forest type	Uncertainty estimate (confidence interval at 95%)								
FSD	16.58%								
FSSV	16.71%								
FF	10.00%								
Any comment:									

Parameter:	$AGB_{after,i}$						
Description:	Aboveground biomass of non-forest type i after conversion						
Data unit:	tons of dry matter per ha						
Source of data or description of the method for developing the data including the spatial level of the data (local, regional, national, international):	<p>For cropland: The values and assumptions of 2006 IPCC GL, Volume 4, Chapter 5 are used because, unfortunately, there aren't country-specific data. The agricultural land in Mozambique is mostly under the annual-crop farming practices that drive conversion of forest land to agricultural lands. So, according to 2006 IPCC GL (Volume 4, Chapter 5, Section 5.28), for lands planted in annual crops, the default value of growth in crops planted after conversion is 5 tonnes of C per hectare, based on the original IPCC Guidelines recommendation of 10 tonnes of dry biomass per hectare (dry biomass has been converted to tonnes carbon in Table 5.9) (2006 IPCC, Volume 4, Chapter 5, Section 5.28).</p> <p>For grassland: The values and assumptions of 2006 IPCC GL, Volume 4, Chapter 6 are used because, unfortunately, there aren't country-specific data. As the climate in most of Mozambique is tropical dry to subtropical dry, the value of peak-above ground biomass for tropical dry of TABLE 6.4 is assumed.</p> <p>For other lands: No default values exist for these conversions.</p> <p>Spatial level: International</p>						
Value applied:	<table border="1"> <tr> <td>Cropland (C)</td> <td>10</td> </tr> <tr> <td>Grassland (P)</td> <td>2.3</td> </tr> <tr> <td>Other lands (A O U)</td> <td>0.0</td> </tr> </table> <p>The values above are recorded in the ranges "B5:B9", "B11:B15" and "B17:B21" of the "BIOMASS" worksheet tab in the "Outside ZILMP Emissions Calculations RL (2005 2015) 28 10 20" workbook. These values</p>	Cropland (C)	10	Grassland (P)	2.3	Other lands (A O U)	0.0
Cropland (C)	10						
Grassland (P)	2.3						
Other lands (A O U)	0.0						

	are then applied in the range "D9:D26" of the "Reference Level" worksheet tab in the " "Outside ZILMP Emissions Calculations RL (2005 2015) 28 10 20" " workbook for estimating emissions.								
QA/QC procedures applied	The adequacy in the use of these default values was confirmed with the experts in GHG Inventory in DINAB.								
Uncertainty associated with this parameter:	<table border="1"> <thead> <tr> <th>Non-forest type</th> <th>Uncertainty estimate (Confidence interval at 95%)</th> </tr> </thead> <tbody> <tr> <td>Cropland (C)</td> <td>75.00%</td> </tr> <tr> <td>Grassland (P)</td> <td>75.00%</td> </tr> <tr> <td>Other lands (A O U)</td> <td>-</td> </tr> </tbody> </table>	Non-forest type	Uncertainty estimate (Confidence interval at 95%)	Cropland (C)	75.00%	Grassland (P)	75.00%	Other lands (A O U)	-
Non-forest type	Uncertainty estimate (Confidence interval at 95%)								
Cropland (C)	75.00%								
Grassland (P)	75.00%								
Other lands (A O U)	-								
Any comment:									

Parameter:	$BGB_{after,i}$						
Description:	Belowground biomass of non-forest type <i>i</i> after conversion						
Data unit:	tons of dry matter per ha						
Source of data or description of the method for developing the data including the spatial level of the data (local, regional, national, international):	<p>For cropland: The values and assumptions of 2006 IPCC GL, Volume 4, Chapter 5 are used. Tier 2 may modify the assumption that carbon stocks immediately following conversion are zero. In this case, it is assumed that conversion leads to annual croplands and in the case the carbon stock in biomass after one year for annual crops provided in TABLE 5.9 is used.</p> <p>For grassland: The values and assumptions of 2006 IPCC GL, Volume 4, Chapter 6, TABLE 6.1 and TABLE 6.4 are used because, unfortunately, there aren't country-specific data. As the climate in most of Mozambique is tropical dry to subtropical dry, the value for semi-arid grassland in tropical dry climate zone is used, therefore a root-shoot ratio of 2.8 (TABLE 6.1) is applied to the value of peak above-ground biomass, 2.3 tonnes of dry biomass per hectare (TABLE 6.4), generating the expected values 6.4 tonnes of dry biomass per hectare.</p> <p>For other lands: No default values exist for these conversions.</p> <p>Spatial level: International</p>						
Value applied:	<table border="1"> <tbody> <tr> <td>Cropland (C)</td> <td>0.00</td> </tr> <tr> <td>Grassland (P)</td> <td>6.44</td> </tr> <tr> <td>Other lands (A O U)</td> <td>0.00</td> </tr> </tbody> </table> <p>The values above are recorded in the ranges "B46:E50", "B52:B56" and "B58:B62" of the "BIOMASS" worksheet tab in the ""Outside ZILMP Emissions Calculations RL (2005 2015) 28 10 20"" workbook. These values are then applied in the range "F9:F26" of the "Reference Level" worksheet tab in the ""Outside ZILMP Emissions Calculations RL (2005 2015) 28 10 20"" workbook for estimating emissions.</p>	Cropland (C)	0.00	Grassland (P)	6.44	Other lands (A O U)	0.00
Cropland (C)	0.00						
Grassland (P)	6.44						
Other lands (A O U)	0.00						
QA/QC procedures applied	The adequacy in the use of these default values was confirmed with the experts in GHG Inventory in DINAB.						

Uncertainty associated with this parameter:		
	Non-forest type	Uncertainty estimate (Confidence interval at 95%)
	Cropland (C)	-
	Grassland (P)	121.04%
	Other lands (A O U)	-
Any comment:		

1.1.3.1.1.3 Calculation of the average annual historical emissions over the Reference Period

The following table shows the average annual historical emissions results obtained per category changes from a forest type to a non-forest type over the Reference Period. The emissions are generated relating the data and parameters described above (Activity data and Emission Factors) and summarized in the Annex-Table 7, by applying Equation 17.

Annex-Table 7: Calculation of the average annual historical emissions over the Reference Period.

Category changes	Average annual historical activity data _{j,i} (ha/yr)	AGB _{before,j} (tdm/ha)	BGB _{before,j} (tdm/ha)	AGB _{after,i} (tdm/ha)	BGB _{after,i} (tdm/ha)	Average annual historical emissions (tCO _{2e} /yr)
Semi-deciduous forest to cropland	17,505.56	144.69	49.4995	10.00	0.00	5,570,060.47
Semi-deciduous forest to grassland	2,435.56	144.69	49.4995	2.30	6.44	780,253.50
Semi-deciduous forest to other lands	0.00	144.69	49.4995	0.00	0.00	0.00
Evergreen forest to cropland	4,566.67	123.13	42.0626	10.00	0.00	1,221,308.32
Evergreen forest to grassland	152.22	123.13	42.066	2.30	6.44	41,040.81
Evergreen forest to other lands	152.22	123.13	42.0626	0.00	0.00	43,333.57
Mangrove to cropland	0.00	269.01	85.43	0.00	0.00	0.00
Mangrove to grassland	152.22	269.01	85.43	2.30	6.44	90,687.38
Mangrove to other lands	304.44	269.01	85.43	0.00	0.00	185,960.28
Total						7,932,644.34

5.2.4 Estimated Reference Level

ER Program Reference level

Crediting Period year t	Average annual historical emissions from deforestation over the Reference Period (tCO _{2-e} /yr)	If applicable, average annual historical emissions from forest degradation over the Reference Period (tCO _{2-e} /yr)	If applicable, average annual historical removals by sinks over the Reference Period (tCO _{2-e} /yr)	Adjustment, if applicable (tCO _{2-e} /yr)	Reference level (tCO _{2-e} /yr)
2018	7,932,644.34	-	-	-	7,932,644.34
2019	7,932,644.34	-	-	-	7,932,644.34
2020	7,932,644.34	-	-	-	7,932,644.34
2021	7,932,644.34	-	-	-	7,932,644.34
2022	7,932,644.34	-	-	-	7,932,644.34
2023	7,932,644.34	-	-	-	7,932,644.34
2024	7,932,644.34	-	-	-	7,932,644.34


5.3 MONITORING AND REPORTING PERIOD

The monitoring and reporting period covers emissions in 2021 outside of ZILMP program.

5.3.1 Measurement, monitoring and reporting approach

1.1.4 Line Diagram

The Annex-Figure 5 illustrates the emissions reductions calculation workflow during the Monitoring Period.



Annex-Figure 5: Emissions reductions calculation workflow.

1.1.5 Calculation

$$ER_{ERP,t} = RL_t - GHG_t \quad \text{Equation 21}$$

Where:

ER_{ERP}	=	Emission Reductions under the area outside the scope of ZILMP in year t ; $tCO_2e*year^{-1}$.
RL_{RP}	=	Gross emissions of the RL from deforestation over the Reference Period; $tCO_2e*year^{-1}$.
GHG_t	=	Monitored gross emissions from deforestation at year t ; $tCO_2e*year^{-1}$;
T	=	Number of years during the monitoring period; <i>dimensionless</i> .

Reference Level (RL_t)

Gross emissions of the RL from deforestation over the Reference Period (RL_{RP}) are estimated as the sum of annual change in total biomass carbon stocks (ΔC_{B_t}) during the reference period.

$$RL_{RP} = \frac{\sum_t^{RP} \Delta C_{B_t}}{RP} \quad \text{Equation 22}$$

Where:

ΔC_{B_t}	=	Annual change in total biomass carbon stocks at year t ; $tC*year^{-1}$;
RP	=	Reference period; <i>years</i> .

Following the 2006 IPCC Guidelines, the annual change in total biomass carbon stocks forest land converted to other land-use category (ΔC_{B_t}) would be estimated through the following equation:

$$\Delta C_{B_t} = \Delta C_G + \Delta C_{CONVERSION} - \Delta C_L \quad \text{Equation 23}$$

Where:

ΔC_{B_t}	Annual change of total biomass carbon stocks during the period, in tC per year;
ΔC_G	Annual increase in carbon stocks in biomass due to growth on land converted to another land-use category, in tC per hectare and year;
$\Delta C_{CONVERSION}$	Initial change in carbon stocks in biomass on land converted to other land-use category, in tC per hectare and year; and
ΔC_L	Annual decrease in biomass carbon stocks due to losses from harvesting, fuel wood gathering and disturbances on land converted to other land-use category, in tC per hectare and year.

Following the recommendations set in chapter 2.2.1 of the GFOI Methods Guidance Document for applying IPCC Guidelines and guidance in the context of REDD+³⁵, the above equation will be simplified and it will be assumed that:

- The annual change in total biomass carbon stocks (ΔC_B) is equal to the initial change in carbon stocks ($\Delta C_{CONVERSION}$);

Considering equation 2.16 of the 2006 IPCC GL for estimating ($\Delta C_{CONVERSION}$) the change of biomass carbon stocks could be expressed with the following equation:

³⁵ <https://www.reddcompass.org/mgd>

$$\Delta C_{B_t} = \sum_{j,i} (B_{Before,j} - B_{After,i}) \times CF \times \frac{44}{12} \times A(j,i)_{RP} \quad \text{Equation 24}$$

Where:

$A(j,i)_{RP}$ Area converted/transited from forest type j to non-forest type i during the Reference Period, in hectares per year. In this case, three forest land conversions are possible:

- (Semi-)deciduous forest to Non-forest type i ;
- (Semi-)evergreen forest to Non-forest type i ; and
- Mangrove forest to Non-forest type i .

Five types of non-forest land are considered:

- Cropland (C);
- Grassland (P);
- Wetland (A);
- Settlement (U); and
- Other lands (O).

$B_{Before,j}$ Total biomass of forest type j before conversion/transition, in tons of dry matter per ha. This is equal to the sum of aboveground ($AGB_{Before,j}$) and belowground biomass ($BGB_{Before,j}$) and it is defined for each forest type.

$B_{After,i}$ Total biomass of non-forest type i after conversion, in tons dry matter per ha. This is equal to the sum of aboveground ($AGB_{After,i}$) and belowground biomass ($BGB_{After,i}$) and it is defined for each of the five non-forest IPCC Land Use categories.

CF Carbon fraction of dry matter in tC per ton dry matter. The value used is:

- **0.47** is the default for (sub)tropical forest as per IPCC AFOLU guidelines 2006, Table 4.3.

44/12 Conversion of C to CO₂

Monitored emissions (GHG_t)

Annual gross GHG emissions over the monitoring period in the Accounting Area (GHG_t) are estimated as the sum of annual change in total biomass carbon stocks (ΔC_{B_t}).

$$GHG_t = \frac{\sum_t \Delta C_{B_t}}{T} \quad \text{Equation 25}$$

Where:

ΔC_{B_t} = Annual change in total biomass carbon stocks at year t ; $tC \cdot year^{-1}$

T = Number of years during the monitoring period; *dimensionless*.

Changes in total biomass carbon stocks

Following the 2006 IPCC Guidelines, the annual change in total biomass carbon stocks forest land converted to other land-use category (ΔC_B) would be estimated through **Equation 23**. Making the same assumptions as described above for the RL the change of biomass carbon stocks could be expressed with the following equation:

$$\Delta C_B = \sum_{j,i} (B_{Before,j} - B_{After,i}) \times CF \times \frac{44}{12} \times A(j,i)_{MP} \quad \text{Equation 26}$$

Where:

$A(j, i)_{MP}$	<p>Area converted/transited from forest type j to non-forest type i during the Monitoring Period, in hectare per year. In this case, three forest land conversions are possible:</p> <ul style="list-style-type: none"> • (Semi-)deciduous forest to Non-forest type i; • (Semi-)evergreen forest to Non-forest type i; and • Mangrove forest to Non-forest type i. <p>Five types of non-forest land are considered:</p> <ul style="list-style-type: none"> • Cropland (C); • Grassland (P); • Wetland (A); • Settlement (U); and • Other lands (O).
$B_{Before,j}$	Total biomass of forest type j before conversion/transition, in tons of dry matter per ha. This is equal to the sum of aboveground ($AGB_{Before,j}$) and belowground biomass ($BGB_{Before,j}$) and it is defined for each forest type.
$B_{After,i}$	Total biomass of non-forest type i after conversion, in <i>tons dry matter per ha</i> . This is equal to the sum of aboveground ($AGB_{After,i}$) and belowground biomass ($BGB_{After,i}$) and it is defined for each of the five non-forest IPCC Land Use categories.
CF	Carbon fraction of dry matter in tC per ton dry matter. The value used is: <ul style="list-style-type: none"> • 0.47 is the default for (sub)tropical forest as per IPCC AFOLU guidelines 2006, Table 4.3.
44/12	Conversion of C to CO ₂

5.3.2 Data and parameters

1.1.6 Fixed Data and Parameters

Parameter:	AGB _{before,j}										
Description:	Aboveground biomass of forest type <i>j</i> before conversion,										
Data unit:	tons of dry matter per ha										
Source of data or description of the method for developing the data including the spatial level of the data (local, regional, national, international):	<p>The data used for the present document are Tier 2 (country specific data or country level estimates or locally derived estimates) and they were sourced from the NFI (for deciduous and evergreen forests) or for Mangrove forests.</p> <p>For semi-deciduous and evergreen forest, data are from the Zambézia Forest Inventory. It includes data that was collected in Zambézia province during the NFI, in 2017 and 2018. Although the inventory covers the whole province of Zambézia this is still representative of the forests located in the ZILMP as forests across the province are homogenous (floristic and structural composition). Moreover, the higher sample size of the inventory covering the whole province will enable more precise estimates for emission factors.</p> <p><i>i. Sampling design</i></p> <p>Carbon stocks before conversion for deciduous and evergreen forests were estimated using data from the National Forest Inventory sample units that were located in Zambézia province. The sample units for surveying carbon stocks were allocated using restricted stratified random sampling, using 4 * 4 km systematic grid superimposed on the agro-ecological zoning map, and stratified among the 12 forest types. Was considered as the strata, the semi-deciduous forest “open and closed”, Miombo forest “open and closed”, semi-evergreen forest “open and closed”, semi-evergreen Mountain forest “open and closed”, Mopane forest “open and closed”, and Mecrusse forest “open and closed”, of which only the first eight types occur in Zambézia province.</p> <p>The total number of sample units was determined using the optimal allocation (assuming a maximum error of 10% for the total volume, and 5% of confidence level). Proportional allocation was used to determine the number of sample units per stratum (Husch, Beers, and Kershaw 2003). For Zambézia province, 128 clusters (512 plots) were distributed between the eight (8) forest types. The cluster was used as a sampling unit, and each cluster has 4 plots of 0.1 ha (20 * 50 m), where each plot was divided into 4 sub-plots of 0.025 ha (10 * 25 m) (Figure 6).</p> <p>For estimating emission factors, the eight strata were aggregated into 2, and the similarity of the strata was used for the aggregation. The aggregation was done with the purpose of harmonizing the forest strata of the activity data with the emission factor data. Below the aggregation table.</p> <table border="1" data-bbox="411 1715 1150 2016"> <thead> <tr> <th>Allocation stratum</th> <th>EF Strata for MR</th> </tr> </thead> <tbody> <tr> <td>semi-deciduous open forest</td> <td rowspan="4">semi-deciduous forest</td> </tr> <tr> <td>semi-deciduous closed forest</td> </tr> <tr> <td>miombo open forest</td> </tr> <tr> <td>miombo closed forest</td> </tr> <tr> <td>semi-evergreen Mountain open forest</td> <td rowspan="2">semi-evergreen forest</td> </tr> <tr> <td>semi-evergreen Mountain closed forest</td> </tr> </tbody> </table>	Allocation stratum	EF Strata for MR	semi-deciduous open forest	semi-deciduous forest	semi-deciduous closed forest	miombo open forest	miombo closed forest	semi-evergreen Mountain open forest	semi-evergreen forest	semi-evergreen Mountain closed forest
Allocation stratum	EF Strata for MR										
semi-deciduous open forest	semi-deciduous forest										
semi-deciduous closed forest											
miombo open forest											
miombo closed forest											
semi-evergreen Mountain open forest	semi-evergreen forest										
semi-evergreen Mountain closed forest											

semi-evergreen open forest	
semi-evergreen closed forest	

ii. Data collection

The plots were used for data collection of adult trees (dbh≥10cm), and the subplots "A" were used for data collection of established regeneration trees (10cm> dbh≥ 5 cm), which were included in the calculation of the carbon stocks. Data collected in the plots and subplots included tree information (dbh, scientific name, total and commercial height, stem quality), soil, forest type (this information was used to validate the information from agro-ecological zoning map), and other important information. Tree data were used to estimate above ground biomass (AGB) and below ground biomass (BGB).

The NFI did not cover Mangrove forests, so, data from the literature was used. For other strata, data from literature were also used.

Details of data collection can be find at <https://www.fnds.gov.mz/mrv/index.php/documentos/guioes/35-directrizes-do-inventario-florestal-nacional/file> .

iii. Prediction at plot level

Above ground biomass (AGB) and below ground biomass (BGB) were estimated using a series of allometric equations adjusted for ecosystems or tree species similar to those in the Zambézia province (Table 1), and this equation was applied at tree level.

The use of the equations meant, applying allometric equations of the specific species (*Millettia stuhlmannii* taub., *Pterocarpus angolensis* DC., *Afzelia quanzensis* Welw.) in all trees of these species to estimate AGB, regardless of forest types. The allometric equation of the semi-deciduous forest was applied for all trees of this forest type (except the species mentioned above), as well as in all trees of the species *Brachystegia spiciformis* Benth., and *Julbernardia globiflora* (Benth.) Troupin to estimate AGB and BGB, because they were the main species used to adjust this equation in this forest type. The equations of the semi-evergreen forest were applied in all remaining trees of this forest type to estimate AGB; and apply the semi-deciduous forest equation in all trees to estimate the BGB in this forest type (including species mentioned above in other forest type), and apply factor 0.28 (shoot ratio) to estimate the BGB of the semi-evergreen forest.

Since Mozambique was not able to propagate this source of error through Monte Carlo (MC) simulation we have increased the sampling uncertainty of AGB and BGB for forest strata by 10% at 90% confidence level using the quadrature approach and the combined error was propagated in the MC simulation.

iv. Estimation

The estimation of mean and their respective uncertainties (standard error, sampling error, and confidence interval) for the variables biomass, carbon and carbon dioxide equivalent (above and below ground) for the two strata (semi-deciduous forest and semi-evergreen forest), were done using the forest inventory data analysis approach proposed by Bechtold & Patterson (2005) chapter 4 of the book "The Enhanced Forest Inventory and Analysis Program-National Sampling Design and Estimation Procedures". Details of this methodology are described in Zambézia inventory report, available at <https://fnds.gov.mz/mrv/index.php/documentos/relatorios/41-relatorio-de-inventario-florestal-da-zambezia-actualizado/file>. For mangrove forests, data are secondary, extracted

	<p>from existing literature. Stringer <i>et al.</i> (2015)³⁶ made an inventory on this ecosystem in the Zambezi delta in Mozambique; we can easily assume that carbon stocks are comparable to those of mangroves in Zambézia province. They divided mangroves into 5 strata and estimated carbon stocks in above and belowground biomass. Since we do not have information on these specific strata for ZILMP, the mean and standard error of biomass (AGB and BGB) of mangrove forest, comes indirectly from table 1 of the article by Stringer <i>et al.</i> (2015). For its determination, first the mean of carbon was found for the two pools (sum of overstory and understory carbon) for each stratum (Height Class 1, ..., Height Class 5), followed by the calculation of the mean of the ecosystem (mean weighted according to the stratum areas). Finally, the carbon was converted to biomass using the conversion factor of 0.47 proposed in the IPCC good practice guide.</p> <p>Spatial level: Regional</p>						
<p>Value applied:</p>	<table border="1" data-bbox="691 647 1153 750"> <tr> <td>Semi-deciduous forest (FSD)</td> <td>144.69</td> </tr> <tr> <td>Evergreen forest (FSSV)</td> <td>123.13</td> </tr> <tr> <td>Mangrove forest (FF)</td> <td>269.01</td> </tr> </table> <p>The values above are estimated and extracted in the "Emission factor v.2" workbook, and then they are recorded in the cells "B4", "B10" and "B16" respectively, of the "BIOMASS" worksheet tab in the "Outside ZILMP Emissions Calculations MR (2021)" and "Outside ZILMP Emissions Calculations MR (2022)" workbooks. These values are then applied in the range "C9:C20" of the "EMISSION MONITORING PERIOD(EMP)" worksheet tab in the "Outside ZILMP Emissions Calculations MR (2021)" and "Outside ZILMP Emissions Calculations MR (2022)" workbooks for estimating emissions.</p>	Semi-deciduous forest (FSD)	144.69	Evergreen forest (FSSV)	123.13	Mangrove forest (FF)	269.01
Semi-deciduous forest (FSD)	144.69						
Evergreen forest (FSSV)	123.13						
Mangrove forest (FF)	269.01						
<p>QA/QC procedures applied</p>	<p>The QA/QC procedures consisted on the following:</p> <ul style="list-style-type: none"> • SOPs were developed as described in <i>Section 2.1 - National Forest Inventory</i>. • A training on the SOPs was conducted prior to the field work. This training lasted for 3 weeks, and consisted of training on the usage of all equipment and evaluating the specific skills of each participant, in order to determine the team and brigade leaders. On the start of the 2nd phase of the IFN (2017) an additional 1-week training was conducted, to refresh the participants and train any new members. • The supervisor of each inventory team conducted a remeasurement of 4 trees per plot which means 16 trees per cluster. This served to ensure that the SOPs were adequately implemented. • An independent measurement of 10% of the plots. This activity was conducted by technicians of the National Directorate of Forests, who had participated in the Provincial Inventories of Gaza and Cabo Delgado. Diameter error must be below 10%. • The adequacy of the allometric models, including root-to-shoot ratios used was confirmed by experts of the Faculty of Agronomy and Forest Engineering (FAEF) and the Department of Biology Sciences (DCB) of the University Eduardo Mondlane (UEM). • The World Bank conducted two regular supervision missions of the National Forest Inventories to confirm the adequate implementation of the SOPs and suggest areas for improvement. The report can be found here. • An independent expert (Jim Alegria, ex-US Forestry Service) was hired in order to evaluate the methodology for the inventory and support in the estimation step. The report can be found here. 						
<p>Uncertainty associated with this parameter:</p>	<table border="1" data-bbox="678 1816 1165 1910"> <thead> <tr> <th>Forest type</th> <th>Uncertainty estimate (confidence interval at 95%)</th> </tr> </thead> <tbody> <tr> <td></td> <td></td> </tr> </tbody> </table>	Forest type	Uncertainty estimate (confidence interval at 95%)				
Forest type	Uncertainty estimate (confidence interval at 95%)						

³⁶ Stringer, C. E.; Trettin, C. C.; Zarnoch, S. J. and Tang, W. 2015. Carbon stocks of mangroves within the Zambezi River Delta, Mozambique. *Forest Ecology Management* 354:139–148.

		FSD	19.72%	
		FSSV	18.33%	
		FF	8.00%	
Any comment:	-			

Parameter:	BGB _{before,j}								
Description:	Belowground biomass of forest type <i>j</i> before conversion,								
Data unit:	tons of dry matter per ha								
Source of data or description of the method for developing the data including the spatial level of the data (local, regional, national, international):	<p>For semi-deciduous and evergreen forest, data are from the Zambézia Forest Inventory. It includes data that was collected in Zambézia province during the NFI, in 2017 and 2018. Please refer to parameter AGB_{before,j} for more information.</p> <p>For mangrove forests, please refer to parameter AGB_{before,j} for more information.</p> <p>Spatial level: Regional</p>								
Value applied:	<table border="1" data-bbox="699 1048 1147 1151"> <tr> <td>Semi-deciduous forest (FSD)</td> <td>49.95</td> </tr> <tr> <td>Evergreen forest (FSSV)</td> <td>42.06</td> </tr> <tr> <td>Mangrove forest (FF)</td> <td>85.43</td> </tr> </table> <p>The values above are estimated and extracted in the workbook "Emission factor v.2" and then they are recorded in the cells "B46", "B52" and "B58" respectively, of the "BIOMASS" worksheet tab in the "Outside ZILMP Emissions Calculations MR (2021)" and "Outside ZILMP Emissions Calculations MR (2022)" workbook. These values are then applied in the range "E9:E20" of the "EMISSION MONITORING PERIOD(ERL)" worksheet tab in the "Outside ZILMP Emissions Calculations MR (2021)" and "Outside ZILMP Emissions Calculations MR (2022)" workbook for estimating emissions.</p>	Semi-deciduous forest (FSD)	49.95	Evergreen forest (FSSV)	42.06	Mangrove forest (FF)	85.43		
Semi-deciduous forest (FSD)	49.95								
Evergreen forest (FSSV)	42.06								
Mangrove forest (FF)	85.43								
QA/QC procedures applied	Please see section QA/QC procedures under parameter AGB _{before,j} .								
Uncertainty associated with this parameter:	<table border="1" data-bbox="678 1579 1165 1765"> <thead> <tr> <th>Forest type</th> <th>Uncertainty estimate (confidence interval at 95%)</th> </tr> </thead> <tbody> <tr> <td>FSD</td> <td>16.58%</td> </tr> <tr> <td>FSSV</td> <td>16.71%</td> </tr> <tr> <td>FF</td> <td>10.00%</td> </tr> </tbody> </table>	Forest type	Uncertainty estimate (confidence interval at 95%)	FSD	16.58%	FSSV	16.71%	FF	10.00%
Forest type	Uncertainty estimate (confidence interval at 95%)								
FSD	16.58%								
FSSV	16.71%								
FF	10.00%								
Any comment:	-								

Parameter:	AGB _{after,i}
Description:	Aboveground biomass of non-forest type <i>i</i> after conversion
Data unit:	tons of dry matter per ha

Source of data or description of the method for developing the data including the spatial level of the data (local, regional, national, international):	<p>For cropland: The values and assumptions of 2006 IPCC GL, Volume 4, Chapter 5 are used because, unfortunately, there aren't country-specific data. The agricultural land in Mozambique is mostly under the annual-crop farming practices that drive conversion of forest land to agricultural lands. So, according to 2006 IPCC GL (Volume 4, Chapter 5, Section 5.28), for lands planted in annual crops, the default value of growth in crops planted after conversion is 5 tonnes of C per hectare, based on the original IPCC Guidelines recommendation of 10 tonnes of dry biomass per hectare (dry biomass has been converted to tonnes carbon in Table 5.9) (2006 IPCC, Volume 4, Chapter 5, Section 5.28).</p> <p>For grassland: The values and assumptions of 2006 IPCC GL, Volume 4, Chapter 6 are used because, unfortunately, there aren't country-specific data. As the climate in most of Mozambique is tropical dry to subtropical dry, the value of peak-above ground biomass for tropical dry of TABLE 6.4 is assumed.</p> <p>For other lands: No default values exist for these conversions.</p> <p>Spatial level: International</p>								
Value applied:	<table border="1" style="margin-left: auto; margin-right: auto;"> <tr> <td style="background-color: #d9ead3;">Cropland (C)</td> <td style="text-align: center;">10</td> </tr> <tr> <td style="background-color: #d9ead3;">Grassland (P)</td> <td style="text-align: center;">2.3</td> </tr> <tr> <td style="background-color: #d9ead3;">Other lands (A O U)</td> <td style="text-align: center;">0.0</td> </tr> </table> <p>The values above are recorded in the ranges "B5:B9", "B11:B15" and "B17:B21" of the "BIOMASS" worksheet tab in the "Outside ZILMP Emissions Calculations MR (2021)" and "Outside ZILMP Emissions Calculations MR (2022)" workbook. These values are then applied in the range "D9:D20" of the "EMISSION MONITORING PERIOD(ERL)" worksheet tab in the "Outside ZILMP Emissions Calculations MR (2021)" and "Outside ZILMP Emissions Calculations MR (2022)" workbook for estimating emissions.</p>	Cropland (C)	10	Grassland (P)	2.3	Other lands (A O U)	0.0		
Cropland (C)	10								
Grassland (P)	2.3								
Other lands (A O U)	0.0								
QA/QC procedures applied	The adequacy in the use of these default values was confirmed with the experts in GHG Inventory in DINAB.								
Uncertainty associated with this parameter:	<table border="1" style="margin-left: auto; margin-right: auto;"> <thead> <tr> <th style="background-color: #d9ead3;">Non-forest type</th> <th style="background-color: #d9ead3;">Uncertainty estimate (confidence interval at 95%)</th> </tr> </thead> <tbody> <tr> <td>Cropland (C)</td> <td style="text-align: center;">75.00%</td> </tr> <tr> <td>Grassland (P)</td> <td style="text-align: center;">75.00%</td> </tr> <tr> <td>Other lands (A O U)</td> <td style="text-align: center;">-</td> </tr> </tbody> </table>	Non-forest type	Uncertainty estimate (confidence interval at 95%)	Cropland (C)	75.00%	Grassland (P)	75.00%	Other lands (A O U)	-
Non-forest type	Uncertainty estimate (confidence interval at 95%)								
Cropland (C)	75.00%								
Grassland (P)	75.00%								
Other lands (A O U)	-								
Any comment:	-								

Parameter:	BGB _{after,i}
Description:	Belowground biomass of non-forest type <i>i</i> after conversion
Data unit:	tons of dry matter per ha
Source of data or description of the method for developing the data including the	<p>For cropland: The values and assumptions of 2006 IPCC GL, Volume 4, Chapter 5 are used because, unfortunately, there aren't country-specific data. Tier 2 may modify the assumption that carbon stocks immediately following conversion are zero. In this case, it is assumed that conversion leads to annual croplands and in the case the carbon stock in biomass after one year for annual crops provided in TABLE 5.9 is used.</p> <p>For grassland: The values and assumptions of 2006 IPCC GL, Volume 4, Chapter 6, TABLE 6.1 and TABLE 6.4 are used because, unfortunately, there aren't country-specific data. As the climate in most of Mozambique is tropical dry to subtropical dry, the value for semi-arid grassland in</p>

spatial level of the data (local, regional, national, international):	tropical dry climate zone is used, therefore a root-shoot ratio of 2.8 (TABLE 6.1) is applied to the value of peak above-ground biomass, 2.3 tonnes of dry biomass per hectare (TABLE 6.4), generating the expected values 6.4 tonnes of dry biomass per hectare. For other lands: No default values exist for these conversions. Spatial level: International								
Value applied:	<table border="1"> <tr> <td>Cropland (C)</td> <td>0.00</td> </tr> <tr> <td>Grassland (P)</td> <td>6.44</td> </tr> <tr> <td>Other lands (A O U)</td> <td>0.00</td> </tr> </table> <p>The values above are recorded in the ranges "B47:E51", "B53:B57" and "B59:B63" of the "BIOMASS" worksheet tab in the "Outside ZILMP Emissions Calculations MR (2021)" and "Outside ZILMP Emissions Calculations MR (2022)" workbooks. These values are then applied in the range "F9:F20" of the "EMISSION MONITORING PERIOD(ERL)" worksheet tab in the "Outside ZILMP Emissions Calculations MR (2021)" and "Outside ZILMP Emissions Calculations MR (2022)" workbook for estimating emissions.</p>	Cropland (C)	0.00	Grassland (P)	6.44	Other lands (A O U)	0.00		
Cropland (C)	0.00								
Grassland (P)	6.44								
Other lands (A O U)	0.00								
QA/QC procedures applied	The adequacy in the use of these default values was confirmed with the experts in GHG Inventory in DINAB.								
Uncertainty associated with this parameter:	<table border="1"> <thead> <tr> <th>Non-forest type</th> <th>Uncertainty estimate (confidence interval at 95%)</th> </tr> </thead> <tbody> <tr> <td>Cropland (C)</td> <td>-</td> </tr> <tr> <td>Grassland (P)</td> <td>121.04%</td> </tr> <tr> <td>Other lands (A O U)</td> <td>-</td> </tr> </tbody> </table>	Non-forest type	Uncertainty estimate (confidence interval at 95%)	Cropland (C)	-	Grassland (P)	121.04%	Other lands (A O U)	-
Non-forest type	Uncertainty estimate (confidence interval at 95%)								
Cropland (C)	-								
Grassland (P)	121.04%								
Other lands (A O U)	-								
Any comment:	-								

1.1.7 Monitored Data and Parameters

Parameter:	$A(j,i)_{MP}$																														
Description:	Area converted from forest type j to non-forest type i during the Monitoring Period.																														
Data unit:	hectare per year.																														
Value monitored during this Monitoring / Reporting Period:	<table border="1"> <thead> <tr> <th>Type of change</th> <th>2021</th> <th>2022</th> </tr> </thead> <tbody> <tr> <td>Semi-deciduous forest to cropland</td> <td>26,873.1</td> <td>23,034.76</td> </tr> <tr> <td>Semi-deciduous forest to grassland</td> <td>1,119.9</td> <td>59.60</td> </tr> <tr> <td>Semi-deciduous forest to other lands</td> <td>0.00</td> <td>811.45</td> </tr> <tr> <td>Evergreen forest to cropland</td> <td>1,557.9</td> <td>1,243.81</td> </tr> <tr> <td>Evergreen forest to grassland</td> <td>296.2</td> <td>532.51</td> </tr> <tr> <td>Evergreen forest to other lands</td> <td>0.00</td> <td></td> </tr> <tr> <td>Mangrove forest to cropland</td> <td>0.00</td> <td></td> </tr> <tr> <td>Mangrove forest to grassland</td> <td>0.00</td> <td></td> </tr> <tr> <td>Mangrove forest to other lands</td> <td>0.00</td> <td></td> </tr> </tbody> </table>	Type of change	2021	2022	Semi-deciduous forest to cropland	26,873.1	23,034.76	Semi-deciduous forest to grassland	1,119.9	59.60	Semi-deciduous forest to other lands	0.00	811.45	Evergreen forest to cropland	1,557.9	1,243.81	Evergreen forest to grassland	296.2	532.51	Evergreen forest to other lands	0.00		Mangrove forest to cropland	0.00		Mangrove forest to grassland	0.00		Mangrove forest to other lands	0.00	
Type of change	2021	2022																													
Semi-deciduous forest to cropland	26,873.1	23,034.76																													
Semi-deciduous forest to grassland	1,119.9	59.60																													
Semi-deciduous forest to other lands	0.00	811.45																													
Evergreen forest to cropland	1,557.9	1,243.81																													
Evergreen forest to grassland	296.2	532.51																													
Evergreen forest to other lands	0.00																														
Mangrove forest to cropland	0.00																														
Mangrove forest to grassland	0.00																														
Mangrove forest to other lands	0.00																														

<p>Source of data and description of measurement/calculation methods and procedures applied:</p>	<p>(i) Source</p> <p>Activity data used for the monitoring period are obtained from a combination of an annual wall-to-wall deforestation map with sampling to generate deforested area estimates through a stratified estimator.</p> <p>(ii) Variable of interest</p> <p>The variable of interest are all the transitions specified above. It is important to note that the variables of interest are not aligned to the strata as this is not required. Strata is linked to the likelihood of presence of deforestation events, whereas the variable of interest is linked to the possible transitions of deforestation per forest type and post-deforestation type.</p> <p>(iii) Annual deforestation map</p> <p>The workflow used to produce annual deforestation map for the area outside the scope of ZILMP follows the steps below:</p> <ol style="list-style-type: none"> 1. Produce two Sentinel-2 satellite imagery composites for the monitoring area, containing all images of wet season (i.e. January - May). The first composite for the 2021 deforestation map comprises the period between January to May 2021 denoted as the reference period and the second composite comprises the period from January to May 2022, referred as actual period. For the 2022 deforestation map the selected images comprises the period between January to May 2022 and January to May 2023. The reason behind the selection of January - May as a reference and actual period of monitoring resides on the fact that it is the wet season, where the NDVI stability is very high in relation to the dry season, which starts in June to October, when most trees lose their foliage and makes it difficult the analysis of deforestation. 2. Generate image features from reference period and actual period from the composites generated in previous step, to identify changes in forest cover. The image features have different vegetation indexes, namely, NDVI, EVI, SAVI, NBR, NDWI with respective sub-products such as NDVI 90th percentile, Normalized NDVI, and variation on NDVI. 3. Generate training data on classes of deforestation, stable forest and stable non-forest by visual interpretation of composites from the reference and actual periods, and NDVI change detection image. The NDVI change detection image is a result of the difference of NDVI from the composites of reference and actual periods. The calculated NDVI change detection image helps the interpreter to locate where the changes of forest cover are occurring. 4. Produce a categorical deforestation map from training data and image features through a process of classification using Random Forest classifier. The Categorical deforestation map includes non-forest stable and stable forest classes. Because errors of omission of deforestation have a very large impact on the final estimates, it is important to reduce these errors as much as possible. 5. To improve the efficacy of the sampling the deforestation class on the map is reclassified as: <ol style="list-style-type: none"> a) High probability deforestation (cluster of more than 10 pixels of deforestation, corresponding to at least 40% of one hectare); b) Low probability of deforestation (cluster of less than 10 pixels and greater than 2 pixels, corresponding at least 12% to 40% of one hectare) and; c) Non-forest (cluster of less than 3 pixels, corresponding to less than 12% of a hectare). 6. To reduce the risk of omission errors, a Buffer of 60 meters is added around the high probability of deforestation class. The result is a deforestation map with five classes: High
---	--

probability of deforestation; buffer; low probability of deforestation; stable forest and stable non-forest.

(iv) Sampling design

Sampling method

Monitoring of activity data for annual reporting is conducted using a stratified estimator, where deforestation map (which includes classes of forest and non-forest) is used for stratification and reference-sampling units are used for estimate activity data and associated confidence intervals.

Sample size determination

The sample size n was determined from the equation:

$$n = \frac{(\sum W_i S_i)^2}{[S(\hat{O})]^2 + \frac{1}{N} \sum W_i S_i} \approx \left(\frac{\sum W_i S_i}{S(\hat{O})} \right)^2 \quad \text{Equation 24}$$

Where:

- N Number of units in the ROI
- $S(\hat{O})$ Standard error of the estimated overall accuracy that we would like to achieve
- W_i Mapped proportion of area of class i ; and
- S_i Standard deviation of stratum i .

The standard deviation of stratum i is given by the formula:

$$S_i = \sqrt{U_i(1 - U_i)} \quad \text{Equation 25}$$

Where:

- U_i Proportion of area of deforestation in stratum i .

In order to obtain approximate values of proportion of deforestation in each stratum (U_i), a pilot sampling is conducted. This pilot consists of 100 sample units per stratum.

Sample units per stratum

After the pilot sampling, sample units may need to be added to each stratum, in order to achieve 20% relative margin error at 95% confidence level. It was decided to use the Optimum (Neyman) allocation for each change stratum, where the stratum standard deviation $S_i = \sqrt{U_i \cdot (1 - U_i)}$ increases the number of plots (ensuring larger numbers of plots in rare classes or strata) and sampling unit costs are constant:

$$n_i = n \frac{w_i \cdot S_i}{\sum_{i=1}^I w_i \cdot S_i} \quad \text{Equation 26}$$

And for each stable stratum, the proportional allocation is applied if deforestation omission errors are completely absent from these strata. In stratified sampling the sample size for proportional allocation is given by:

$$n_i = n \cdot w_i$$

Equation 27

The number of reference points is presented in Annex-Table 8.

Annex-Table 8: Number of reference sampling units per map stratum for monitoring period (2021 and 2022).

Stratum	Number of sample units 2021	Number of sample units 2022
High probability of deforestation	100	100
60 m Buffer	277	300
Low probability of deforestation	100	100
Forest	300	300
Non-forest	300	300
Total	1,077	1,100

(v) Response design

Sampling unit and spatial support

The sampling unit is a 20 m pixel of the stratification map that was produced. The spatial support used is a 100m x 100m plot (1ha). Each Spatial sampling unit contains an internal grid of 5 x 5 points (20m x 20m grid) to aid in the labelling attribution (Annex-Figure 6).

Source of reference data

Each sampling unit was evaluated using Collect Earth (<http://www.openforis.org/>). This tool enables access to high-resolution images in Google Earth, Bing Maps and Planet Labs, as well as a medium resolution image repository available through Google Earth Engine Explorer and Code Editor (Landsat and Sentinel-2). The tool enables to display digital forms designed to collect the Land-Use Land Cover Change and Forestry (LULCCF) information on the sampling points (Annex-Figure 7). The Earth Engine Code Editor facilitates the interpretation of the vegetation type and the determination of LULC changes, by displaying the historical MOD13Q1 (NDVI 16-day Global Modis 250 m) graphic as well as monthly mosaics of Sentinel-2 images. The main source of data to identify changes in land cover, is Sentinel-2 15 days TOA reflectance composites. However, Planet data is also used in cases of doubt or excessive cloud cover with Sentinel-2.

Reference labelling protocol

The activity data was generated considering the national land use and land cover classification system, which reflects the six broad IPCC Land Use categories.

A set of hierarchical rules were established and used to determine the LULCCF category based on a certain percentage and taking into account the national forest definition as well (Annex-Figure 8). A single land use class is easier to classify, but it becomes challenging when there is a combination of two or more land use classes within the area of interest. Thus, this is where the hierarchical rules are important to determine the land use. Any sampling unit that has 30% of tree canopy cover is considered a forest, according to the national forest definition, even if it has more than 20% of settlements, crops or other land use, the forest is priority.

In the case the sampling unit was classified as forest land and different forest types were present in the sampling unit, a majority rule was used in this case, i.e. the largest forest class is chosen (please click [here](#) for more details).

(vi) Analysis

Applying the methodology described in Olofsson *et al.* (2014)³⁷ and the GFOI MGD the estimations of the areas corresponding to land-use and land-cover change categories, more specifically the activity data for deforestation, in the framework of this stratified random sampling approach (based on the visual assessment of the 1 ha plots) was based on assessments of area proportions. A sample error matrix is constructed where the map classes ($h=1, 2, \dots, q$) are represented by rows and the reference data ($k=1, 2, \dots, q$) by columns as shown in Annex-Table 9.

Annex-Table 9: Error matrix of area proportions.

Map data	Reference data					Total	User's accuracy (\hat{U}_i)
	Deforestation			Stable forest	Stable non-forest		
	High probability of deforestation	40 m Buffer	Low probability of deforestation				
High probability of deforestation	\hat{p}_{11}	\hat{p}_{12}	\hat{p}_{13}	\hat{p}_{14}	\hat{p}_{15}	$\hat{p}_{.1}$	$\hat{p}_{11}/\hat{p}_{.1}$
40 m Buffer	\hat{p}_{21}	\hat{p}_{22}	\hat{p}_{23}	\hat{p}_{24}	\hat{p}_{25}	$\hat{p}_{.2}$	$\hat{p}_{22}/\hat{p}_{.2}$
Low probability of deforestation	\hat{p}_{31}	\hat{p}_{32}	\hat{p}_{33}	\hat{p}_{34}	\hat{p}_{35}	$\hat{p}_{.3}$	$\hat{p}_{33}/\hat{p}_{.3}$
Stable forest	\hat{p}_{41}	\hat{p}_{42}	\hat{p}_{43}	\hat{p}_{44}	\hat{p}_{45}	$\hat{p}_{.4}$	$\hat{p}_{44}/\hat{p}_{.4}$
Stable non-forest	\hat{p}_{51}	\hat{p}_{52}	\hat{p}_{53}	\hat{p}_{54}	\hat{p}_{55}	$\hat{p}_{.5}$	$\hat{p}_{55}/\hat{p}_{.5}$
Total	$\hat{p}_{.1}$	$\hat{p}_{.2}$	$\hat{p}_{.3}$	$\hat{p}_{.4}$	$\hat{p}_{.5}$	1	
Producer's accuracy (P_i)	$\hat{p}_{11}/\hat{p}_{.1}$	$\hat{p}_{22}/\hat{p}_{.2}$	$\hat{p}_{33}/\hat{p}_{.3}$	$\hat{p}_{44}/\hat{p}_{.4}$	$\hat{p}_{55}/\hat{p}_{.5}$		Overall accuracy (\hat{O}) = $\hat{p}_{11} + \hat{p}_{22} + \hat{p}_{33} + \hat{p}_{44} + \hat{p}_{55}$

The mean estimator for the area of each class can be directly obtained from the error matrix. Unbiased stratified estimators are provided using reference class area proportions ($\hat{p}_{.k}$):

$$\hat{p}_{.k} = \sum_{h=1}^H w_h \cdot \frac{n_{hk}}{n_h} = \sum_{h=1}^H \hat{p}_{hk} \quad \text{Equation 28}$$

Where:

$\hat{p}_{.k}$ Area proportions of reference data class k . These proportions of reference data for deforestation classes as a whole are collapsed in three possible types of conversions/transitions from forest type j to non-forest type i , namely:

- Broadleaved (Semi-) deciduous to Non-forest type i ;
- Broadleaved (Semi-) evergreen to Non-forest type i ; and
- Mangrove to Non-forest type i .

³⁷ Olofsson, P., Foody, G.M., Herold, M., Stehman, S.V., Woodcock, C.E., & Wulder, M.A. 2014. Good practices for estimating area and assessing accuracy of land change. *Remote Sensing of Environment*. 148:42-57.

	<p>Five types of non-forest land are considered:</p> <ul style="list-style-type: none"> ● Cropland (C); ● Grassland (P); ● Wetland (A); ● Settlement (U); and ● Other lands (O). <p>w_h Proportion of area mapped as class h; n_{hk} Sample count at cell (h,k); n_h Sum of sample counts across row h; and \hat{p}_{hk} Proportion of area in cell (h,k).</p> <p>Once the estimated reference class area proportions (\hat{p}_k) are obtained, the mean total area per class is calculated by multiplying them with the total reporting area a:</p> $\hat{A}_j = \hat{p}_k \cdot a \quad \text{Equation 29}$ <p>The estimated standard error for the reference class area proportions was given by:</p> $S(\hat{p}_j) = \sqrt{\sum_{h=1}^H w_h^2 \cdot \frac{\hat{p}_{hj} \cdot (1 - \hat{p}_{hj})}{n_h - 1}} \quad \text{Equation 30}$ <p>where the term inside the root is the variance of the reference class area proportion. Translated to actual area,</p> $S(\hat{A}_j) = S(\hat{p}_j) \cdot a \quad \text{Equation 31}$ <p>Given the confidence level (i.e., 95%, expressed as a fraction, that is, 0.95), the significance level is $\alpha = 1 - \text{confidence level}$, one must use Student's t given α and the degrees of freedom, $df = n_h - 1$. For large samples, $df \rightarrow 1.96$. Then the confidence interval of the estimated area per class was given by:</p> $CI(\hat{A}_j) = t_{\alpha, df} \cdot S(\hat{p}_j) \quad \text{Equation 32}$ <p>The uncertainty, usually represented as a percentage, then becomes:</p> $U(\hat{A}_j) = \frac{CI(\hat{A}_j)}{\hat{A}_j} \cdot 100 \quad \text{Equation 33}$
<p>QA/QC procedures applied:</p>	<p>The QA/QC procedures consisted on the following:</p> <ul style="list-style-type: none"> ● SOPs were developed as described in Section 2.1 - Satellite and land monitoring system and training; and ● Interpretation is done by highly qualified professionals which are specialized in land cover interpretation with satellite imagery. They were trained and a robust control system is in place to ensure that they are correctly calibrated throughout the data collection process. ● All reference data interpreted as deforestation or forest degradation, and an additional 20% of the remaining reference data were evaluated. The quality control is carried out by two independent supervisors, who after the independent evaluation compare the two evaluations and consensually compile a single comment for each sample. The parameters to be taken into account in the evaluation for identifying errors are: a) the

	<p>percentage of coverage for each element within the plot; b) the current land cover/land use class (levels 1 and 2); c) the land cover/land use change class; d) the former land cover/land use class (levels 1 and 2); and e) the date of occurrence of land cover/land use change, or evidence date of remaining land cover/land use. If there are gross errors related to the parameters b), c) and d) in at least 20% of samples from the 20% mentioned initially, the respective interpreter should review all samples from the batch, otherwise the interpreter reviews only the samples evaluated by the supervisors, that present gross errors. On the other hand, in relation to all samples interpreted as deforestation, the interpreter reviews only the samples that present gross errors according to the evaluation from the supervisors. The process is cyclical until the interpreter achieves values less than 20% of gross errors in the batch.</p> <ul style="list-style-type: none"> • The sampling design and estimation was reviewed by an international renowned expert (Steve Stehman), a statistics professor of State University of New York. 																		
<p>Uncertainty for this parameter:</p>	<table border="1" data-bbox="491 689 1422 1003"> <thead> <tr> <th data-bbox="491 689 718 797">Category change</th> <th data-bbox="718 689 1067 797">Uncertainty estimate (Confidence interval at 95%) 2021</th> <th data-bbox="1067 689 1422 797">Uncertainty estimate (Confidence interval at 95%) 2022</th> </tr> </thead> <tbody> <tr> <td data-bbox="491 797 718 840">FSD>C</td> <td data-bbox="718 797 1067 840">20.49%</td> <td data-bbox="1067 797 1422 840">± 21.42%</td> </tr> <tr> <td data-bbox="491 840 718 882">FSD>P</td> <td data-bbox="718 840 1067 882">82.13%</td> <td data-bbox="1067 840 1422 882">± 196.21%</td> </tr> <tr> <td data-bbox="491 882 718 925">FSD>(A O U)</td> <td data-bbox="718 882 1067 925">-</td> <td data-bbox="1067 882 1422 925">± 140.00%</td> </tr> <tr> <td data-bbox="491 925 718 967">FSSV>C</td> <td data-bbox="718 925 1067 967">92.42%</td> <td data-bbox="1067 925 1422 967">± 119.69%</td> </tr> <tr> <td data-bbox="491 967 718 1003">FSSV>P</td> <td data-bbox="718 967 1067 1003">159.06%</td> <td data-bbox="1067 967 1422 1003">± 196.21%</td> </tr> </tbody> </table>	Category change	Uncertainty estimate (Confidence interval at 95%) 2021	Uncertainty estimate (Confidence interval at 95%) 2022	FSD>C	20.49%	± 21.42%	FSD>P	82.13%	± 196.21%	FSD>(A O U)	-	± 140.00%	FSSV>C	92.42%	± 119.69%	FSSV>P	159.06%	± 196.21%
Category change	Uncertainty estimate (Confidence interval at 95%) 2021	Uncertainty estimate (Confidence interval at 95%) 2022																	
FSD>C	20.49%	± 21.42%																	
FSD>P	82.13%	± 196.21%																	
FSD>(A O U)	-	± 140.00%																	
FSSV>C	92.42%	± 119.69%																	
FSSV>P	159.06%	± 196.21%																	
<p>Any comment:</p>	<p>-</p>																		

5.4 QUANTIFICATION OF EMISSION REDUCTIONS

5.4.1 Reference level for the Monitoring / Reporting Period covered in this report

Year of Monitoring/Reporting period t	Average annual historical emissions from deforestation over the Reference Period (tCO _{2-e} /yr)	If applicable, average annual historical emissions from forest degradation over the Reference Period (tCO _{2-e} /yr)	If applicable, average annual historical removals by sinks over the Reference Period (tCO _{2-e} /yr)	Adjustment, if applicable (tCO _{2-e} /yr)	Reference level (tCO _{2-e} /yr)
2021	7,932,644.34	-	-	-	7,932,644.34
2022	7,932,644.34				7,932,644.34
Total	15,865,288.68	-	-	-	15,865,288.68

5.4.2 Estimation of emissions by sources and removals by sinks included

The following table shows the emissions results obtained per category changes from a forest type to a non-forest type during the Monitoring Period. The emissions are generated relating the data and parameters described in [Subsection 5.3.2](#) and summarized in the Annex-Table 10, by applying [Equation 14](#).

Annex-Table 10: Calculation of the emissions during the Monitoring Period (2021 and 2022).

Category changes	AGB _{before,j} (tdm/ha)	BGB _{before,j} (tdm/ha)	AGB _{after,i} (tdm/ha)	BGB _{after,i} (tdm/ha)	2021		2022	
					A _{(j,i)MP} (ha)	Emissions (tCO _{2e})	A _{(j,i)MP} (ha)	Emissions (tCO _{2e})
Semi-deciduous forest to cropland	144.69	49.95	10	0	26,873.1	8,550,698	23,034.76	7,329,385
Semi-deciduous forest to grassland	144.69	49.95	2.3	6.4	1,119.9	358,762	59.60	19,093
Semi-deciduous forest to other lands	144.69	49.95	0	0	0	0	811.45	272,176
Evergreen forest to cropland	123.13	42.06	10	0	1,557.9	416,643	1,243.81	332,644
Evergreen forest to grassland	123.13	42.06	2.3	6.4	296,2	79,870	532.51	143,570

Evergreen forest to other lands	123.13	42.06	0	0	0	0	0	0
Mangrove to cropland	269.01	85.43	10	0	0	0	0	0
Mangrove to grassland	269.01	85.43	2.3	6.4	0	0	0	0
Mangrove to other lands	269.01	85.43	0	0	0	0	0	0
Total						9,405,973		8,096,868

Year of Monitoring/Reporting Period	Emissions from deforestation (tCO ₂ -e/yr)	If applicable, emissions from forest degradation (tCO ₂ -e/yr)*	If applicable, removals by sinks (tCO ₂ -e/yr)	Net emissions and removals (tCO ₂ -e/yr)
2021	9,405,973	-	-	9,405,973
2022	8,096,868			8,096,868
Total	17,502,841	-	-	17,502,841

5.4.3 Calculation of emission reductions

Total Reference Level emissions during the Reporting Period (tCO₂-e)	15,865,288.68
Net emissions and removals under the ER Program during the Reporting Period (tCO₂-e)	17,502,841.14
Emission Reductions during the Reporting Period (tCO₂-e)	-1,637,552.46

5.5 UNCERTAINTY OF THE ESTIMATE OF EMISSION REDUCTIONS

Uncertainties were propagated using the Approach 1 of the 2019 Refinement to the 2006 IPCC GL, i.e. propagation of uncertainties. The following equations were used for addition or multiplication.

For addition or subtraction:

$$U_{total} = \frac{\sqrt{(U_1 \cdot x_1)^2 + \dots + (U_i \cdot x_i)^2 + \dots + (U_n \cdot x_n)^2}}{|x_1 + \dots + x_i + \dots + x_n|} \quad \text{Equation 34}$$

Where:

- U_i The percentage uncertainties associated with each of the quantities;
- X_i Quantities to be combined; x_i may be a positive or a negative number; and
- U_{total} The percentage uncertainty in the sum of the quantities (half the 95 percent confidence interval divided by the total (i.e., mean) and expressed as a percentage)

For multiplication:

$$U_{total} = \sqrt{U_1^2 + \dots + U_i^2 + \dots + U_n^2} \quad \text{Equation 35}$$

Where:

- X_i The percentage uncertainties associated with each of the quantities; and
- U_{total} The percentage uncertainty in the product of the quantities (half the 95 percent confidence interval divided by the total and expressed as a percentage)

	2021	2022
Uncertainty of Reference Level emissions during the Monitoring Period (%)	19.34	19.34
Uncertainty of net emissions and removals under the ER Program during the Monitoring Period (%)	24.57	25.71
Uncertainty of Emission Reductions during the Reporting Period (%)	-188.3	-1575

ANNEX 7: DEGRADATION IN THE SCOPE OF ZAMBÉZIA INTEGRATED LANDSCAPE MANAGEMENT PROGRAM (ZILMP)

Since 2020, Mozambique has been developing a methodology for estimating forest degradation. This methodology was developed taking into account the process used to estimate deforestation to allow for consistency over time. This chapter (Annex 7) was prepared only to demonstrate the country's process and progress regarding estimates of emissions from forest degradation for 2021.

1.1. Overview of the approach to estimate degradation

FCPF methodological guidance provides few limits to the approach for monitoring degradation. Degradation is required to be accounted for where emissions are estimated to be significant (>10% of total forest-related emissions). Where included degradation should be estimated using 'best available data (including proxy activities or data)'. While 'direct' methods are preferred, 'activity data may be derived using indirect methods such as survey data, proxies derived from landscape ecology, or statistical data on timber harvesting and regrowth if no alternative methods are available'.

Mozambique has been testing a range of approaches for expanding its existing monitoring procedures to include estimates of degradation. Five broad options for estimation emissions from degradation were identified:

1. Remote sensing based estimates (e.g. changes in land cover maps)
2. Forestry statistics (e.g. timber production statistics)
3. Forest inventory plot data (e.g. cyclical NFI plots)
4. Model-based estimates (e.g. modelling fuelwood demand)
5. Sample-based estimation

Of these, only sample based area estimation was considered feasible for ZILMP. Remote sensing based estimates [1] were not yet considered sufficiently reliable for degradation, a problem that is particularly great in the dry tropics where satellite monitoring of forest cover can be problematic. Forestry statistics [2] and forest inventory plot data [3] are not currently systematically collected for the ZILMP region, and would require an unrealistically large data collection effort to implement. Model or proxy based estimates [4] were considered challenging for ZILMP on account of difficulty in implementation and given limited evidence that they could work.

Sample based area estimation [5] fits well within existing MRV workflows, and has the potential to be applied to existing monitoring datasets. Methods are also well-developed for emissions estimation using sample-based data, and development of a transparent process for estimating emissions uncertainty was considered possible.

Methodological development work is described in more detail in the report: 'FNDS degradation: final report' (March 2023). This technical annex describes the finalized approach for measuring degradation, plans for future improvements of estimates, and provides provisional estimates of degradation-related emissions for ZILMP in 2021.

1.2. Methods descriptions

In summary, degradation is defined as the loss of canopy cover that transforms a 'closed canopy' forest (greater than or equal to 65% canopy cover) to an 'open canopy' forest (less than 65% canopy cover, but still at least 30%). Emissions are estimated from the difference of carbon density between a 'closed' and 'open' canopy woodland. Activity data are derived from systematic (reference period) or stratified random (monitoring period) sampling, using high-resolution imagery to record degradation where canopy cover of closed canopy forest is reduced to that of open canopy forest. Uncertainty estimates are derived using a Monte Carlo method adapted from the existing deforestation emissions methods.

1.2.1. Definition of degradation

Degradation is defined as the loss of canopy cover such that a formerly ‘closed-canopy’ forest ($\geq 65\%$ canopy cover) is modified to ‘open-canopy’ forest ($\geq 30\%$ to $< 65\%$ canopy cover) as the result of human action. Thresholds for closed- and open-canopy forest are derived from Mozambique’s existing forest cover definitions. The degradation is recorded in the case that a dense forest transitions to open forest. A separate carbon density is applied to each of open and closed forests, and transitions between these two classes are associated with emissions or removals equivalent to the difference between the carbon stocks of a dense vs sparse forest. This method should prevent double-counting of emissions between deforestation and degradation, as deforestation that follows from degradation will be associated with lowered emissions.

1.2.2. Emission factor

ZILMP uses a single emission factor for each forest stratum in the region, consisting of Semi-deciduous forest (FSD), Evergreen forest (FSSV), and Mangrove forest (FF) strata. Above-ground biomass is estimated using data from Mozambique’s National Forest Inventory in combination with a series of allometric models. Below-ground biomass is estimated using root:shoot ratios, and emissions are assumed to be instantaneous following deforestation.

The updated method uses a similar approach, but rather than a single biomass density per forest stratum, separate biomass densities are estimated for closed-canopy and open-canopy forests (Annex 7 - Table 1). No changes are made to other elements of emission factor estimation, including allometric models, root:shoot ratios, or uncertainty estimation methods. Forests in the FF stratum are not currently given separate emission factors for closed and open canopy areas, as no suitable data currently exist for this assessment, and examples of degradation in this stratum have not yet been observed.

The Emission factors were previously estimated in a complex [Excel spreadsheet](#), which outputs estimates of above- and below- ground biomass in each relevant forest type, along with an estimate of uncertainty (t-distribution). These methods provide functionality to re-estimate emission factors given permuted root:shoot ratios.

Annex 7 - Table 1: Biomass densities used for emission factors in the existing ZILMP emissions estimation procedures (level 1) and updated biomass densities for open-canopy and closed-canopy forests (level 2)

Stratum	Biomass (tonnes/ha)		Uncertainty (%)		Degrees of freedom	
	Above-ground	Below-ground	Above-ground	Below-ground	Above-ground	Below-ground
Level 1						
Semi-deciduous forest	144.69	49.95	13.3	9.97	28.7	26.0
Evergreen forest	123.13	45.06	8.71	7.83	5.2	4.01
Mangrove	269.01	85.43	10	85.4	-	-
Level 2						
Semi-deciduous open forest	128.47	43.79	14.4	43.8	17.8	11.8

Semi-deciduous closed forest	174.11	61.11	10.9	61.1	15.1	15.5
Evergreen open forest	114.65	41.02	6.59	41.0	3.52	3.43
Evergreen closed forest	174.11	47.23	19.8	47.2	3.37	3.14
Non-forest						
Cultivation	10	0	37.5	0	-	-
Grassland	2.3	6.44	37.5	60.5	-	-
(A O U)	0	0	0	0	-	-

1.2.3. Activity data

Activity data derive from the same sample-based method that is used to quantify rates of deforestation (see [section 3](#) of this report). Collect Earth is used to label land use and land use changes using high-resolution satellite imagery over a series of sample points. Sample points are based either on a systematic sample (reference period) or a stratified sample (monitoring period). Stratification is based on an existing deforestation mapping tool and buffer regions designed to minimise errors of omission.

Two methods are used to identify degradation using outputs from Collect Earth:

Pre-2021 Historically FNDS-MRV have recorded all canopy cover changes in each Collect Earth sample unit, in addition to monitoring of degradation. These records are somewhat limited, only recording final canopy cover (to the nearest 10%), and canopy cover losses and gains in 10% bins. From this information points of degradation can be flagged where final canopy cover falls between 30% - 65% (open-canopy), and canopy cover losses were sufficiently large to mean that initial forest cover must have been $\geq 65\%$ (closed-canopy). This approach is somewhat limited, as little quality assurance was conducted on this aspect of the data, and historical satellite imagery is sometimes insufficient to identify small-scale forest changes.

Post-2021 FNDS-MRV updated its [Collect Earth survey](#) to require that operators record the initial canopy cover of each point in addition to the final canopy cover. Alongside, operators are asked to flag degradation where canopy cover losses are such that a sample unit meets the definition of degradation. This approach can be considered considerably more robust, and quality assurance procedures have been updated to ensure that data quality is high.

In 2021 a further change was implemented for mapping procedures to improve monitoring of degradation. Errors of omission were observed of degradation points from the map change classes, resulting in a high degree of uncertainty in emissions estimates. Larger buffer regions were included to capture minor changes on the periphery of deforestation, and smaller areas of forest change (previously excluded) were included in change classes. For consistency, these same changes were rolled out for deforestation monitoring from 2021 onwards. It is not anticipated this will impact uncertainty of deforestation estimates. The Annex 7 - Table 2 show the modifications to standard operating procedures for map production to improve monitoring of canopy forest change for degradation

Annex 7 - Table 2. Modifications to standard operating procedures for map production to improve monitoring of canopy forest change for degradation

Class	Colour	Label	Modifications
1	#CE0000	High probability deforestation	-
2	#FFA500	Buffer deforestation	Buffer size increased from 40 m to 60 m
3	#FFFF00	Low probability deforestation	Minimum of 3 connected pixels in place of 6

4	#367D49	Stable forest	-
5	#FFBCC	Stable non-forest	-

The Sample data collected across the five map strata are summarized in Annex 7 - Table 3. Notably, no errors of omission were detected in either deforestation or degradation classes in 2021 (map strata 4 and 5). Without this property, attaining reasonably constrained estimates on emissions reductions associated with degradation is not possible. Also notable is the absence of enhancement points detected in the monitoring period. This is likely down to monitoring on a year-to-year basis not being sufficiently long to reliably identify forest regrowth.

Annex 7 - Table 3: Sample counts from systematic sampling (reference period) and stratified random sampling (monitoring period).

Stratum	Number of samples	Deforestation	Degradation	Enhancement	Area (ha)
Reference period					
-	3,318	117	46	22	5,310,265
Monitoring period 2021					
1 - high probability deforestation	100	69	4	0	25,013
2 - buffer deforestation	309	21	9	0	226,309
3 - low probability deforestation	100	32	5	0	9,891
4 - stable forest	300	0	0	0	2,225,307
5 - stable non-forest	300	0	0	0	2,823,745

Activity data resulting from stratified area estimation (2021) are shown in Annex 7 - Table 4. Outputs indicate that the area of forest impacted by degradation is around a quarter to one third of that subject to deforestation, and that it impacts FSD and FSSV forest strata in a proportion similar to deforestation. Uncertainties associated with deforestation and forest degradation are of comparable magnitude. The area of deforestation is larger in 2021, while degradation appears roughly stable.

Annex 7 - Table 4: Areas of deforestation, degradation and enhancement observed in the ZILMP reference period and the monitoring period of 2021.

	Reference period		Monitoring period 2021	
	Area (ha/yr)	SE (ha/yr)	Area (ha/yr)	SE (ha/yr)
Deforestation total	17,023	-	35,805	-
Semi-deciduous open forest > other land (A O U)	145	145	-	-
Semi-deciduous open forest >	7,857	1,060	27,097	3,365

cultivation				
Semi-deciduous open forest > grassland	1,309	436	250	250
Semi-deciduous closed forest > cultivation	3,928	753	6,480	1,277
Semi-deciduous closed forest > grassland	436	252	-	-
Evergreen open forest > cultivation	1,600	482	1,681	860
Evergreen open forest > grassland	145	145	-	-
Evergreen closed forest > cultivation	1,600	482	198	139
Mangrove> cultivation	-	-	99	99
Degradation total	6,256	-	8,087	-
Semi-deciduous closed forest > Semi-deciduous open forest	5,529	892	8,087	2,234
Evergreen closed forest > Evergreen open forest	727	325	-	-
Enhancement total	3,201	-	0	-
Semi-deciduous open forest > Semi-deciduous closed forest	2,764	632	-	-
Evergreen open forest > Evergreen closed forest	436	252	-	-

1.2.4. Emission estimation

Methods for emissions estimation remain unchanged from existing documentation for ZILMP. The only difference is that now more land cover transitions are possible, including to and from open- and closed-canopy forest types. New forest transition types are summarised in Annex 7 - Table 5.

Summary of forest change classes following from the inclusion of degradation.

Annex 7 - Table 5: Summary of forest change classes following from the inclusion of degradation.

From ↓	To →	Closed-canopy forest	Open-canopy forest	Non-forest
Closed-canopy forest		-	Degradation	Deforestation
Open-canopy forest		Enhancement	-	Deforestation
Non-forest		Afforestation	Afforestation	-

1.2.5. Uncertainty estimation

Uncertainty estimation follows the same approach as estimates for deforestation. Uncertainties in emission factors are estimated using National Forest Inventory, but with the addition of further forest classes to account for closed and open canopy variants. Uncertainties in activity data are derived from non-parametric bootstrapping (resampling with replacement), again with new classifications for the transitions between open- and closed-canopy forests. Uncertainty in emissions reductions estimates are estimated using Monte Carlo analysis, using at least 10,000 parameter sets. Methods for uncertainty estimation are described in more detail in [section 5](#) of this report.

1.2.6. Results

Emission

Emissions increased in 2021, with a marked increase in deforestation emissions (please see the section [4.3](#) of this report). Emissions from degradation are an order of magnitude smaller than those from deforestation (~10%), and show a small increase in 2021 relative to the reference period (Annex 7 - Table 6).

Annex 7 - Table 6: Nominal emissions reductions for deforestation and degradation. *Note: negative emissions reductions indicate increased emissions in the monitoring period.

	Emissions (tCO ₂ /yr)		
	Reference period	Monitoring period 2021	Reductions*
Degradation total	671,556	877,431	-205,875
Semi-deciduous closed forest > Semi-deciduous open forest	599,899	877,431	-277,532
Evergreen close forest > Evergreen open forest	71,657	-	71,657

Reference level for the Monitoring / Reporting Period covered in this report

Year of Monitoring/Reporting period <i>t</i>	Average annual historical emissions from deforestation over the Reference Period (tCO _{2-e} /yr)	If applicable, average annual historical emissions from forest degradation over the Reference Period (tCO _{2-e} /yr)	If applicable, average annual historical removals by sinks over the Reference Period (tCO _{2-e} /yr)	Adjustment, if applicable (tCO _{2-e} /yr)	Reference level (tCO _{2-e} /yr)
2021	5,253,267.99	671,556	-	-	5,924,823.99
Total	5,253,267.99	671,556	-	-	5,924,823.99

Estimation of emissions by sources and removals by sinks included

Year of Monitoring/Reporting Period	Emissions from deforestation (tCO ₂ -e/yr)	If applicable, emissions from forest degradation (tCO ₂ -e/yr)*	If applicable, removals by sinks (tCO ₂ -e/yr)	Net emissions and removals (tCO ₂ -e/yr)
2021	11,325,034	877,431	-	12,202,465
Total	11,325,034	877,431	-	12,202,465

Calculation of emission reductions

Total Reference Level emissions during the Reporting Period (tCO₂-e)	5,924,823.99
Net emissions and removals under the ER Program during the Reporting Period (tCO₂-e)	12,202,465
Emission Reductions during the Reporting Period (tCO₂-e)	-6,277,641.01

Monte Carlo analysis

Uncertainties are estimated using a Monte Carlo approach, where model parameters are repeatedly resampled and overall uncertainty estimated from the resulting distribution (Figure 3). This process, like with deforestation, indicates with a high degree of confidence that emissions have increased in 2021 relative to the reference period (Annex 7 - Table 7).

Annex 7 - Table 7: Monte Carlo analysis

Model parameters	Median estimate (tCO ₂)	95% C.I.	Relative margin (%)
Reference period	5,540,513	4,542,735 to 6,615,100	18.7%
Monitoring period	11,489,514	9,402,131 to 13,693,718	18.7%
Emissions reductions	-5,936,199	-7,963,573 to -3,972,248	33.6%

A model sensitivity analysis was conducted using the same procedures as the ER monitoring report, fixing parameters at their nominal values one-at-a-time and reporting the reduction in overall uncertainty (Annex 7 - Table 8). The greatest source of uncertainty is from monitoring period activity data, followed by reference period activity data and above-ground biomass estimates. Notable is that the main sources of uncertainty are identical to those associated with deforestation emissions estimates. Therefore, any efforts to reduce uncertainty in these estimates are likely to also have a positive impact on reducing uncertainty in emissions from degradation.

Annex 7 - Table 8: Sensitivity analysis

Model parameters	Reduction in uncertainty (%)	
	Deforestation only	Deforestation + degradation
Nominal	0	0
Reference area	11.0	8.3
Monitoring area	44.2	34.9
AGB	17.0	5.7
BGB	7.2	1.7
CF	6.4	-0.8